

CONTENTS VOLUME 86 DECEMBER 1986

- page 1 HESLOP-HARRISON, J., HESLOP-HARRISON, Y., CRESTI, M., TIEZZI, A. and CIAMPOLINI, F. Actin during pollen germination
- 9 CURTIS, A. S. G., FORRESTER, J. V. and CLARK, P. Substrate hydroxylation and cell adhesion
- 25 CURTIS, A. S. G. and McMURRAY, H. Conditions for fibroblast adhesion without fibronectin
- 35 UKIL, M., CHATTERJEE, K., DEY, A., GHOSH, S. and MUKHERJEE, A. S. Affinity binding of non-histone chromatin proteins to the X chromosome of *Drosophila* by *in situ* reconstitution and its significance
- 47 ENGSTRÖM, W. Differential effects of epidermal growth factor (EGF) on cell locomotion and cell proliferation in a cloned human embryonal carcinoma-derived cell line *in vitro*
- 57 LELKES, G., FODOR, I., LELKES, G. and HOLLÁN, S. R. The mobility of intramembrane particles in non-haemolysed human erythrocytes: factors affecting acridine-orange-induced particle aggregation
- 69 CLAVIEZ, M., BRINK, M. and GERISCH, G. Cytoskeletons from a mutant of *Dictyostelium discoideum* with flattened cells
- 83 PAYE, M., ETIEVANT, CH., MICHIELS, F., PIERARD, D., NUSGENS, B. and LAPIÈRE, CH. M. Characterization of a spontaneously transformed pulmonary embryonic rat (PER) epithelial cell line
- 95 PAYE, M. and LAPIÈRE, CH. M. The lack of attachment of transformed embryonic lung epithelial cells to collagen I is corrected by fibronectin and FXIII
- 109 NAKATSUJI, N. Presumptive mesoderm cells from *Xenopus laevis* gastrulae attach to and migrate on substrata coated with fibronectin or laminin
- 119 WARBURTON, M. J., FERNS, S. A., KIMBELL, R. and MONAGHAN, P. Dissociation of basement membrane protein deposition and cell spreading in virally transformed rat mammary myoepithelial cells
- 133 STOTT, D. and WYLIE, C. C. Invasive behaviour of mouse primordial germ cells *in vitro*
- 145 ZIPARO, E., ZANI, B. M., FILIPPINI, A., STEFANINI, M. and MARCHESI, V. T. Proteins of the membrane skeleton in rat Sertoli cells
- 155 BEKERS, A. G. M., PIECK, A. C. M., RIJKEN, A. A. M. and WANKA, F. Evidence for the persistence of a decondensed chromosome scaffold in the interphase nucleus
- 173 VERHEIJEN, R., KUIJPERS, H., VOOLJS, P., VAN VENROOIJ, W. and RAMAEKERS, F. Distribution of the 70K U1 RNA-associated protein during interphase and mitosis: correlation with other U RNP particles and proteins of the nuclear matrix
- 191 NOVAK, B. and MITCHISON, J. M. Change in the rate of CO₂ production in synchronous cultures of the fission yeast *Schizosaccharomyces pombe*: a periodic cell cycle event that persists after the DNA-division cycle has been blocked
- 207 CREANOR, J. and MITCHISON, J. M. Nucleoside diphosphokinase, an enzyme with step changes in activity during the cell cycle of the fission yeast *Schizosaccharomyces pombe*. I. Persistence of steps after a block to the DNA-division cycle
- 217 KOCH, G., SMITH, M., MACER, D., WEBSTER, P. and MORTARA, R. Endoplasmic reticulum contains a common, abundant calcium-binding glycoprotein, endoplasmic reticulum chaperone
- 233 ATHERTON, B. T., MEYER, D. M. and SIMPSON, D. G. Assembly and remodelling of myofibrils and intercalated discs in cultured neonatal rat heart cells
- Continued overleaf*

