

## PUBLISHER'S NOTE

# Expression of Concern: Expression of junctional adhesion molecule-A prevents spontaneous and random motility

Gianfranco Bazzoni, Paolo Tonetti, Luca Manzi, Maria R. Cera, Giovanna Balconi and Elisabetta Dejana

This Expression of Concern relates to *J. Cell Sci.* (2005) **118**, 623-632 (doi:10.1242/jcs.01661).

An issue initially raised by a reader and on [PubPeer](#) suggests that there is a duplication of the tubulin blots for samples from 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells in Fig. 4B of this article.

The journal has been unable to get in touch with the authors of this study and have learnt that the first and last authors have recently retired.

We are therefore publishing this Expression of Concern to make readers aware of this issue and our efforts to resolve it.

# Expression of junctional adhesion molecule-A prevents spontaneous and random motility

Gianfranco Bazzoni<sup>1,\*</sup>, Paolo Tonetti<sup>1</sup>, Luca Manzi<sup>1</sup>, Maria R. Cera<sup>2</sup>, Giovanna Balconi<sup>3</sup> and Elisabetta Dejana<sup>2,3,4</sup>

<sup>1</sup>Laboratory of Systems Biology, Department of Immunology and Cell Biology, Istituto di Ricerche Farmacologiche Mario Negri, 20157 Milano, Italy

<sup>2</sup>Department of Vascular Biology, FIRG Institute of Molecular Oncology, 20139 Milano, Italy

<sup>3</sup>Laboratory of Vascular Biology, Department of Immunology and Cell Biology, Istituto di Ricerche Farmacologiche Mario Negri, 20157 Milano, Italy

<sup>4</sup>Department of Biomolecular and Biotechnological Sciences, School of Sciences, University of Milan, 20133 Milano, Italy

\*Author for correspondence (e-mail: bazzoni@marionegri.it)

Accepted 24 November 2004

Journal of Cell Science 118, 623-632 Published by The Company of Biologists 2005

doi:10.1242/jcs.01661

## Summary

Junctional adhesion molecule-A (JAM-A) is a cell-surface glycoprotein that localizes to intercellular junctions and associates with intracellular proteins via PSD95-*Dlg*-ZO1-binding residues. To define the functional consequences of JAM-A expression, we have produced endothelial cells from *JAM-A*-deficient mice. We report here that the absence of JAM-A enhanced spontaneous and random motility. In turn, the enhanced motility of JAM-A-negative cells was abrogated either on transfection of exogenous JAM-A or on treatment with inhibitors of glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ). In addition, in JAM-A-positive cells, motility was enhanced on inactivation of protein kinase C $\zeta$  (PKC $\zeta$ ), which is an inhibitor of GSK-3 $\beta$ . Although these findings suggested that JAM-A might inhibit GSK-3 $\beta$ , we found that expression per se of JAM-A did not change the levels of inactive GSK-3 $\beta$ . Thus, JAM-

A expression may regulate effectors of motility that are also downstream of the PKC $\zeta$ /GSK-3 $\beta$  axis. In support of this view, we found that JAM-A absence increased the number of actin-containing protrusions, reduced the stability of microtubules and impaired the formation of focal adhesions. Notably, all the functional consequences of JAM-A absence were reversed either on treatment with GSK-3 $\beta$  inhibitors or on transfection of full-length JAM-A, but not on transfection of a JAM-A deletion mutant devoid of the PSD95-*Dlg*-ZO1-binding residues. Thus, by regulating cytoskeletal and adhesive structures, JAM-A expression prevents cell motility, probably in a PSD95-*Dlg*-ZO1-dependent manner.

Key words: Junction adhesion molecule, Motility, Glycogen Synthase kinase 3, PDZ domains, Microtubules, Focal adhesions

## Introduction

Junctional adhesion molecule-A (JAM-A), which is also known as JAM, JAM-1 and F11R, belongs to a family of cell adhesion molecules that contain two immunoglobulin-like folds in the extracellular domain (Bazzoni, 2003). JAM-A localizes to intercellular junctions (Martin-Padura et al., 1998) and mediates processes as diverse as junction assembly (Liu et al., 2000; Liang et al., 2000), leukocyte transmigration (Martin-Padura et al., 1998; Del Maschio et al., 1999), platelet activation (Sobocka et al., 2000) and pathogen infection (Barton et al., 2001; Amieva et al., 2003).

JAM-A comprises an extracellular domain, a transmembrane segment and a cytoplasmic tail. The extracellular domain forms parallel dimers (Kostrewa et al., 2001) and binds several ligands, such as JAM-A itself (Liu et al., 2000; Bazzoni et al., 2000a), the leukocyte integrin  $\alpha$ L $\beta$ 2 (Ostermann et al., 2002) and the Reovirus protein  $\sigma$ -1 (Barton et al., 2001). The cytoplasmic tail contains the carboxyl terminal residues F<sup>298</sup>-L<sup>299</sup>-V<sup>300</sup> (FLV<sub>COOH</sub>), which represent a recognition motif for type II PSD95-*Dlg*-ZO-1 (PDZ) domains. JAM-A associates directly with at least five different PDZ proteins, namely ZO-1 (Bazzoni et al., 2000b; Ebnet et al., 2000), AF6/Afadin (Ebnet et al., 2000), CASK/Lin-2 (Martinez-Estrada et al., 2001), Par3/ASIP (Ebnet et al., 2001; Itoh et al., 2001) and

MUPP-1 (Hamazaki et al., 2002), as evaluated by in vitro binding assays. Importantly, all these interactions are abolished on deletion of the FLV<sub>COOH</sub> motif.

Although the junctional localization of most of these PDZ proteins suggests a role in the assembly or stabilization of JAM-A-containing junctions, they may bind other molecules and play additional roles. For example, ZO-1 (Fanning et al., 2002) and AF6/Afadin (Boettner et al., 2000) bind directly F-actin and profilin, respectively. CASK/Lin-2 interacts with protein 4.1, which nucleates actin and spectrin filaments (Cohen et al., 1998). Finally, Par3 associates with Par6 and atypical protein kinase C (PKC), which in turn regulate cell polarity (Ohno, 2001). To date, however, the functional role of JAM-A and its PDZ-dependent associations remains largely unknown.

To further define the function of JAM-A, we have produced and characterized endothelial cells from *JAM-A*-deficient mice. We report here that JAM-A absence enhanced spontaneous and random motility. In turn, the enhanced motility of JAM-A-defective cells was abolished on transfection of full-length JAM-A (JAM-A FL), but not a deletion mutant (JAM-A delFLV) that lacks the FLV<sub>COOH</sub> motif. Thus, JAM-A inhibits cell motility in a PDZ-dependent way. Finally, we report that JAM-A controls cell motility by regulating cytoskeletal and

adhesive structures that are also regulated by the atypical PKC $\zeta$  substrate glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ).

## Materials and Methods

### Cell lines

The 164-JAM-A<sup>+/+</sup> and 160-JAM-A<sup>-/-</sup> cells were obtained from *floxP-JAM-A-floxP* mice and *JAM-A<sup>-/-</sup>* mice (the latter derived from mating *floxP-JAM-A-floxP* with *Cre* transgenic mice), respectively. The *floxP-JAM-A-floxP* and *JAM-A<sup>-/-</sup>* mice have been generated as described previously (Cera et al., 2004). To prepare the cells, 3-month old mice were killed by cervical dislocation and exsanguination. Then, lungs were explanted and digested with collagenase (Sigma, St Louis, MO). Endothelial cells were immortalized by polyoma middle-T antigen to establish the 164-JAM-A<sup>+/+</sup> and 160-JAM-A<sup>-/-</sup> cells. Finally, the 160-JAM-A FL and 160-JAM-A delFLV cells were produced by transfection of recombinant constructs (Bazzoni et al., 2000b). Cells were kept in culture (with 1:2 splitting every second day) in D-MEM supplemented with 10% (vol:vol) FBS (Gibco Invitrogen, Carlsbad, CA), 10  $\mu$ g/ml heparin (Sigma) and 5  $\mu$ g/ml endothelial cell growth supplement.

### Antibodies and reagents

The following antibodies were used: rat anti-JAM-A mAbs BV12, BV11, and BV19; rat anti-PECAM-1 MEC 13.3 (kindly provided by A. Vecchi, Mario Negri Institute, Milan); mouse anti-vinculin mAb hVIN-1 (Sigma); mouse anti-paxillin mAb 349 (Transduction Laboratories, Belgium); mouse anti- $\alpha$ -tubulin mAb B-5-1-2 and anti-acetylated- $\alpha$ -tubulin mAb 6-11B-1 (Sigma); mouse anti-GSK3 $\beta$  mAb 7 (Transduction). Rabbit GSK-3 $\beta$  Ser<sup>9</sup>-P antiserum (Biosource International, Camarillo, CA); rabbit PKC $\zeta$  antiserum sc-216 (Santa Cruz Biotechnology, Santa Cruz, CA); rabbit ZO-1 and occludin antisera (Zymed Laboratories, San Francisco, CA). TRITC- and FITC-labeled anti-IgG (H+L) antibodies were from Jackson ImmunoResearch Laboratories (West Grove, PA). FITC-Phalloidin, colloidal gold, lithium chloride, SB216763, taxol and nocodazole were from Sigma. Myristoylated PKC $\zeta$  pseudo-substrate (PKC $\zeta$ -PS) was from Biosource International.

### Fluorescence flow cytometric analysis

Fluorescence flow cytometric analysis was performed with a FACStar Plus apparatus (Becton Dickinson, Mountainview, CA). Cells were detached by incubation with 0.25% Trypsin (Gibco) in phosphate-buffered saline. Aliquots (100  $\mu$ l) of cell suspensions (10<sup>6</sup> cells/ml) were incubated with 2% (wt:vol) BSA in PBS (for 60 minutes on ice), then washed with PBS plus 0.2% BSA, and incubated with primary antibodies (for 45 minutes on ice) at a concentration of 20  $\mu$ g/ml. Cells were then washed three times, incubated (for 45 minutes on ice) with secondary fluorescein isothiocyanate-conjugated goat F(ab<sup>2</sup>) anti-rat IgG (Caltag Laboratories, Burlingame, CA) and washed three times before analysis.

### Immunofluorescence microscopy

Single-cell suspensions were seeded at different densities onto glass coverslips (that had been coated with human plasma fibronectin; 7  $\mu$ g/ml) and incubated for 120 minutes at 37°C. Cells were then fixed with 3% (wt:vol) paraformaldehyde for 15 minutes, permeabilized with 0.5% (vol:vol) Triton-X-100 for 3 minutes, and stained with primary and TRITC-labeled secondary antibodies, as previously described (Martinez-Estrada et al., 2001). Coverslips were mounted in 488-Mowiol and analyzed with a Zeiss Axiophot microscope (equipped with 100 $\times$  Plan-Neofluar Ph3 objective lens). Images were recorded with a Zeiss 456070 camera on 3200 ASA Kodak films and processed using Adobe Photoshop.

### Phagokinetic track assay

The assay was performed as described (Albrecht-Buhler, 1977). Briefly, cells were plated onto glass coverslips (previously coated with colloidal gold) and incubated for 24 hours at 37°C. Then, cells were fixed with 3% paraformaldehyde. Pictures of individual cells were analyzed with the NIH Image<sup>TM</sup> software for measuring particle-free area and number of tracks per cell. For these measurements, only particle-free surfaces containing one cell were analyzed, in order to exclude surfaces produced by either intersection of two (or more) cells or cells derived from cell division during the 24 hours.

### Microtubule polymerization assay

The assay was performed as described previously (Minotti et al., 1991). Briefly, cells were grown in a 6-well plate. Before reaching confluence, cells were lysed with taxol-containing lysis buffer (20 mM Tris-HCl, pH 6.8; 140 mM NaCl; 0.5% NP-40; 1 mM MgCl<sub>2</sub>; 2 mM EGTA; 4  $\mu$ g/ml taxol). Wells were scraped and washed with lysis buffer, while any remaining insoluble residue was solubilized with Laemmli sample buffer. Then, the aliquot of taxol-containing lysis buffer was centrifuged, and the pellet was combined with Laemmli-solubilized material to obtain the NP40-insoluble fraction. Samples were separated by SDS-PAGE. Filters were analyzed by western blotting with anti-tubulin antibody, secondary HRP-conjugated antimurine IgG antibody and chemiluminescence.


### Wound assay

Cells were seeded (at a density of 10<sup>5</sup>/cm<sup>2</sup>) onto gelatin-coated 24-well plates and grown to confluence. Then, the culture medium was removed, and scratch wounds were produced using a plastic tip for 1000  $\mu$ l pipettes. After two washes with D-PBS, cells were incubated with D-MEM medium containing 10% serum. Finally, cells were fixed and stained (5 minutes at room temperature) with a solution of crystal violet in methanol/water (80:20; vol:vol). The migrated distance was obtained by subtracting the width of the wound at 24 hours from the width at time 0 (i.e. immediately after wounding) and by averaging the results obtained at the different 1 mm intervals. At time 0, wounds measured 1118 $\pm$ 36 (164-JAM-A<sup>+/+</sup>), 1110 $\pm$ 25 (160-JAM-A<sup>-/-</sup>), 1204 $\pm$ 45 (160-JAM-A FL) and 1108 $\pm$ 63  $\mu$ m (160-JAM-A delFLV; mean $\pm$ s.e.m., *n*=15).

## Results

### JAM-A expression in endothelial cells

We have produced JAM-A-positive (164-JAM-A<sup>+/+</sup>) and JAM-A-negative (160-JAM-A<sup>-/-</sup>) endothelial cells from wild-type and *JAM-A*-deficient mice, respectively (Cera et al., 2004). In addition, we have transfected the 160-JAM-A<sup>-/-</sup> cells with either full-length JAM-A (160-JAM-A FL) or a deletion mutant lacking the FLV<sub>COOH</sub> motif (160-JAM-A delFLV). As assessed by fluorescence flow cytometric analysis, endogenous JAM-A was expressed in 164-JAM-A<sup>+/+</sup> but not in 160-JAM-A<sup>-/-</sup> cells. Also, exogenous JAM-A FL and delFLV were expressed at similar levels in 160-JAM-A FL and 160-JAM-A delFLV cells, respectively (Fig. 1). By comparison, the expression levels of PECAM-1 and occludin (two other molecules that localize at intercellular contacts in endothelial cells) did not differ among the four cell lines, as assessed by fluorescence flow cytometric analysis. Specifically, in 164-JAM-A<sup>+/+</sup>, 160-JAM-A<sup>-/-</sup>, 160-JAM-A FL and 160-JAM-A delFLV cells, the values of mean fluorescence intensity for PECAM-1 were 43.3, 44.4, 47.6 and 40.5 units, respectively. Also, the values for occludin were 96.3, 106.8, 104.9 and 114.5 units, respectively.



**Fig. 1.** JAM-A expression. Expression of JAM-A was evaluated by FACS analysis using anti-JAM-A mAb BV12 (shaded area). Other anti-JAM-A antibodies (mAbs BV19 and BV11) gave similar results. The white area indicates negative control antibodies. Results are expressed as arbitrary units of fluorescence (on the x-axis) and as cell count (on the y-axis).


As assessed by IF microscopy, JAM-A localized to intercellular junctions in all cell lines, with the exception of 160-JAM-A<sup>-/-</sup>. Nonetheless, all cell lines displayed similar expression of junctional markers and similar levels of paracellular permeability (data not shown), thus indicating that JAM-A is dispensable for junction formation and function in contacting cells. Next, in search for additional functions of JAM-A, we analyzed noncontacting cells.

#### JAM-A expression reduces the number of membrane protrusions

When plated onto fibronectin, the four cell lines formed actin-containing protrusions of variable length and shape (Fig. 2A). However, 160-JAM-A<sup>-/-</sup> cells formed a higher number of long protrusions than 164-JAM-A<sup>+/+</sup> cells. Also, on transfection of 160-JAM-A<sup>-/-</sup> cells with JAM-A FL (but not JAM-A delFLV), the number of long protrusions decreased to the level observed in 164-JAM-A<sup>+/+</sup> cells. Then, cells were analyzed individually for measuring the distribution of this type of protrusion (Fig. 2B). Specifically, we counted protrusions that were longer than 2 μm. The average number of long protrusions per cell was about 1.5-fold higher in 160-JAM-A<sup>-/-</sup> (8.3±0.4 protrusions; mean±s.e.m.; n=168 cells) and 160-JAM-A delFLV (8.6±0.3; n=123), compared with 164-JAM-A<sup>+/+</sup> (5.5±0.3; n=176) and 160-JAM-A FL cells (5.9±0.3; n=129). Thus, expression of JAM-A (but not JAM-A delFLV) reduced the protrusive activity at the membrane. As actin-based protrusions are associated with cell motility (Condeelis, 1993), we next evaluated the effect of JAM-A expression on motility.

#### JAM-A expression prevents spontaneous and random motility

To study motility, we deployed the phagokinetic track assay (Albrecht-Buehler, 1977). Over a 24-hour interval, most 164-JAM-A<sup>+/+</sup> cells produced a small particle-free surface, with no



**Fig. 2.** JAM-A expression reduces the number of membrane protrusions. (A) Cells were plated onto fibronectin-coated coverslips (for 2 hours), stained with FITC-phalloidin and examined by IF microscopy. (B) Individual cells were analyzed for measuring the distribution of the number of long protrusions (i.e. length >2 μm) per cell. Results are derived from three experiments.

(or very few) tracks (Fig. 3A). By contrast, most 160-JAM-A<sup>-/-</sup> cells produced a broader particle-free surface, with many more tracks. As above, on transfection of 160-JAM-A<sup>-/-</sup> cells with JAM-A FL (but not JAM-A delFLV), the phagokinesis pattern became similar to the one observed in 164-JAM-A<sup>+/+</sup> cells.

Then, cells were analyzed individually for measuring the distribution of particle-free area (Fig. 3B) and number of tracks (Fig. 3C). The average particle-free area per cell was greater in 160-JAM-A<sup>-/-</sup> (522±32 μm<sup>2</sup>; mean±s.e.m.; n=61 cells) and 160-JAM-A delFLV cells (530±34 μm<sup>2</sup>; n=58), compared with 164-JAM-A<sup>+/+</sup> (239±14 μm<sup>2</sup>; n=72) and 160-JAM-A FL cells (328±19 μm<sup>2</sup>; n=53). Similarly, the average number of tracks per cell was higher in 160-JAM-A<sup>-/-</sup> (2.9±0.2; mean±s.e.m.; n=181) and 160-JAM-A delFLV cells (3.3±0.2; n=182), compared with 164-JAM-A<sup>+/+</sup> (0.2±0.1; n=180) and 160-JAM-A FL cells (1.4±0.1; n=182).

#### Inactivation of GSK-3β mimics JAM-A-dependent inhibition of motility

Similarly to the expression of JAM-A in our system, inactivation of GSK-3β was shown to reduce long protrusions in keratinocytes (Koivisto et al., 2003). Thus, we surmised that JAM-A expression might inhibit protrusions (and possibly motility) by inactivating GSK-3β. To address this issue, we first treated cells with chemical inhibitors of GSK-3β. In 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells, either lithium (40 mM;

Table 1A) or SB216763 (20  $\mu$ M; Table 1B) reduced the number of protrusions, the particle-free area and the number of tracks per cell almost to the levels observed in 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells. Thus, inactivation of GSK-3 $\beta$  mimicked the inhibitory effect of JAM-A expression on protrusions and motility.

PKC $\zeta$  is one of the endogenous regulators of GSK-3 $\beta$ . Notably, PKC $\zeta$ -dependent inactivation of GSK-3 $\beta$  (via phosphorylation on Ser<sup>9</sup>) is instrumental in controlling cell migration (Etienne-Manneville and Hall, 2003). To test the role of PKC $\zeta$  in our system, we treated cells with a cell-permeable and PKC $\zeta$ -specific pseudo-substrate (PKC $\zeta$ -PS). In 160-JAM-A FL cells, PKC $\zeta$ -PS (20  $\mu$ M; Table 1C) increased the number of protrusions, the particle-free area and the number of tracks per cell to the levels observed in 160-JAM-A<sup>-/-</sup> cells. Thus, complementing the GSK-3 $\beta$  inhibitor data, PKC $\zeta$ -PS prevented the inhibitory effect of JAM-A expression on protrusions and motility. The effect of PKC $\zeta$ -PS was probably due to GSK-3 $\beta$  activation, as an inhibitor of GSK-3 $\beta$  could prevent it. For instance, in PKC $\zeta$ -PS-treated 160-JAM-A FL cells, SB216763 (5  $\mu$ M) strongly reduced the effect of PKC $\zeta$ -PS (20  $\mu$ M) on the particle-free area (610 $\pm$ 64  $\mu$ m<sup>2</sup> and 312 $\pm$ 56  $\mu$ m<sup>2</sup>, in the absence and presence of SB216763, respectively).


### JAM-A expression does not influence the activation state of GSK-3 $\beta$

To examine the direct effect of the inhibitors on GSK-3 $\beta$ , we analyzed inactive GSK-3 $\beta$  (GSK-3 $\beta$  Ser<sup>9</sup>-P) by western blotting. In 160-JAM-A FL cells, constitutive levels of GSK-3 $\beta$  Ser<sup>9</sup>-P were detectable even in absence of the inhibitors. Either lithium (40 mM) or SB216763 (20  $\mu$ M) increased the levels of GSK-3 $\beta$  Ser<sup>9</sup>-P, while PKC $\zeta$ -PS (20  $\mu$ M) reduced them. In all conditions tested, no changes in total GSK-3 $\beta$  were observed (data not shown).

Then, to test whether JAM-A expression per se influenced the activation state of GSK-3 $\beta$ , we compared the constitutive levels of GSK-3 $\beta$  Ser<sup>9</sup>-P among the cell lines. We found similar levels of GSK-3 $\beta$  Ser<sup>9</sup>-P in 160-JAM-A<sup>-/-</sup>, FL and delFLV cells (data not shown). Thus, although JAM-A expression (see Figs 2 and 3) and GSK-3 $\beta$  inhibitors alike (see Table 1) reduced protrusions and motility, JAM-A expression did not affect the activation state of GSK-3 $\beta$ . Rather, we hypothesized that JAM-A expression regulated motility-related responses downstream of the PKC $\zeta$ -GSK-3 $\beta$  axis.

### JAM-A expression stabilizes microtubules

As shown above, one of these responses is the formation of actin-



**Fig. 3.** JAM-A expression prevents cell motility. (A) Cells were plated onto gold particles, incubated for 24 hours and fixed. Individual cells were analyzed for measuring the distribution of (B) particle-free area and (C) number of tracks per cell. Results are derived from three experiments.

based protrusions, which are reduced in number on either expression of JAM-A (Fig. 2) or treatment with GSK-3 $\beta$  inhibitors (Table 1). Stabilization of microtubules is another effect of GSK-3 $\beta$  inactivation that is important for cell motility (Zumbrunn et al., 2001). To test whether JAM-A affected microtubule dynamics, we first studied microtubule assembly (by IF microscopy) and tubulin polymerization (by western blotting) following nocodazole withdrawal. By IF microscopy (Fig. 4A), microtubules disappeared from all cell lines after 16-hour treatment with 10  $\mu$ M nocodazole. However, by 2 hours after nocodazole withdrawal, microtubules were more numerous in 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells than in 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells. In addition, by western blotting (Fig. 4B), NP40-insoluble (i.e. polymerized) tubulin was strongly reduced in all cell lines after nocodazole treatment. However, after nocodazole withdrawal, much more insoluble tubulin was detectable in 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells than in 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells. As a loading control, similar amounts of NP40-insoluble ZO-1 were detectable in the four cell lines on nocodazole withdrawal. On the basis of these results, we next investigated whether JAM-A expression might actually stabilize microtubules.

Cold temperature causes microtubule disassembly (Bershadsky et al., 1979), even though diverse cell types differ

**Table 1. PKC- $\zeta$ -dependent inactivation of GSK-3 $\beta$  mimics JAM-A-dependent inhibition of protrusions and motility**

**A. Lithium**

| Treatment and cell line  | Number of protrusions/cell |         | Particle-free area/cell ( $\mu\text{m}^2$ ) |         | Number of tracks/cell |         |
|--------------------------|----------------------------|---------|---------------------------------------------|---------|-----------------------|---------|
|                          | Control                    | Lithium | Control                                     | Lithium | Control               | Lithium |
| 164-JAM-A <sup>+/+</sup> | 5.3±0.6                    | 5.7±0.6 | 324±53                                      | 332±23  | 1.2±0.3               | 1.5±0.4 |
| 160-JAM-A <sup>-/-</sup> | 9.7±0.6                    | 6.7±0.5 | 818±103                                     | 354±17  | 4.8±0.4               | 2.3±0.3 |
| 160-JAM-A FL             | 4.7±0.4                    | 4.9±0.5 | 312±64                                      | 306±35  | 1.9±0.3               | 1.9±0.3 |
| 160-JAM-A delFLV         | 9.0±0.5                    | 6.2±0.5 | 1027±128                                    | 337±33  | 4.1±0.4               | 2.3±0.3 |

**B. SB216763**

| Treatment and cell line  | Number of protrusions/cell |          | Particle-free area/cell ( $\mu\text{m}^2$ ) |          | Number of tracks/cell |          |
|--------------------------|----------------------------|----------|---------------------------------------------|----------|-----------------------|----------|
|                          | Control                    | SB216763 | Control                                     | SB216763 | Control               | SB216763 |
| 164-JAM-A <sup>+/+</sup> | n.d.                       | n.d.     | 319±20                                      | 325±22   | 1.4±0.3               | 1.4±0.3  |
| 160-JAM-A <sup>-/-</sup> | 8.6±0.8                    | 4.8±1.2  | 884±94                                      | 308±18   | 3.8±0.4               | 1.5±0.4  |
| 160-JAM-A FL             | 4.5±0.6                    | 3.8±0.6  | 324±34                                      | 338±29   | 1.7±0.5               | 1.6±0.4  |
| 160-JAM-A delFLV         | 8.9±0.8                    | 3.9±0.8  | 934±111                                     | 317±21   | 3.7±0.5               | 1.4±0.3  |



**C. PKC $\zeta$ -PS**

| Treatment and cell line  | Number of protrusions/cell |                 | Particle-free area/cell ( $\mu\text{m}^2$ ) |                 | Number of tracks/cell |                 |
|--------------------------|----------------------------|-----------------|---------------------------------------------|-----------------|-----------------------|-----------------|
|                          | Control                    | PKC $\zeta$ -PS | Control                                     | PKC $\zeta$ -PS | Control               | PKC $\zeta$ -PS |
| 164-JAM-A <sup>+/+</sup> | n.d.                       | n.d.            | n.d.                                        | n.d.            | n.d.                  | n.d.            |
| 160-JAM-A <sup>-/-</sup> | 9.3±1.5                    | 9.3±1.3         | 735±73                                      | 622±118         | 3.1±0.4               | 2.8±0.3         |
| 160-JAM-A FL             | 3.3±1.0                    | 9.8±2.0         | 312±28                                      | 575±14          | 1.2±0.3               | 2.6±0.4         |
| 160-JAM-A delFLV         | n.d.                       | n.d.            | 731±136                                     | 715±105         | 3.2±0.5               | 2.9±0.3         |


Cells were incubated in the absence (Control) or presence of lithium (40 mM), SB216763 (20  $\mu\text{M}$ ), and PKC $\zeta$ -PS (20  $\mu\text{M}$ ). Cells were then analyzed for number of actin-based protrusions, particle-free area and number of tracks. Data are expressed as mean±s.e.m. ( $n=30-32$  cells; n.d.=not done).

in the cold-stability of microtubules (Lieuvin et al., 1994). In addition, it is well established that stable microtubules accumulate acetylated tubulin (Westermann and Weber, 2003). So, to analyze microtubule stability, we examined whether JAM-A expression enhanced the amount of either cold-stable microtubules or acetylated tubulin, or both. To this purpose, we incubated cells at either 4°C or 37°C for 30 minutes. Then we

used antibodies against bulk tubulin (Fig. 5A) and acetylated tubulin (Fig. 5B) to examine cold-stability and tubulin acetylation, respectively. At 37°C, similar amounts of microtubules were detectable in both 164-JAM-A<sup>+/+</sup> and 160-JAM-A<sup>-/-</sup> cells (Fig. 5A, left panels). Remarkably, however, more acetylated tubulin was detectable in 164-JAM-A<sup>+/+</sup> than in 160-JAM-A<sup>-/-</sup> cells (Fig. 5B, left panels).



**Fig. 4.** JAM-A expression enhances microtubule polymerization. Cells were incubated for 16 hours in either the absence (Control/C) or the presence (Nocodazole/N) of 10  $\mu\text{M}$  nocodazole. In some samples, nocodazole was removed after the 16 hours incubation, and cells incubated for an additional 2 hours (Nocodazole withdrawal/NW). Using the anti- $\alpha$ -tubulin mAb B-5-1-2, microtubules were analyzed by IF microscopy (A), and NP40-insoluble tubulin was analyzed by western blotting (B). In (B), ZO-1 is shown as a loading control for NP40-insoluble material upon nocodazole withdrawal.



**Fig. 5.** JAM-A expression stabilizes microtubules. Cells were plated onto fibronectin-coated coverslips and incubated for 90 minutes at 37°C. Then, after the addition of 10 nM taxol (where indicated), cells were further incubated for 30 minutes at 37°C. Finally, cells were incubated at either 37°C or 4°C for an additional 30 minutes, stained with either anti- $\alpha$ -tubulin mAb B-5-1-2 (A) or anti-acetylated- $\alpha$ -tubulin mAb 6-11B-1 (B).

Next we examined the effect of cold temperature. Following exposure to 4°C, the amount of microtubules was only slightly reduced in 164-JAM-A<sup>+/+</sup> cells, whereas it was reduced to a greater extent in 160-JAM-A<sup>-/-</sup> cells (Fig. 5A, middle panels). In addition, in 164-JAM-A<sup>+/+</sup> cells, greater amounts of acetylated tubulin were detectable at 37°C than at 4°C. At variance, in 160-JAM-A<sup>-/-</sup> cells, acetylated tubulin was barely detectable at both 37°C and 4°C (Fig. 5B, left and middle panels). As a control, pretreatment with 10 nM taxol (for 30 minutes before exposure to cold) preserved microtubules and acetylated tubulin in both cell types (right panels). Finally, we found that 160-JAM-A FL cells displayed higher amounts of cold-resistant and acetylated tubulin than 160-JAM-A delFLV cells (not shown). Thus, expression of JAM-A (but not JAM-A delFLV) enhanced microtubule stability.

#### JAM-A expression increases the number of focal adhesions

As highly dynamic microtubules cause disassembly of focal adhesions (Small and Kaverina, 2003), we examined whether JAM-A expression affected focal adhesions as well. By IF

microscopy of vinculin- (Fig. 6A) and paxillin- (Fig. 6B) stained cells, we found that, in 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells, adhesions were fewer in number and more peripheral in location, compared with 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells. Cells were analyzed individually for measuring the distribution of the number of vinculin-based focal adhesions (Fig. 6C). The average number of focal adhesions per cell was greater in 164-JAM-A<sup>+/+</sup> (141±9; mean±s.e.m.; *n*=15 cells) and 160-JAM-A FL cells (121±6), compared with 160-JAM-A<sup>-/-</sup> (53±4) and 160-JAM-A delFLV cells (52±4).

In 160-JAM-A<sup>-/-</sup> cells, 10 nM taxol (a dose that stabilizes microtubules; see above) (Jordan et al., 1993) induced numerous adhesions. Similarly, lithium (40 mM) increased the number of adhesions in 160-JAM-A<sup>-/-</sup> cells, whereas PKC $\zeta$ -PS reduced it in 160-JAM-A FL cells (Fig. 6D). Hence, the effect of JAM-A expression on focal adhesions (Fig. 6) was related to the effect on microtubules (Figs 4 and 5), probably downstream of the PKC $\zeta$ /GSK-3 $\beta$  axis.


#### JAM-A expression causes morphological changes in migrating cells at a wound edge

Next, we analyzed cell migration into a wounded monolayer. Using this system, we found that all cell lines migrated with similar efficiency. For instance, at 24 hours after wounding, the migrated distances were 767±43  $\mu$ m (164-JAM-A<sup>+/+</sup>), 808±29  $\mu$ m (160-JAM-A<sup>-/-</sup>), 807±35  $\mu$ m (160-JAM-A FL) and 751±35  $\mu$ m (160-JAM-A delFLV; mean±s.e.m., *n*=15). Hence, expression of JAM-A (but not JAM-A delFLV) reduced spontaneous motility (see particle-free area) and randomness of motion (see number of tracks), while leaving migration efficiency (see migrated distances in the wound) unaffected.

However, morphological differences among the cell lines were detectable at the wound edge (Fig. 7A). First, compared with 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells, 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells aligned cell body (first row) and actin filaments (second row) in a much more orderly fashion and more in parallel to the wound. Second, vinculin-containing focal adhesions were more numerous and central in 164-JAM-A<sup>+/+</sup> and 160-JAM-A FL cells than in 160-JAM-A<sup>-/-</sup> and 160-JAM-A delFLV cells (third row). Third, the percentage of cells orienting flat protrusions in parallel to the major axis of the wound was about 2.6-fold greater in 164-JAM-A<sup>+/+</sup> cells (41.2±1.4%) than in 160-JAM-A<sup>-/-</sup> cells (15.7±1.0%; mean±s.e.m.; *n*=900 cells). Thus, JAM-A absence induced morphological changes in cell motility not only in noncontacting cells (Figs 2 and 6), but also in cohesive cell sheets (Fig. 7).

#### JAM-A effect is independent of cell-cell contact and expression levels

To evaluate this issue in further detail, we examined membrane protrusions and focal adhesions in small groups of contacting cells (two or three cells). We found that, similarly to single noncontacting cells (Figs 2 and 6), contacting 164-JAM-A<sup>+/+</sup> cells formed fewer actin-based protrusions at the membrane (Fig. 7B, left panels) and more vinculin-based focal adhesions (right panels) compared with 160-JAM-A<sup>-/-</sup> cells. Thus, the



**Fig. 6.** JAM-A expression increases the number of focal adhesions. Cells were plated onto fibronectin, incubated for 2 hours, and then stained with either anti-vinculin mAb hVIN-1 (A) or anti-paxillin mAb 349 (B). (C) Individual cells were analyzed for measuring the distribution of the number of vinculin-based focal adhesions per cell; results are derived from three experiments. In (D), cells were plated onto fibronectin for 30 minutes, incubated for an additional 90 minutes in either the absence (Control) or presence of 10 nM taxol, 40 mM lithium and 20  $\mu$ M PKC $\zeta$ -PS, and finally stained with anti-vinculin mAb hVIN-1.

morphological changes associated with JAM-A expression were not influenced by cell-cell contact.

Finally, we examined whether the functional consequences of JAM-A expression were influenced by the levels of JAM-A expression at the cell surface. To this purpose, we sorted the 160-JAM-A FL cells and obtained the 160-JAM-A FL1 and 160-JAM-A FL2 cells, which express JAM-A at medium and high levels, respectively (Table 2). Using these cells (in parallel with 164-JAM-A<sup>+/+</sup> and 160-JAM-A<sup>-/-</sup> cells for comparison), we found that expression of JAM-A at either medium (FL1) or

high (FL2) levels similarly decreased the migrated area, reduced the number of membrane protrusions and increased the number of focal adhesions. Thus, JAM-A reversed the functional consequences of JAM-A absence independently of its levels of expression.

## Discussion

The major findings of this manuscript are as follows. First, the absence of JAM-A enhances cell motility, increases membrane

**Table 2. JAM-A effect is not dependent on the expression levels at the cell surface**

| Cell line                | JAM-A expression (MFI) | Number of protrusions/cell | Particle-free area/cell ( $\mu\text{m}^2$ ) | Number of adhesions/cell |
|--------------------------|------------------------|----------------------------|---------------------------------------------|--------------------------|
| 164-JAM-A <sup>+/+</sup> | 54                     | 4.5 $\pm$ 0.2              | 258 $\pm$ 24                                | 116 $\pm$ 5              |
| 160-JAM-A <sup>-/-</sup> | 0                      | 9.5 $\pm$ 0.3              | 1021 $\pm$ 162                              | 63 $\pm$ 4               |
| 160-JAM-A FL1            | 70                     | 4.3 $\pm$ 0.2              | 269 $\pm$ 36                                | 120 $\pm$ 4              |
| 160-JAM-A FL2            | 850                    | 4.0 $\pm$ 0.2              | 403 $\pm$ 60                                | 118 $\pm$ 7              |

The 160-JAM-A FL cells were sorted by fluorescence flow cytometry to obtain the 160-JAM-A FL1 and FL2 cells, which express JAM-A at medium and high levels, respectively. Cells were then analyzed for JAM-A expression (by flow cytometry), number of actin-based membrane protrusions, gold particle-free area and number of vinculin-based focal adhesions. Data are expressed as mean $\pm$ s.e.m. ( $n=15,000$  for JAM-A expression;  $n=30$  for protrusion number;  $n=9$  for particle-free area and adhesion number).



protrusions, affects microtubule stability, and reduces focal adhesions. Second, the consequences of JAM-A absence are reversed either on transfection of JAM-A (in a way that critically requires the PDZ-binding motif) or on treatment with GSK-3 $\beta$  inhibitors.

### JAM-A and motility


Although JAM-A expression modulates pleiotropic targets as diverse as protrusions, adhesions and microtubules, the final outcomes are all consistent with a coordinated inhibition of spontaneous and random motility. First, the formation of a single lamellipodial protrusion at the cell front is one of the earliest steps of cell migration that ensures regular and directional motility (Lauffenburger and Horwitz, 1996). At variance, the formation of numerous protrusions (such as those that appear around the periphery of JAM-A-defective cells) is predictive of irregular patterns of motility (as observed in these

cells). In this context, transfection of JAM-A not only restricts overall motility (as reflected in the smaller particle-free area) but also prevents frequent changes in direction (as reflected in the lower number of tracks). Second, focal adhesions are another key factor of motility (Webb et al., 2002). As shown in other systems, numerous adhesions strengthen the restraint to free motion that the substrate exerts on the cells (Ilic et al., 1995). Thus, the higher number of focal adhesions in JAM-A-expressing cells is consistent with the lower motility of these cells. In partial agreement with this view, other authors have reported that JAM-A blockade reduced adhesion to vitronectin (Naik et al., 2003). Third, microtubules play a role in cell motility. Notably, focal adhesions are among the key effectors of motility that are also potential targets for microtubules. In particular, microtubule instability (with microtubules that repeatedly direct to and retract from the adhesions) has been associated with adhesion disassembly (Kaverina et al., 1999; Small and Kaverina, 2003). Conversely, taxol, besides stabilizing microtubules, reduces cell motility (Liao et al., 1995). Hence, the ability of JAM-A to stabilize microtubules provides a probable explanation for the higher number of adhesions and the lower motility of JAM-A-expressing cells (even though we cannot exclude a direct effect of JAM-A on the adhesions). The observation that taxol fully reverses the defective formation of focal adhesions in JAM-A<sup>-/-</sup> cells further supports the notion of a JAM-A-dependent and coordinated regulation of microtubules and adhesions in the inhibition of cell motility.

Finally, JAM-A expression (besides favoring microtubule assembly on nocodazole withdrawal) was associated with enhanced amounts of acetylated tubulin, which is a known marker of microtubule stability (Westermann and Weber, 2003). To further strengthen the link between microtubule stability and cell motility, it is worth mentioning that reduced tubulin acetylation (in cells that overexpress histone deacetylase-6) was associated with enhanced cell motility (Hubbert et al., 2002), even though the precise molecular mechanism remains unclear (Palazzo et al., 2003).

### JAM-A and the PKC $\zeta$ /GSK-3 $\beta$ axis

In addition to its key role in metabolism and proliferation (Manoukian and Woodgett, 2002), GSK-3 $\beta$  regulates protrusions, motility and microtubules. First, GSK-3 $\beta$  inhibitors prevent the formation of long lamellipodia in keratinocytes (Koivisto et al., 2003) and filopodia in neurons (Owen and Gordon-Weeks, 1999). In addition, inactive GSK-3 $\beta$  localizes to the leading edge of growth cones (Eickholt et al., 2002). Second, following PKC $\zeta$ -dependent phosphorylation, inactive GSK-3 $\beta$  Ser<sup>9</sup>-P specifically localizes to the leading edge of migrating astrocytes (Etienne-Manneville and Hall, 2003). Third, GSK-3 $\beta$  inactivation



**Fig. 7.** JAM-A expression causes morphological changes in directionally migrating cells and in cell groups. In (A), confluent cell monolayers were scratch-wounded and incubated for an additional 24 hours. In (B), cells were seeded at high density and incubated for 2 hours to allow the formation of small groups of contacting cells. Finally, cells were stained with crystal violet (Phase contrast), FITC-phalloidin (F-actin), and anti-vinculin mAb hVIN-1 (Vinculin).

stabilizes microtubules by preventing the phosphorylation of adenomatous polyposis coli (Zumbrunn et al., 1999) and microtubule-associated protein 1B (Goold et al., 1999). Building on this evidence, our data (with chemical inhibitors) further show that GSK-3 $\beta$  inhibition reverses the protrusive and motile activity of JAM-A-negative cells, thus establishing a link between JAM-A, the PKC $\zeta$ /GSK-3 $\beta$  axis and motility.

All the herein-described effects of JAM-A expression required the PDZ-binding residues FLV<sub>COOH</sub>, thus suggesting the involvement of PDZ partner(s) of JAM-A. Among these molecules, it is known that Par3 associates with the Par6-PKC $\zeta$  complex (Lin et al., 2000; Joberty et al., 2000), thus representing a putative molecular linkage between JAM-A and GSK-3 $\beta$ . In addition, a JAM-A deletion mutant lacking the PDZ-binding residues redistributes Par3 from cell contacts in the cytosol (Ebnet et al., 2001). Finally, Par3 overexpression causes the loss of contact-dependent inhibition of cell motility (Mishima et al., 2002), and soluble Par3 inhibits PKC $\zeta$  activity (Lin et al., 2000). On the basis of this evidence, we surmised that, in the absence of JAM-A, increased availability of Par3 might result in PKC $\zeta$  inactivation and, consequently, in GSK-3 $\beta$  activation. However, JAM-A expression did not increase the absolute levels of inactive GSK-3 $\beta$  Ser<sup>9</sup>-P (at least within the sensitivity limits of western blotting), thus ruling out a general effect of JAM-A (and Par3) on GSK-3 $\beta$  activation. Interestingly, it has been recently shown that JAM-C (another member of the JAM family) regulates the polarized distribution of a PKC $\lambda$ -containing complex in differentiating spermatids (Gliki et al., 2004). Thus, it remains possible that, by analogy, JAM-A might regulate the polarized distribution of PKC $\zeta$  and/or inactive GSK-3 $\beta$  in our cells (without changing the absolute levels of inactive GSK-3 $\beta$ ).

At first sight, the observation that JAM-A expression did not decrease the levels of GSK-3 $\beta$  activation is at odds with the finding that JAM-A expression and GSK-3 $\beta$  inhibitors both inhibited motility. As a plausible explanation for this apparent paradox, we propose that JAM-A controls a regulator of motility, which is also downstream of GSK-3 $\beta$ . Also, the ability of PKC $\zeta$ -PS to induce motility in JAM-A-expressing cells strengthens the idea that such a putative regulator of motility lies at the crossroads of signals that originate from both JAM-A and the PKC $\zeta$ /GSK-3 $\beta$  axis. However, given the existence of several mechanisms of microtubule stabilization, we cannot rule out at present that JAM-A and GSK-3 $\beta$  have no other target in common, apart from the microtubules.

We also report that JAM-A expression affected protrusions and focal adhesions in both isolated and cohesive cells (at the wound edge). Interestingly, however, JAM-A expression limited the extent of motility (in isolated cells) but not the efficiency of migration (in cohesive cells). Migration efficiency is dependent on instantaneous velocity and directional persistence (Lauffenburger and Horwitz, 1996). Thus, we speculate that intercellular cohesion may enhance either the instantaneous velocity of JAM-A-positive cells or the directional persistence of JAM-A-negative cells, even though real-time measurements are essential to test this hypothesis. Finally, regardless of the cause, the discrepancy between isolated and cohesive cells also suggests that different mechanisms may underlie different types of motility. In this respect, it should be stressed that some of the aforementioned reports analyzed wound edge migration (e.g. Etienne-

Manneville and Hall, 2003) and thus the relevant mechanisms (e.g. localization of inactive GSK-3 $\beta$ ) may not necessarily apply to our experimental system, which uses isolated cells.

With these caveats, however, our findings point to a novel link between JAM-A expression and cell motility. On the basis of these findings, we propose the following model. JAM-A expression favors microtubule stability, probably in a PDZ-dependent manner. As a consequence, more focal adhesions form. In turn, more adhesions may restrain spontaneous cell motility, possibly by providing stronger anchorage to the matrix. Although JAM-A expression does not directly affect GSK-3 $\beta$  activation, inhibitors of GSK3 $\beta$  mimic the effect of JAM-A expression, thus suggesting the existence of an as-yet-unidentified mediator at a crossroad of signals that originate from both JAM-A/PDZ and PKC $\zeta$ /GSK3 $\beta$  axis.

It is also noteworthy that the effect of JAM-A on motility was contact independent, as it was detectable in noncontacting cells. In this respect, the inhibitory mechanism reported here is novel and unrelated to the well established mechanism of contact-dependent inhibition of motility (Abercrombie and Heaysman, 1953).

We thank G. Albrecht-Buehler (Northwestern University, Chicago, IL) for advice on the phagokinetic assay. G. Bazzoni was supported by AICR (Association International for Cancer Research, UK; 04-095), AIRC (Italian Association for Cancer Research) and MIUR (Italian Ministry of University and Research; FIRB RBNE01T8C8\_004, FIRB RBAU01E5F5). E. Dejana was supported by AICR, MIUR (RBNE01F8LT-007, RBNE01MAWA-009); Italian Ministry of Health (Ricerca Finalizzata, 2002; Convenzione 192), Italian NRC (04.00149.ST 97); Fondazione Cariplo (2003.1697/10.6399); ISS (CS36); and European Community (LSHG-CT-2004-503254, 502935, 503573). M.R.C. was supported by a FIRC Fellowship.

## References

- Abercrombie, M. and Heaysman, J.** (1953). Observation on the social behavior of cells in tissue culture. I. Speed of movement of chick heart fibroblasts in relation to their mutual contacts. *Exp. Cell Res.* **5**, 111-131.
- Albrecht-Buehler, G.** (1977). The Phagokinetic tracks of 3T3 cells. *Cell* **11**, 395-404.
- Amieva, M. R., Vogelmann, R., Covacci, A., Tompkins, L. S., Nelson, W. J. and Falkow, S.** (2003). Disruption of the epithelial apical-junctional complex by *Helicobacter pylori* CagA. *Science* **300**, 1430-1434.
- Barton, E. S., Forrest, J. C., Connolly, J. L., Chappell, J. D., Liu, Y., Schnell, F. J., Nusrat, A., Parkos, C. A. and Dermody, T. S.** (2001). Junction adhesion molecule is a receptor for reovirus. *Cell* **104**, 441-451.
- Bazzoni, G.** (2003). The JAM family of junctional adhesion molecules. *Curr. Opin. Cell Biol.* **15**, 525-530.
- Bazzoni, G., Martinez-Estrada, O. M., Mueller, F., Nelboeck, P., Schmid, G., Bartfai, T., Dejana, E. and Brockhaus, M.** (2000a). Homophilic interaction of junctional adhesion molecule. *J. Biol. Chem.* **275**, 30970-30976.
- Bazzoni, G., Martinez-Estrada, O. M., Orsenigo, F., Cordenonsi, M., Citi, S. and Dejana, E.** (2000b). Interaction of junctional adhesion molecule with the tight junction components ZO-1, cingulin, and occludin. *J. Biol. Chem.* **275**, 20520-20526.
- Bershadsky, A. D., Gelfand, V. I., Svitkina, T. M. and Tint, I. S.** (1979). Cold-stable microtubules in the cytoplasm of mouse embryo fibroblasts. *Cell Biol. Int. Rep.* **3**, 45-50.
- Boettner, B., Govek, E. E., Cross, J. and van Aelst, L.** (2000). The junctional multidomain protein AF-6 is a binding partner of the Rap1A GTPase and associates with the actin cytoskeletal regulator profilin. *Proc. Natl. Acad. Sci. USA* **97**, 9064-9069.
- Cera, M. R., del Prete, A., Vecchi, A., Corada, M., Martin-Padura, I., Motoike, T., Tonetti, P., Bazzoni, G., Vermi, W., Gentili, F. et al.** (2004). Increased DC trafficking to lymph nodes and contact hypersensitivity in

- junctional adhesion molecule-A-deficient mice. *J. Clin. Invest.* **114**, 729-738.
- Cohen, A. R., Woods, D. F., Marfatia, S. M., Walther, Z., Chishti, A. H., Anderson, J. M. and Wood, D. F. (1998). Human CASK/LIN-2 binds syndecan-2 and protein 4.1 and localizes to the basolateral membrane of epithelial cells. *J. Cell Biol.* **142**, 129-138.
- Condeelis, J. (1993). Life at the leading edge: the formation of cell protrusions. *Annu. Rev. Cell Biol.* **9**, 411-444.
- Del Maschio, A., de Luigi, A., Martin-Padura, I., Brockhaus, M., Bartfai, T., Fruscella, P., Adorini, L., Martino, G., Furlan, R., de Simoni, M. G. et al. (1999). Leukocyte recruitment in the cerebrospinal fluid of mice with experimental meningitis is inhibited by an antibody to junctional adhesion molecule (JAM). *J. Exp. Med.* **190**, 1351-1356.
- Ebnet, K., Schulz, C. U., Meyer zu Brickwedde, M.-K., Pendl, G. G. and Vestweber, D. (2000). Junctional adhesion molecule interacts with the PDZ domain-containing proteins AF-6 and ZO-1. *J. Biol. Chem.* **275**, 27979-27988.
- Ebnet, K., Suzuki, A., Horikoshi, Y., Hirose, T., Meyer Zu Brickwedde, M. K., Ohno, S. and Vestweber, D. (2001). The cell polarity protein ASIP/PAR-3 directly associates with junctional adhesion molecule (JAM). *EMBO J.* **20**, 3738-3748.
- Eickholt, B. J., Walsh, F. S. and Doherty, P. (2002). An inactive pool of GSK-3 at the leading edge of growth cones is implicated in Semaphorin 3A signaling. *J. Cell Biol.* **157**, 211-217.
- Etienne-Manneville, S. and Hall, A. (2003). Cdc42 regulates GSK-3 $\beta$  and adenomatous polyposis coli to control cell polarity. *Nature* **421**, 753-756.
- Fanning, A. S., Ma, T. Y. and Anderson, J. M. (2002). Isolation and functional characterization of the actin binding region in the tight junction protein ZO-1. *FASEB J.* **16**, 1835-1837.
- Gliki, G., Ebnet, K., Aurrand-Lions, M., Imhof, B. A. and Adams, R. H. (2004). Spermatid differentiation requires the assembly of a cell polarity complex downstream of junctional adhesion molecule-C. *Nature* **431**, 320-324.
- Goold, R. G., Owen, R. and Gordon-Weeks, P. R. (1999). Glycogen synthase kinase 3 $\beta$  phosphorylation of microtubule-associated protein 1B regulates the stability of microtubules in growth cones. *J. Cell Sci.* **112**, 3373-3384.
- Hamazaki, Y., Itoh, M., Sasaki, H., Furuse, M. and Tsukita, S. (2002). Multi-PDZ domain protein 1 (MUPP1) is concentrated at tight junctions through its possible interaction with claudin-1 and junctional adhesion molecule. *J. Biol. Chem.* **277**, 455-461.
- Hubbert, C., Guardiola, A., Shao, R., Kawaguchi, Y., Ito, A., Nixon, A., Yoshida, M., Wang, X. F. and Yao, T. P. (2002). HDAC6 is a microtubule-associated deacetylase. *Nature* **417**, 455-458.
- Ilic, D., Furuta, Y., Kanazawa, S., Takeda, N., Sobue, K., Nakatsuji, N., Nomura, S., Fujimoto, J., Okada, M. and Yamamoto, T. (1995). Reduced cell motility and enhanced focal adhesion contact formation in cells from FAK-deficient mice. *Nature* **377**, 539-544.
- Itoh, M., Sasaki, H., Furuse, M., Ozaki, H., Kita, T. and Tsukita, S. (2001). Junctional adhesion molecule (JAM) binds to PAR-3: a possible mechanism for the recruitment of PAR-3 to tight junctions. *J. Cell Biol.* **154**, 491-497.
- Joberty, G., Petersen, C., Gao, L. and Macara, I. G. (2000). The cell-polarity protein Par6 links Par3 and atypical protein kinase C to Cdc42. *Nat. Cell Biol.* **2**, 531-539.
- Jordan, M. A., Toso, R. J., Thrower, D. and Wilson, L. (1993). Mechanism of mitotic block and inhibition of cell proliferation by taxol at low concentrations. *Proc. Natl. Acad. Sci. USA* **90**, 9552-9556.
- Kaverina, I., Krylyshkina, O. and Small, J. V. (1999). Microtubule targeting of substrate contacts promotes their relaxation and dissociation. *J. Cell Biol.* **146**, 1033-1044.
- Koivisto, L., Alavian, K., Hakkinen, L., Pelech, S., McCulloch, C. A. and Larjava, H. (2003). Glycogen synthase kinase-3 regulates formation of long lamellipodia in human keratinocytes. *J. Cell Sci.* **116**, 3749-3460.
- Kostrewa, D., Brockhaus, M., D'Arcy, A., Dale, G. E., Nelboeck, P., Schmid, G., Mueller, F., Bazzoni, G., Dejana, E., Bartfai, T. et al. (2001). X-ray structure of junctional adhesion molecule: structural basis for homophilic adhesion via a novel dimerization motif. *EMBO J.* **20**, 4391-4398.
- Lauffenburger, D. A. and Horwitz, A. F. (1996). Cell migration: a physically integrated molecular process. *Cell* **84**, 359-369.
- Liang, T. W., DeMarco, R. A., Mrsny, R. J., Gurney, A., Gray, A., Hooley, J., Aaron, H. L., Huang, A., Klassen, T., Tumas, D. B. et al. (2000). Characterization of huJAM: evidence for involvement in cell-cell contact and tight junction regulation. *Am. J. Physiol. Cell. Physiol.* **279**, 1733-1743.
- Liao, G., Nagasaki, T. and Gundersen, G. G. (1995). Low concentrations of nocodazole interfere with fibroblast locomotion without significantly affecting microtubule level: implications for the role of dynamic microtubules in cell locomotion. *J. Cell Sci.* **108**, 3473-3483.
- Lieuvain, A., Labbe, J. C., Doree, M. and Job, D. (1994). Intrinsic microtubule stability in interphase cells. *J. Cell Biol.* **124**, 985-996.
- Lin, D., Edwards, A. S., Fawcett, J. P., Mbamalu, G., Scott, J. D. and Pawson, T. (2000). A mammalian PAR-3-PAR-6 complex implicated in Cdc42/Rac1 and aPKC signalling and cell polarity. *Nat. Cell Biol.* **2**, 540-547.
- Liu, Y., Nusrat, A., Schnell, F. J., Reaves, T. A., Walsh, S., Pochet, M. and Parkos, C. A. (2000). Human junction adhesion molecule regulates tight junction resealing in epithelia. *J. Cell Sci.* **113**, 2363-2374.
- Manoukian, A. S. and Woodgett, J. R. (2002). Role of glycogen synthase kinase-3 in cancer: regulation by Wnts and other signaling pathways. *Adv. Cancer Res.* **84**, 203-229.
- Martin-Padura, I., Lostaglio, S., Schneemann, M., Williams, L., Romano, M., Fruscella, P., Panzeri, C., Stoppacciaro, A., Ruco, L., Villa, A. et al. (1998). Junctional adhesion molecule, a novel member of the immunoglobulin superfamily that distributes at intercellular junctions and modulates monocyte transmigration. *J. Cell Biol.* **142**, 117-127.
- Martinez-Estrada, O. M., Villa, A., Breviario, F., Orsenigo, F., Dejana, E. and Bazzoni, G. (2001). Association of junctional adhesion molecule with calcium/calmodulin-dependent serine protein kinase (CASK/LIN-2) in human epithelial caco-2 cells. *J. Biol. Chem.* **276**, 9291-9296.
- Minotti, A. M., Barlow, S. B. and Cabral, F. (1991). Resistance to antimetabolic drugs in Chinese hamster ovary cells correlates with changes in the level of polymerized tubulin. *J. Biol. Chem.* **266**, 3987-3994.
- Mishima, A., Suzuki, A., Enaka, M., Hirose, T., Mizuno, K., Ohnishi, T., Mohri, H., Ishigatsubo, Y. and Ohno, S. (2002). Over-expression of PAR-3 suppresses contact-mediated inhibition of cell migration in MDCK cells. *Genes Cells* **7**, 581-596.
- Naik, M. U., Vuppalandhi, D. and Naik, U. P. (2003). Essential role of junctional adhesion molecule-1 in basic fibroblast growth factor-induced endothelial cell migration. *Arterioscler. Thromb. Vasc. Biol.* **23**, 2165-2171.
- Ohno, S. (2001). Intercellular junctions and cellular polarity: the PAR-aPKC complex, a conserved core cassette playing fundamental roles in cell polarity. *Curr. Opin. Cell Biol.* **13**, 641-648.
- Ostermann, G., Weber, K. S., Zerneck, A., Schroder, A. and Weber, C. (2002). JAM-1 is a ligand of the beta(2) integrin LFA-1 involved in transendothelial migration of leukocytes. *Nat. Immunol.* **3**, 151-158.
- Owen, R. and Gordon-Weeks, P. R. (2003). Inhibition of glycogen synthase kinase 3 $\beta$  in sensory neurons in culture alters filopodia dynamics and microtubule distribution in growth cones. *Mol. Cell. Neurosci.* **23**, 626-637.
- Palazzo, A., Ackerman, B. and Gundersen, G. G. (2003). Tubulin acetylation and cell motility. *Nature* **421**, 230.
- Small, J. V. and Kaverina, I. (2003). Microtubules meet substrate adhesions to arrange cell polarity. *Curr. Opin. Cell Biol.* **15**, 40-47.
- Soboocka, M. B., Soboocki, T., Banerjee, P., Weiss, C., Rushbrook, J. I., Norin, A. J., Hartwig, J., Salifu, M. O., Markell, M. S., Babinska, A. et al. (2000). Cloning of the human platelet F11 receptor: a cell adhesion molecule member of the immunoglobulin superfamily involved in platelet aggregation. *Blood* **95**, 2600-2609.
- Webb, D. J., Parsons, J. T. and Horwitz, A. F. (2002). Adhesion assembly, disassembly and turnover in migrating cells – over and over and over again. *Nat. Cell Biol.* **4**, E97-E100.
- Westermann, S. and Weber, K. (2003). Post-translational modifications regulate microtubule function. *Nat. Rev. Mol. Cell Biol.* **4**, 938-947.
- Zumbrunn, J., Kinoshita, K., Hyman, A. A. and Nathke, I. S. (2001). Binding of the adenomatous polyposis coli protein to microtubules increases microtubule stability and is regulated by GSK3 beta phosphorylation. *Curr. Biol.* **11**, 44-49.