

New molecular players in capacitative Ca^{2+} entry

James W. Putney, Jr

National Institute of Environmental Health Sciences – NIH, PO Box 12233, Research Triangle Park, NC 27709, USA
e-mail: putney@niehs.nih.gov

Accepted 19 April 2007

Journal of Cell Science 120, 1959–1965 Published by The Company of Biologists 2007
doi:10.1242/jcs.03462

Summary

Capacitative Ca^{2+} entry links the emptying of intracellular Ca^{2+} stores to the activation of store-operated Ca^{2+} channels in the plasma membrane. In the twenty years since the inception of the concept of capacitative Ca^{2+} entry, a number of activation mechanisms have been proposed, and there has been considerable interest in the possibility that TRP channels function as store-operated channels. However, in the past two years, two major players in both the signaling and permeation mechanisms for store-operated channels have been discovered: Stim1 and the

Orai proteins. Stim1 is an endoplasmic reticulum Ca^{2+} sensor. It appears to act by redistributing within a small component of the endoplasmic reticulum, approaching the plasma membrane, but does not seem to translocate into the plasma membrane. Stim1 signals to plasma membrane Orai proteins, which constitute pore-forming subunits of store-operated channels.

Key words: Store-operated Ca^{2+} entry, Ca^{2+} channels, Stim1, Orai1, Ca^{2+} -release-activated Ca^{2+} current

Introduction

In a variety of cell types, activation of phospholipase C by G-protein-coupled receptors results in release of intracellular Ca^{2+} stores and activation of Ca^{2+} entry across the plasma membrane. The intracellular release of Ca^{2+} is most commonly signaled by the second messenger inositol 1,4,5-trisphosphate [$\text{Ins}(1,4,5)\text{P}_3$]. Early work on mechanisms of Ca^{2+} mobilization revealed that Ca^{2+} entry involves signaling from depleted intracellular stores to plasma membrane Ca^{2+} channels, a process referred to as capacitative Ca^{2+} entry or store-operated Ca^{2+} entry. In 2001, my colleagues and I reviewed the state of knowledge about capacitative Ca^{2+} entry for *Journal of Cell Science*, emphasizing theories on the activation mechanism and the nature of the channels (Putney et al., 2001). Embarrassingly, while the phenomenology we summarized at that time was correct, almost all of the favored ideas about activation mechanisms and channels have turned out to be either incorrect or, at the very least, incomplete. In just the past two years, major breakthroughs have revealed key players that initiate signaling from depleted stores and form store-operated channels themselves. In this Commentary, I provide a snapshot of current knowledge about the new players, Stim1 and the Orai, speculate on how they act, and make some guesses as to where this rapidly moving field is headed. I first, however, briefly review the concept of capacitative Ca^{2+} entry and describe some of the pharmacological and biophysical properties that are important for assessing the actions of the newly discovered components.

Capacitative Ca^{2+} entry

Capacitative Ca^{2+} entry, also called store-operated Ca^{2+} entry, simply refers to a phenomenon in which depletion of intracellular Ca^{2+} stores leads to activation of Ca^{2+} -permeable channels in the plasma membrane. The Ca^{2+} entering through the store-operated channels can then be pumped into the stores and thus replenish them. Actions of relatively specific inhibitors of sarcoplasmic-endoplasmic reticulum Ca^{2+}

(SERCA) pumps (discussed below) and clear localization of structures sensitive to the Ca^{2+} -releasing action of $\text{Ins}(1,4,5)\text{P}_3$ indicate that the intracellular store involved is the endoplasmic reticulum, or a component of it. The function of this pathway is, at the very least, to maintain sufficient Ca^{2+} in the endoplasmic reticulum, where it is required for proper protein synthesis and folding (Sambrook, 1990; Teasdale and Jackson, 1996). Indeed, prolonged depletion of endoplasmic Ca^{2+} leads to a stress response and ultimately to apoptosis (Tsukamoto and Kaneko, 1993; Bian et al., 1997). The pathway is also needed to support ligand-activated Ca^{2+} signaling, essential for a wide variety of regulated physiological pathways (Berridge et al., 2000).

Thapsigargin

Thapsigargin is a specific inhibitor of SERCA pumps on the endoplasmic reticulum. Its application to cells results in passive depletion of intracellular Ca^{2+} stores and thereby activation of the store-operated channels (Takemura et al., 1989). The ability to activate Ca^{2+} entry simply by blocking these pumps led to a number of important conclusions. First, entry can be fully activated with no discernible activation of phospholipase C. Second, additivity experiments showed that, at least in the parotid cells initially studied, there is no additional pathway activated by phospholipase-C-linked receptors. Third, because thapsigargin activates Ca^{2+} entry to the same extent that phospholipase C does, but without increasing $\text{Ins}(1,4,5)\text{P}_3$ levels, passage of Ca^{2+} through the endoplasmic reticulum is not required for it to access the cytoplasm (i.e. the permeability of the endoplasmic reticulum membrane to Ca^{2+} does not determine the rate at which Ca^{2+} enters the cytoplasm). This and other arguments favoring a transmembrane flux of Ca^{2+} rather than transit through intracellular stores were summarized in an earlier review (Putney, Jr, 1990).

But perhaps the most significant outcome of the discovery of thapsigargin and other SERCA-inhibiting drugs was that it

provided a functional and pharmacological diagnosis of capacitative entry: a thapsigargin-induced sustained elevation of intracellular Ca^{2+} that is dependent upon extracellular Ca^{2+} is generally attributed to capacitative Ca^{2+} entry – although additional pharmacological criteria may sometimes be considered (Putney, 2001). Notably, the ability of thapsigargin to specifically activate capacitative Ca^{2+} entry, while minimizing the roles of other upstream players in the pathway (receptors, G proteins, phospholipase C, etc.), formed the basis for the high-throughput assays that led to the discoveries of the functions of both Stim and Orai.

I_{crac}

A decade and a half ago, Hoth and Penner discovered and characterized a very small but highly Ca^{2+} -selective current, which they termed I_{crac} for Ca^{2+} -release-activated Ca^{2+} (CRAC) current (Hoth and Penner, 1992; Hoth and Penner, 1993). This current was identified earlier by Lewis and Cahalan although it was not recognized as a store-operated current (Lewis and Cahalan, 1989). I_{crac} shares some properties with previously studied Ca^{2+} -selective voltage-activated currents. For example, I_{crac} demonstrates an anomalous mole-fraction effect^{*}; removing external Ca^{2+} leads to loss of current, unless all external divalent cations are removed by use of chelators, in which case the CRAC channels become permeable to monovalent cations and the current actually becomes larger. However, I_{crac} differs from other known currents in one important way: the single-channel conductance is too small to be measured directly and can only be estimated indirectly by noise analysis (Zweifach and Lewis, 1993). The molecular structure of CRAC channels might therefore be very different from those of other known Ca^{2+} channels (this has turned out to be the case). In addition, I_{crac} shows unusual regulation by Ca^{2+} ; intracellular Ca^{2+} inhibits CRAC channels by multiple mechanisms (Hoth and Penner, 1993; Zweifach and Lewis, 1995), whereas extracellular Ca^{2+} potentiates channel function (Zweifach and Lewis, 1996). These and a number of other findings have provided a rigorous fingerprint for the CRAC channel that eventually aided its correct molecular identification.

Activation mechanisms abound

Numerous reviews have summarized the ideas that have come in and out of vogue in the last twenty years regarding the signaling mechanism for capacitative Ca^{2+} entry (Putney and Bird, 1993; Sage et al., 1994; Berridge, 1995; Favre et al., 1996; Putney, 1997; Parekh and Penner, 1997; Putney and McKay, 1999; Barritt, 1999; Birnbaumer et al., 2000; Elliott, 2001; Gill and Patterson, 2004; Parekh and Putney, 2005; Bolotina and Csutora, 2005; Rosado et al., 2005). Two

fundamental mechanisms have been proposed for transmitting the signal from intracellular stores to the plasma membrane: (1) a diffusible message (Randriamampita and Tsien, 1993; Bolotina and Csutora, 2005) and, (2) conformational coupling of proteins in the intracellular store membranes to plasma membrane channels (Irvine, 1990; Berridge, 1995). It has also been suggested that the activation mechanism might involve the insertion of active channels into the plasma membrane by vesicle secretion (Somasundaram et al., 1995; Yao et al., 1999; Scott et al., 2003). However, this idea does not address the more fundamental question of how the signaling is initiated from Ca^{2+} -depleted stores. This concept is therefore not discussed further here.

Early ideas about capacitative Ca^{2+} entry envisioned specialized structures permitting direct access of external Ca^{2+} to the intracellular stores (Putney, 1977; Casteels and Droogmans, 1981). When it was later realized that Ca^{2+} enters the cell directly across the plasma membrane into the cytoplasm, there was no apparent connection between the stores and the channels, leading to the idea that a diffusible signal might be involved (Putney, 1990). However, there is precedent for direct plasma-membrane–sarcoplasmic-reticulum communication in the case of the very specialized skeletal muscle cells. In the process of excitation-contraction coupling in skeletal muscle, plasma membrane dihydropyridine receptors (voltage-dependent Ca^{2+} channels) interact with, and directly activate, intracellular ryanodine Ca^{2+} release channels. This led Irvine (Irvine, 1990) to propose an analogous conformational coupling model to explain regulated Ca^{2+} entry in cells that utilize the phospholipase-C–Ins(1,4,5) P_3 signaling mechanism: endoplasmic reticulum Ins(1,4,5) P_3 receptors would interact directly with plasma membrane capacitative Ca^{2+} entry channels (see also Berridge, 1995).

In the case of skeletal muscle, information flows from the t-tubule membrane to the sarcoplasmic reticulum; in the case of the Ins(1,4,5) P_3 receptor and capacitative Ca^{2+} entry, a fall in luminal Ca^{2+} levels in the endoplasmic reticulum was proposed to induce a conformational change in the Ins(1,4,5) P_3 receptor, and this would be conveyed directly to the plasma membrane channel by protein-protein interaction. This was an intriguing idea, but there was little direct evidence for it. There is, however, evidence for a requirement for close spatial association between endoplasmic reticulum and plasma membrane store-operated channels. Jaconi et al. used centrifugation to redistribute the organellar contents of oocytes and found that entry only occurred in regions with closely apposed endoplasmic reticulum (Jaconi et al., 1997). Patterson et al. (Patterson et al., 1999) used drugs to stimulate peripheral actin polymerization and disrupted communication between Ca^{2+} stores and plasma membrane store-operated channels (see also Bakowski et al., 2001). Golovina et al. demonstrated that in astrocytes, store-operated Ca^{2+} entry occurred at sites of close plasma membrane–endoplasmic-reticulum apposition (Golovina et al., 2005).

Numerous publications have argued that a diffusible signal couples depletion of intracellular Ca^{2+} stores to Ca^{2+} entry, and some evidence, albeit more controversial, supports the involvement of specific mediators. Two laboratories published findings suggesting cyclic GMP is involved (Pandolfi and Schoeffield-Payne, 1990; Xu et al., 1994); others have

^{*}The anomalous mole fraction effect refers to a phenomenon whereby a mixture of two permeant species produces a current less than that obtained with either component of mixture alone (Almers and McCleskey, 1984). In the case of Ca^{2+} channels, the most striking instance of such behavior is seen with divalent-cation-selective channels that pass essentially no monovalent cations unless the concentration of divalent cations is very low, generally in the low μM range or less. The favored interpretation is that when divalent cations are present, they bind to and reside for relatively long times on a site within the channel known as a selectivity filter (Hess and Tsien, 1984). Ions are thought to permeate such channels in single-file, the rate for a particular species depending both on the permeability of the pore for that species as well as the affinity or off-rate of the selectivity filter. Thus, when Ca^{2+} concentration is reduced below its affinity for the selectivity filter, the site is no longer occupied for the majority of the time, and more permeable Na^+ ions can freely permeate.

implicated arachidonic acid (or one of its metabolites) (Graier et al., 1995; Gailly, 1998; Rzigalinski et al., 1999). However, it now seems likely that these mediators act on channels distinct from the store-operated ones (Thompson, 1997; Mignen et al., 2005). The one candidate for a diffusible signal for store-operated channels that has withstood the test of time is one whose structure is not yet known: a Ca²⁺-entry activator partially purified from store-depleted cells called CIF (for Ca²⁺ influx factor) (Randriamampita and Tsien, 1993; Thomas and Hanley, 1995; Bolotina and Csutora, 2005). A role for such a messenger within the context of the new Stim1 and Orai story is certainly well within the realm of possibility.

The TRP story

Hardie and Minke first pointed out that the *Drosophila* photoreceptor Ca²⁺ channel, TRP, is activated downstream of phospholipase C and thus is a candidate for a store-operated channel (Hardie and Minke, 1992; Hardie and Minke, 1993). When its closest mammalian relatives were cloned (the seven canonical TRPs, TRPCs), a number of laboratories aggressively pursued this possibility. Early results were encouraging (Zhu et al., 1996; Zitt et al., 1996; Kiselyov et al., 1998). However, it soon became apparent that many of the early findings either resulted from misinterpretation of constitutive activity of overexpressed channels or could not be reproduced in other laboratories (Zhu et al., 1998; Trebak et al., 2003). In addition, subsequent work on the *Drosophila* TRP channel demonstrated that this is clearly not a store-operated channel (Acharya et al., 1997). Nonetheless, there are numerous reports of diminished store-operated entry when TRPC expression is suppressed or eliminated (e.g. Mori et al., 2002), and others that are reviewed in Parekh and Putney (Parekh and Putney, 2005). Also, it is clear that, under some circumstances, TRPC channels can exhibit what appears to be store-operated activity when ectopically expressed (Vazquez et al., 2001; Vazquez et al., 2003; Lievremont et al., 2004; Zeng et al., 2004). The major problem is that, when experimentally introduced into cells, TRPC channels do not recapitulate the properties of I_{crac} ; rather, they form non-selective cation channels or, at best, channels that show only modest Ca²⁺ selectivity. More significantly, they have single channel conductances of conventional size (tens of picosiemens). A number of laboratories have demonstrated that TRPC channels can function physiologically as non-store-operated channels activated downstream of phospholipase C, like the original *Drosophila* channel (Putney, 2005; Dietrich et al., 2005). This does not, however, rule out the possibility that TRPCs also function as store-operated channels in different contexts, for example, when assembled in signaling complexes of differing composition. There are a number of examples of store-operated Ca²⁺ entry that have properties distinct from I_{crac} (e.g. Krause et al., 1996; Trepakova et al., 2000), others are reviewed in Parekh and Putney (Parekh and Putney, 2005), and perhaps TRPCs function as store-operated channels in such instances.

Stim1

Almost simultaneously, two laboratories discovered the role of Stim1 (initially *Drosophila* Stim) in capacitative Ca²⁺ entry by use of limited RNAi screens for modulators of thapsigargin-activated Ca²⁺ entry in *Drosophila* S2 cells (Roos et al., 2005)

or in mammalian HeLa cells (Liou et al., 2005). *Drosophila* has a single *Stim* gene, whereas mammals have two, *Stim1* and *Stim2*. Roos et al. found that knocking down Stim1, but not Stim2, reduces store-operated entry and I_{crac} in mammalian cells. However, Liou et al. reported a slight reduction in Ca²⁺ entry in HeLa cells in response to knocking down Stim2 as well. We have found no effect of knockdown of Stim2 in HEK293 cells (Wedel et al., 2007) and no effect of expression of Stim2 in HEK293 cells, even when co-expressed with Orai1 (Mercer et al., 2006). Soboloff et al. reported that Stim2 can act as an inhibitor of store-operated entry (Soboloff et al., 2006a); however, this action is only seen when Stim2 is expressed at very high levels. Thus, the true physiological function of Stim2 requires further investigation.

It seems clear that the function of Stim1 is to act as the initial sensor of Ca²⁺ levels in the endoplasmic reticulum or the component of it involved in regulation of I_{crac} . The Ca²⁺-sensing domain is an EF-hand that is N-terminal to the single transmembrane segment, and oriented in the lumen of the endoplasmic reticulum. Several laboratories have demonstrated that mutations in the EF-hand region, which presumably reduce Ca²⁺ affinity, result in constitutive activation of store-operated entry and I_{crac} . (Liou et al., 2005; Zhang et al., 2005; Spassova et al., 2006; Mercer et al., 2006).

Experiments examining the cellular distribution of Stim1 by histochemistry or fluorescently tagged Stim1 with real-time imaging techniques have provided intriguing information on the cell biology of this Ca²⁺ sensor. Liou et al. reported that Stim1 appeared to be located in the endoplasmic reticulum (Liou et al., 2005). However, we observed Stim1 with enhanced yellow fluorescent protein fused to its N-terminus (EYFP-Stim1) in fibrillar, or possibly tubular, patterns in HEK293 cells (Mercer et al., 2006); these patterns are not apparent from labeling of generic endoplasmic reticulum in the same cell type (Ribeiro et al., 2000). A similar conclusion was reached by Baba et al., who imaged Stim1 and endoplasmic reticulum markers within the same cell (Baba et al., 2006). In that study, the unique distribution of Stim1 was apparently dependent on the microtubular cytoskeleton (Baba et al., 2006). Thus, Stim1 may reside on a distinct organelle or, more likely, a distinct compartment within the endoplasmic reticulum. This would be consistent with a number of previous studies that concluded that the pool of Ca²⁺ that regulates store-operated channels is a small component of the total endoplasmic reticulum (Ribeiro and Putney, 1996; Parekh et al., 1997; Huang and Putney, 1998; Broad et al., 1999; Turner et al., 2003; Wisnoskey et al., 2003).

An even more exciting finding is that dissociation of Ca²⁺ causes Stim1 to redistribute into punctate structures and move closer to the plasma membrane (Liou et al., 2005; Zhang et al., 2005; Spassova et al., 2006; Mercer et al., 2006). However, a major point of controversy remains the subcellular localization of Stim1 when it redistributes into punctae. Liou et al. reported that EYFP-Stim1 moved close to the plasma membrane but, using an antibody directed against the EYFP on the N-terminus, they failed to detect it on the plasma membrane (Liou et al., 2005). By contrast, on the basis of surface biotinylation studies, Zhang et al. reported that Stim1 actually translocates into the plasma membrane (Zhang et al., 2005). A third position was taken by Spassova et al. (Spassova et al., 2006). They reported that a fraction of Stim1 is present in the plasma

membrane, where it appears to function in store-operated entry (this conclusion was based on the inhibition of entry by extracellular application of an anti-Stim1 antibody), but they failed to see any change in the amount of plasma membrane Stim1 following Ca^{2+} store depletion.

Mercer et al. (Mercer et al., 2006) have repeated the experiments performed by Liou et al. (Liou et al., 2005), using the same EYFP-tagged Stim1. These authors confirmed that no EYFP-Stim1 can be detected on the plasma membrane of HEK293 cells by surface antibody staining and confocal microscopy or by flow cytometry (Mercer et al., 2006). Xu et al. reached a similar conclusion, on the basis of studies using a Stim1 N-terminally tagged with a pH-sensitive fluorescent indicator (Xu et al., 2006). Likewise, Wu et al. failed to observe plasma membrane Stim1 by immuno-electronmicroscopy, by using an N-terminally horseradish-peroxidase-tagged Stim1 (Wu et al., 2006). However, Spassova et al. clearly observed surface labeling of Stim1 by flow cytometry, using an untagged construct and an antibody directed against the N-terminus (Spassova et al., 2006). Furthermore, the first report on Stim1, prior to the discovery that it acts as an initiator of capacitative Ca^{2+} entry, identified it as a surface protein on stromal cells involved in their interaction with B lymphocytes (Oritani and Kincade, 1996), and this surface localization was substantiated in subsequent studies (Manji et al., 2000; Williams et al., 2002).

A recent report by (Hauser and Tsien, 2007) provides an explanation for these disparate findings: the large additions, such as EYFP, to the N-terminus of Stim1 prevent its trafficking to the plasma membrane. This is actually fortuitous, since it allows us to assess the function of Stim1 by using a construct restricted to intracellular sites. EYFP-Stim1 is fully functional, both rescuing store-operated entry after RNAi-mediated knockdown of Stim1, and generating constitutive Ca^{2+} entry following mutation of the EF hand domain (Liou et al., 2005; Mercer et al., 2006). Similarly, Baba et al. found that N-terminally Flag-tagged Stim1 could not be detected in the plasma membrane of a B-cell line but still fully supports store-operated Ca^{2+} entry (Baba et al., 2006). In this case, they knocked out Stim1 by targeted gene disruption, leaving no doubt that the N-terminally tagged Stim1 can function in the absence of any residual wild-type protein. As I discuss below, co-expression of Stim1 with Orai1 greatly increases store-operated entry; however, even these very large I_{crac} -like currents are supported by EYFP-Stim1, and again no surface expression can be detected (Mercer et al., 2006). Finally, Huang et al. demonstrated that the cytoplasmic C-terminal region of Stim1, expressed in the cytoplasm, was sufficient to activate store-operated entry (Huang et al., 2006). Clearly this construct cannot produce any plasma membrane Stim1 in the correct orientation. My conclusion is that Stim1 does not translocate to the plasma membrane in response to Ca^{2+} -store depletion, and that plasma membrane Stim1, although probably present, does not play an obligatory role in activating store-operated channels.

Orai

Orai1 was first reported (and named) by Feske et al. through a combination of gene mapping in a family with an immunodeficiency attributed to loss of I_{crac} and a whole-genome screen of *Drosophila* S2 cells (Feske et al., 2006).

Almost simultaneously, two other groups, also using a similar whole-genome screen of S2 cells, reported that Orai [termed CRACM1 by Vig et al. (Vig et al., 2006a)] is essential for store-operated entry and for I_{crac} (Zhang et al., 2006; Vig et al., 2006a). Orai1 is located in the plasma membrane and appears to have four transmembrane domains. Unlike the case for Stim1, which was included in restricted screens based on its signaling domains, Orai1 has no recognizable signaling or channel-like domains and thus required the full genome screen for its detection. When Orai and Stim are co-expressed, they synergize to generate unusually large I_{crac} -like currents (Zhang et al., 2006; Mercer et al., 2006; Peinelt et al., 2006; Soboloff et al., 2006b), which are indistinguishable from native I_{crac} in both biophysical and pharmacological properties (Zhang et al., 2006; Mercer et al., 2006; Peinelt et al., 2006; Soboloff et al., 2006b). These two proteins can thus fully recapitulate the properties of I_{crac} . This does not necessarily mean that no other players are involved; however, other proteins functioning in a stoichiometric complex with Stim1 and/or Orai1 would have to be constitutively present in considerable excess.

The large I_{crac} -like currents observed upon co-expression of Stim1 and Orai1 immediately led to speculation that Orai1 is the CRAC channel itself or a subunit of it. However, there are no obvious channel-pore-like sequences in Orai1. Three laboratories focused on a string of acidic residues near the extracellular boundary of the first transmembrane domain (Prakriya et al., 2006; Yeromin et al., 2006; Vig et al., 2006b). The most interesting mutations affect glutamate 106 in mammalian Orai1, and the corresponding residue in *Drosophila* Orai. Mutation to alanine results in a non-functional channel; however, mutation to an aspartate (E106D; E180D in *Drosophila* Stim) results in a functional channel that has markedly reduced selectivity for Ca^{2+} . Acidic residues just downstream from E180 in *Drosophila* Orai confer the well-known high sensitivity of I_{crac} to Gd^{3+} (Yeromin et al., 2006). These findings provide strong evidence that this region functions as part of the Ca^{2+} selectivity filter and indicate that Orai1 is indeed a pore-forming subunit of the CRAC channel.

Prakriya et al. (Prakriya et al., 2006) and Vig et al. (Vig et al., 2006b) have also investigated a glutamate residue at position 190. Mutation of this residue to aspartate or even alanine had no effect on channel function; however, alteration to glutamine (E190Q) resulted in diminished Ca^{2+} selectivity. Since the complete loss of charge from this amino acid (the mutation to alanine) had no effect on channel selectivity, the mutation to glutamine may alter the secondary or tertiary structure of the channel, rather than indicating a function as part of the Ca^{2+} -binding site in the channel pore.

Mammalian cells have two additional proteins, Orai2 and Orai3. Orai2 appears to function similarly to Orai1, at least when expressed with Stim1 in HEK293 cells (Mercer et al., 2006). The Orai2 Ca^{2+} currents were somewhat smaller, however, and whether this is an intrinsic property of Orai2 channels or indicates a lower level of expression in this particular study is not known. Orai3 Ca^{2+} currents were even smaller; those resulting from co-expression of Stim1 and Orai3 in HEK293 cells were below the limits of detection. However, Orai3-dependent currents can be observed when Na^+ carries the current (DeHaven et al., 2007), and Orai3 can rescue store-operated entry when Orai1 is knocked down in HEK293 cells (Mercer et al., 2006). Whether Orai2 and Orai3 form distinct

store-operated channels in specific cell types, or function as subunits of heteromeric channels with Orai1 will be a topic of future investigation.

Stim1-Orai1 communication

The evidence is very strong that the signaling pathway for capacitative Ca²⁺ entry begins with the Ca²⁺ sensor Stim1, and culminates in the activation of channels composed partly or wholly of Orai subunits. The obvious question is how Stim1 conveys information from depleted Ca²⁺ stores to Orai channels. The simplest answer is that, when Stim1 coalesces into punctae and approaches the plasma membrane, it directly interacts with Orai channels there. In support of this idea, Yeromin et al. reported that *Drosophila* Stim and Orai can be co-immunoprecipitated, and this association is increased by depletion of Ca²⁺ stores (Yeromin et al., 2006). However, Feske et al. failed to observe any co-immunoprecipitation of mammalian Stim1 and Orai1 (Feske et al., 2006). Vig et al. reported co-immunoprecipitation of transfected and tagged Stim1 and Orai1 but did not assess effects of store depletion (Vig et al., 2006b). Nonetheless, it is clear from the work of Luik et al. (Luik et al., 2006) and Xu et al. (Xu et al., 2006) that communication between Stim1 and Orai1 occurs over very short distances. These investigators observed that Orai1 is recruited to sites of Stim1 punctae formation, and Luik et al. (Luik et al., 2006) showed that Ca²⁺ entry is spatially restricted to these plasma membrane sites as well. Wu et al. estimated that Stim1 localizes in junctional endoplasmic reticulum within 10–25 nm of the plasma membrane (Wu et al., 2006). Finally, although a direct interaction between Stim1 and Orai seems the simplest mechanism for signaling from intracellular stores to the plasma membrane channels, there is nothing to rule out the generation of a secondary message downstream of Stim1 that acts on Orai – for example, the proposed CIF and iPLA₂ pathway (Bolotina and Csutora, 2005).

Conclusions

In just the past two years, the discoveries of Stim and Orai proteins have revolutionized our thinking about capacitative Ca²⁺ entry and *I*_{crac}. It is remarkable how quickly different laboratories confirmed the dependence of *I*_{crac} on these two proteins and confirmed their ability to recapitulate the long-known properties of CRAC channels. In fact, it could be said that these are the only aspects of the *I*_{crac} story upon which all (or most) seem to agree. The current findings about Stim1 and Orai action are consistent with a number of earlier ideas about store-operated entry. For example, the presence of a small, specialized component of the endoplasmic reticulum dedicated to communicating with plasma membrane channels; the observation that biophysical properties of CRAC channels differ substantially from conventional ion channels, which indicated that they might also differ substantially in their structure; the general finding that Ca²⁺ refills the stores efficiently without causing a rise in global cytoplasmic Ca²⁺; and the suggested close physical association between sites of endoplasmic reticulum signaling and plasma membrane Ca²⁺ entry. One might argue that Irvine's original idea (Irvine, 1990)




Fig. 1. The roles of Stim1 and Orai. Agonist (Ag) activation of a plasma membrane receptor (R) results in formation of Ins(1,4,5)P₃, which activates the Ins(1,4,5)P₃ receptor (IP3R) causing discharge of stored Ca²⁺ from a subcompartment of the endoplasmic reticulum. Within this subcompartment, Ca²⁺ binds reversibly to an EF hand motif in Stim1; depletion of Ca²⁺ results in dissociation of Ca²⁺ from Stim1, which causes Stim1 to redistribute within the endoplasmic reticulum to areas near Orai channels that reside in the plasma membrane. Stim1 then activates Ca²⁺-selective Orai channels; the mechanism by which this activation is accomplished is unknown at present. Stim1 is also present in the plasma membrane, although its function there is unclear. TRPC channels can also be activated by phospholipase C (PLC)-dependent mechanisms. There is evidence that in some instances, perhaps when combined with other subunits, they can function as store-operated channels, perhaps also involving Stim1 as an activator. Figure modified from Putney, 2007 (Putney, 2007).

of conformational coupling is at least partially vindicated by the current model. However, it appears to be Stim1 rather than Ins(1,4,5)P₃ receptors that are responsible for this coupling.

Fig. 1 summarizes the store-operated signaling pathway, incorporating the roles of Stim1 and Orai. Several general questions remain. (1) Are any other players or proteins necessary for activation of capacitative Ca²⁺ entry and/or *I*_{crac}? (2) What is the role, if any, of plasma membrane Stim1? (3) What is the composition of native CRAC channels – are these Orai homo- or heteromultimers or combinations with TRPs? (4) Are there mechanisms of store-operated entry involving other modes of activation and other store-operated channels? The availability of Stim1 and Orai cDNAs will lead to a detailed structural understanding of the various modes of regulation and activation of this pathway. The development of animal models will lead to a better understanding of the roles of these proteins in specific physiological and pathological processes. Such information may ultimately prove useful in design of novel pharmacological reagents to aid in the treatment of a number of diseases in which the store-operated entry pathway is thought to play a role.

David Miller and Stephen Shears read the manuscript and provided helpful comments. Research from the author's laboratory described in this review was supported by the Intramural Research Program of the NIH, National Institute of Environmental Health Sciences.

References

- Acharya, J. K., Jalink, K., Hardy, R. W., Hartenstein, V. and Zuker, C. S. (1997). InsP₃ receptor is essential for growth and differentiation but not for vision in *Drosophila*. *Neuron* **18**, 881–887.
- Almers, W. and McCleskey, E. W. (1984). Non-selective conductance in calcium

- channels of frog muscle: calcium selectivity in a single-file pore. *J. Physiol.* **353**, 585-608.
- Baba, Y., Hayashi, K., Fujii, Y., Mizushima, A., Watarai, H., Wakamori, M., Numaga, T., Mori, Y., Iino, M., Hikida, M. et al. (2006). Coupling of STIM1 to store-operated Ca^{2+} entry through its constitutive and inducible movement in the endoplasmic reticulum. *Proc. Natl. Acad. Sci. USA* **103**, 16704-16709.
- Bakowski, D., Glitsch, M. D. and Parekh, A. B. (2001). An examination of the secretion-like coupling model for the activation of the Ca^{2+} release-activated Ca^{2+} current I_{crac} in RBL-1 cells. *J. Physiol. Lond.* **532**, 55-71.
- Barritt, G. J. (1999). Receptor-activated Ca^{2+} inflow in animal cells: a variety of pathways tailored to meet different intracellular Ca^{2+} signalling requirements. *Biochem. J.* **337**, 153-169.
- Berridge, M. J. (1995). Capacitative calcium entry. *Biochem. J.* **312**, 1-11.
- Berridge, M. J., Lipp, P. and Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell Biol.* **1**, 11-21.
- Bian, X., Hughes, F. M., Jr, Huang, Y., Cidlovski, J. A. and Putney, J. W., Jr (1997). Roles of cytoplasmic Ca^{2+} and intracellular Ca^{2+} stores in induction and suppression of apoptosis in S49 cells. *Am. J. Physiol.* **272**, C1241-C1249.
- Birnbaumer, L., Boulay, G., Brown, D., Jiang, M., Dietrich, A., Mikoshiba, K., Zhu, X. and Qin, N. (2000). Mechanism of capacitative Ca^{2+} entry (CCE): interaction between IP₃ receptor and TRP links the internal calcium storage compartment to plasma membrane CCE channels. *Recent Prog. Horm. Res.* **55**, 127-161.
- Bolotina, V. M. and Csutora, P. (2005). CIF and other mysteries of the store-operated Ca^{2+} -entry pathway. *Trends Biochem. Sci.* **30**, 378-387.
- Broad, L. M., Armstrong, D. L. and Putney, J. W., Jr (1999). Role of the IP₃ receptor in Ca^{2+} feedback inhibition of calcium release-activated calcium current (I_{crac}). *J. Biol. Chem.* **274**, 32881-32888.
- Casteels, R. and Droogmans, G. (1981). Exchange characteristics of the noradrenaline-sensitive calcium store in vascular smooth muscle cells of rabbit ear artery. *J. Physiol. Lond.* **317**, 263-279.
- DeHaven, W. I., Smyth, J. T., Boyles, R. R., and Putney, J. W. (2007). Calcium inhibition and calcium potentiation of orai1, orai2 and orai3 calcium-release-activated calcium channels. *J. Biol. Chem.* in press, doi/10.1074/jbc.M611374200, PMID: 17452328.
- Dietrich, A., Schnitzler, M., Kalwa, H., Storch, U. and Gudermann, T. (2005). Functional characterization and physiological relevance of the TRPC3/6/7 subfamily of cation channels. *Naunyn Schmiedeberg's Arch. Pharmacol.* **371**, 257-265.
- Elliott, A. C. (2001). Recent developments in non-excitatory cell calcium entry. *Cell Calcium* **30**, 73-93.
- Favre, C. J., Nüsse, O., Lew, D. P. and Krause, K.-H. (1996). Store-operated Ca^{2+} influx: what is the message from the stores to the membrane? *J. Lab. Clin. Med.* **128**, 19-26.
- Feske, S., Gwack, Y., Prakriya, M., Srikanth, S., Puppel, S. H., Tanasa, B., Hogan, P. G., Lewis, R. S., Daly, M. and Rao, A. (2006). A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* **441**, 179-185.
- Gailly, P. (1998). Ca^{2+} entry in CHO cells, after Ca^{2+} stores depletion, is mediated by arachidonic acid. *Cell Calcium* **24**, 293-304.
- Gill, D. L. and Patterson, R. W. L. (2004). Toward a consensus on the operation of receptor-induced calcium entry signals. *Sci. STKE* **2004**, pe39.
- Golovina, V. A. (2005). Visualization of localized store-operated calcium entry in mouse astrocytes. Close proximity to the endoplasmic reticulum. *J. Physiol.* **564**, 737-749.
- Graier, W. F., Simecek, S. and Sturek, M. (1995). Cytochrome P450 mono-oxygenase-regulated signalling of Ca^{2+} entry in human and bovine endothelial cells. *J. Physiol. Lond.* **482**, 259-274.
- Hardie, R. C. and Minke, B. (1992). The trp gene is essential for a light-activated Ca^{2+} channel in drosophila photoreceptors. *Neuron* **8**, 643-651.
- Hardie, R. C. and Minke, B. (1993). Novel Ca^{2+} channels underlying transduction in *Drosophila* photoreceptors: implications for phosphoinositide-mediated Ca^{2+} mobilization. *Trends Neurosci.* **16**, 371-376.
- Hausser, C. T. and Tsien, R. Y. (2007). A hexahistidine-Zn²⁺-dye label reveals STIM1 surface exposure. *Proc. Natl. Acad. Sci. USA* **104**, 3693-3697.
- Hess, P. and Tsien, R. W. (1984). Mechanism of ion permeation through calcium channels. *Nature* **309**, 453-456.
- Hoth, M. and Penner, R. (1992). Depletion of intracellular calcium stores activates a calcium current in mast cells. *Nature* **355**, 353-355.
- Hoth, M. and Penner, R. (1993). Calcium release-activated calcium current in rat mast cells. *J. Physiol. Lond.* **465**, 359-386.
- Huang, G. N., Zeng, W., Kim, J. Y., Yuan, J. P., Han, L., Muallem, S. and Worley, P. F. (2006). STIM1 carboxyl-terminus activates native SOC, I_{crac} and TRPC1 channels. *Nat. Cell Biol.* **8**, 1003-1010.
- Huang, Y. and Putney, J. W., Jr (1998). Relationship between intracellular calcium store depletion and calcium release-activated calcium current (I_{crac}) in a mast cell line (RBL-1). *J. Biol. Chem.* **273**, 19554-19559.
- Irvine, R. F. (1990). "Quantal" Ca^{2+} release and the control of Ca^{2+} entry by inositol phosphates - a possible mechanism. *FEBS Lett.* **263**, 5-9.
- Jaconi, M., Pyle, J., Bortolon, R., Ou, J. and Clapham, D. (1997). Calcium release and influx colocalize to the endoplasmic reticulum. *Curr. Biol.* **7**, 599-602.
- Kiselyov, K., Xu, X., Mozhayeva, G., Kuo, T., Pessah, I., Mignery, G., Zhu, X., Birnbaumer, L. and Muallem, S. (1998). Functional interaction between InsP₃ receptors and store-operated Htrp3 channels. *Nature* **396**, 478-482.
- Krause, E., Pfeiffer, F., Schmid, A. and Schulz, I. (1996). Depletion of intracellular calcium stores activates a calcium conducting nonselective cation current in mouse pancreatic acinar cells. *J. Biol. Chem.* **271**, 32523-32528.
- Lewis, R. S. and Cahalan, M. D. (1989). Mitogen-induced oscillations of cytosolic Ca^{2+} and transmembrane Ca^{2+} current. *Cell Regul.* **1**, 99-112.
- Lievremont, J. P., Bird, G. St J. and Putney, J. W., Jr (2004). Canonical transient receptor potential TRPC7 can function as both a receptor- and store-operated channel in HEK-293 cells. *Am. J. Physiol. Cell Physiol.* **287**, C1709-C1716.
- Liou, J., Kim, M. L., Heo, W. D., Jones, J. T., Myers, J. W., Ferrell, J. E., Jr and Meyer, T. (2005). STIM is a Ca^{2+} sensor essential for Ca^{2+} -store-depletion-triggered Ca^{2+} influx. *Curr. Biol.* **15**, 1235-1241.
- Luik, R. M., Wu, M. M., Buchanan, J. and Lewis, R. S. (2006). The elementary unit of store-operated Ca^{2+} entry: local activation of CRAC channels by STIM1 at ER-plasma membrane junctions. *J. Cell Biol.* **174**, 815-825.
- Manji, S. S., Parker, N. J., Williams, R. T., Van Stekelenburg, L., Pearson, R. B., Dziadek, M. and Smith, P. J. (2000). STIM1: a novel phosphoprotein located at the cell surface. *Biochim. Biophys. Acta* **1481**, 147-155.
- Mercer, J. C., DeHaven, W. I., Smyth, J. T., Wedel, B., Boyles, R. R., Bird, G. S. and Putney, J. W., Jr (2006). Large store-operated calcium-selected currents due to co-expression of orai1 or orai2 with the intracellular calcium sensor, stim1. *J. Biol. Chem.* **281**, 24979-24990.
- Mignen, O., Thompson, J. L., Yule, D. I. and Shuttleworth, T. J. (2005). Agonist activation of arachidonate-regulated Ca^{2+} -selective (ARC) channels in murine parotid and pancreatic acinar cells. *J. Physiol.* **564**, 791-801.
- Mori, Y., Wakamori, M., Miyakawa, T., Hermosura, M., Hara, Y., Nishida, M., Hirose, K., Mizushima, A., Kurosaki, M., Mori, E. et al. (2002). Transient receptor potential 1 regulates capacitative Ca^{2+} entry and Ca^{2+} release from endoplasmic reticulum in B lymphocytes. *J. Exp. Med.* **195**, 673-681.
- Oritani, K. and Kincade, P. W. (1996). Identification of stromal cell products that interact with pre-B cells. *J. Cell Biol.* **134**, 771-782.
- Pandolf, S. J. and Schoeffield-Payne, M. S. (1990). Cyclic GMP mediates the agonist-stimulated increase in plasma membrane calcium entry in the pancreatic acinar cell. *J. Biol. Chem.* **265**, 12846-12853.
- Parekh, A. B. and Penner, R. (1997). Store depletion and calcium influx. *Physiol. Rev.* **77**, 901-930.
- Parekh, A. B. and Putney, J. W., Jr (2005). Store-operated calcium channels. *Physiol. Rev.* **85**, 757-810.
- Parekh, A. B., Fleig, A. and Penner, R. (1997). The store-operated calcium current I_{CRAC}: nonlinear activation by InsP₃ and dissociation from calcium release. *Cell* **89**, 973-980.
- Patterson, R. L., van Rossum, D. B. and Gill, D. L. (1999). Store-operated Ca^{2+} entry: evidence for a secretion-like coupling model. *Cell* **98**, 487-499.
- Peinelt, C., Vig, M., Koomoa, D. L., Beck, A., Nadler, M. J. S., Koblan-Huberson, M., Lis, A., Fleig, A., Penner, R. and Kinet, J. P. (2006). Amplification of CRAC current by STIM1 and CRACM1 (Orai1). *Nat. Cell Biol.* **8**, 771-773.
- Prakriya, M., Feske, S., Gwack, Y., Srikanth, S., Rao, A. and Hogan, P. G. (2006). Orai1 is an essential pore subunit of the CRAC channel. *Nature* **443**, 230-233.
- Putney, J. W., Jr (1977). Muscarinic, alpha-adrenergic and peptide receptors regulate the same calcium influx sites in the parotid gland. *J. Physiol. Lond.* **268**, 139-149.
- Putney, J. W., Jr (1990). Capacitative calcium entry revisited. *Cell Calcium* **11**, 611-624.
- Putney, J. W., Jr (1997). *Capacitative Calcium Entry*. Austin, TX: Landes Biomedical Publishing.
- Putney, J. W., Jr (2001). Pharmacology of capacitative calcium entry. *Mol. Interv.* **1**, 84-94.
- Putney, J. W., Jr (2005). Physiological mechanisms of TRPC activation. *Pflugers Arch.* **451**, 29-34.
- Putney, J. W., Jr (2007). Recent breakthroughs in the molecular mechanism of capacitative calcium entry (with thoughts on how we got here). *Cell Calcium* in press, doi:10.1016/j.ccc.2007.01.011, PMID: 17349691
- Putney, J. W., Jr and Bird, G. St J. (1993). The signal for capacitative calcium entry. *Cell* **75**, 199-201.
- Putney, J. W., Jr and McKay, R. R. (1999). Capacitative calcium entry channels. *BioEssays* **21**, 38-46.
- Putney, J. W., Jr, Broad, L. M., Braun, F.-J., Lievremont, J.-P. and Bird, G. St J. (2001). Mechanisms of capacitative calcium entry. *J. Cell Sci.* **114**, 2223-2229.
- Randriamampita, C. and Tsien, R. Y. (1993). Emptying of intracellular Ca^{2+} stores releases a novel small messenger that stimulates Ca^{2+} influx. *Nature* **364**, 809-814.
- Ribeiro, C. M. P. and Putney, J. W., Jr (1996). Differential effects of protein kinase C activation on calcium storage and capacitative calcium entry in NIH 3T3 cells. *J. Biol. Chem.* **271**, 21522-21528.
- Ribeiro, C. M. P., McKay, R. R., Hosoki, E., Bird, G. St J. and Putney, J. W., Jr (2000). Effects of elevated cytoplasmic calcium and protein kinase C on endoplasmic reticulum structure and function in HEK293 cells. *Cell Calcium* **27**, 175-185.
- Roos, J., DiGregorio, P. J., Yeromin, A. V., Ohlsen, K., Lioudyno, M., Zhang, S., Safrina, O., Kozak, J. A., Wagner, S. L., Cahalan, M. D. et al. (2005). STIM1, an essential and conserved component of store-operated Ca^{2+} channel function. *J. Cell Biol.* **169**, 435-445.
- Rosado, J. A., Redondo, P. C., Sage, S. O., Pariente, J. A. and Salido, G. M. (2005). Store-operated Ca^{2+} entry: vesicle fusion or reversible trafficking and de novo conformational coupling? *J. Cell. Physiol.* **205**, 262-269.
- Rzagalinski, B. A., Willoughby, K. A., Hoffman, S. W., Falck, J. R. and Ellis, E. F. (1999). Calcium influx factor, further evidence it is 5,6-epoxyeicosatrienoic acid. *J. Biol. Chem.* **274**, 175-185.
- Sage, S. O., Sargeant, P., Jenner, S. and Fardale, R. W. (1994). Tyrosine phosphorylation and Ca^{2+} influx. A further mechanism for store-dependent Ca^{2+} entry? *Trends Pharmacol. Sci.* **15**, 282.

- Sambrook, J. F.** (1990). The involvement of calcium in transport of secretory proteins from the endoplasmic reticulum. *Cell* **61**, 197-199.
- Scott, C. C., Furuya, W., Trimble, W. S. and Grinstein, S.** (2003). Activation of store-operated calcium channels: assessment of the role of snare-mediated vesicular transport. *J. Biol. Chem.* **278**, 30534-30539.
- Soboloff, J., Spassova, M. A., Hewavitharana, T., He, L. P., Xu, W., Johnstone, L. S., Dziadek, M. A. and Gill, D. L.** (2006a). STIM2 is an inhibitor of STIM1-mediated store-operated Ca²⁺ entry. *Curr. Biol.* **16**, 1465-1470.
- Soboloff, J., Spassova, M. A., Tang, X. D., Hewavitharana, T., Xu, W. and Gill, D. L.** (2006b). Orai1 and STIM reconstitute store-operated calcium channel function. *J. Biol. Chem.* **281**, 20661-20665.
- Somasundaram, B., Norman, J. C. and Mahaut-Smith, M. P.** (1995). Primaquine, an inhibitor of vesicular transport, blocks the calcium release activated current in rat megakaryocytes. *Biochem. J.* **309**, 725-729.
- Spassova, M., Soboloff, J., He, L.-P., Xu, W., Dziadek, M. and Gill, D. L.** (2006). STIM1 has a plasma membrane role in the activation of store-operated Ca²⁺ channels. *Proc. Natl. Acad. Sci. USA* **103**, 4040-4045.
- Takemura, H., Hughes, A. R., Thastrup, O. and Putney, J. W., Jr** (1989). Activation of calcium entry by the tumor promoter, thapsigargin, in parotid acinar cells. Evidence that an intracellular calcium pool, and not an inositol phosphate, regulates calcium fluxes at the plasma membrane. *J. Biol. Chem.* **264**, 12266-12271.
- Teasdale, R. D. and Jackson, M. R.** (1996). Signal-mediated sorting of membrane proteins between the endoplasmic reticulum and the golgi apparatus. *Annu. Rev. Cell Dev. Biol.* **12**, 27-54.
- Thomas, D. and Hanley, M. R.** (1995). Evaluation of calcium influx factors from stimulated Jurkat T-lymphocytes by microinjection into *Xenopus* oocytes. *J. Biol. Chem.* **270**, 6429-6432.
- Thompson, S. H.** (1997). Cyclic GMP-gated channels in a sympathetic neuron cell line. *J. Gen. Physiol.* **110**, 155-164.
- Trebak, M., Bird, G. St J., McKay, R. R., Birnbaumer, L. and Putney, J. W., Jr** (2003). Signaling mechanism for receptor-activated TRPC3 channels. *J. Biol. Chem.* **278**, 16244-16252.
- Trepakova, E. S., Csutora, P., Hunton, D. L., Marchase, R. B., Cohen, R. A. and Bolotina, V. M.** (2000). Calcium influx factor (CIF) directly activates store-operated cation channels in vascular smooth muscle cells. *J. Biol. Chem.* **275**, 26158-26163.
- Tsukamoto, A. and Kaneko, Y.** (1993). Thapsigargin, a Ca²⁺-ATPase inhibitor, depletes the intracellular Ca²⁺ pool and induces apoptosis in human hepatoma cells. *Cell Biol. Int.* **17**, 969-970.
- Turner, H., Fleig, A., Stokes, A., Kinert, J. P. and Penner, R.** (2003). Discrimination of intracellular calcium store subcompartments using TRPV1 (transient receptor potential channel, vanilloid subfamily member 1) release channel activity. *Biochem. J.* **371**, 341-350.
- Vazquez, G., Lièvre, J.-P., Bird, G. St J. and Putney, J. W., Jr** (2001). Human Trp3 forms both inositol trisphosphate receptor-dependent and receptor-independent store-operated cation channels in DT40 avian B-lymphocytes. *Proc. Natl. Acad. Sci. USA* **98**, 11777-11782.
- Vazquez, G., Wedel, B. J., Trebak, M., Bird, G. St J. and Putney, J. W., Jr** (2003). Expression level of TRPC3 channel determines its mechanism of activation. *J. Biol. Chem.* **278**, 21649-21654.
- Vig, M., Peinelt, C., Beck, A., Koomoa, D. L., Rabah, D., Koblan-Huberson, M., Kraft, S., Turner, H., Fleig, A., Penner, R. et al.** (2006a). CRACM1 is a plasma membrane protein essential for store-operated Ca²⁺ entry. *Science* **312**, 1220-1223.
- Vig, M., Beck, A., Billingsley, J. M., Lis, A., Parvez, S., Peinelt, C., Koomoa, D. L., Soboloff, J., Gill, D. L. and Fleig, A.** (2006b). CRACM1 multimers form the ion-selective pore of the CRAC channel. *Curr. Biol.* **16**, 2073-2079.
- Wedel, B., Boyles, R. R., Putney, J. W., and Bird, G. S.** (2007). Role of the Store-operated Calcium Entry Proteins, Stim1 and Orai1, in Muscarinic-Cholinergic Receptor Stimulated Calcium Oscillations in Human Embryonic Kidney Cells. *J. Physiol.* **579**, 679-689.
- Williams, R. T., Senior, P. V., Van Stekelenburg, L., Layton, J. E., Smith, P. J. and Dziadek, M. A.** (2002). Stromal interaction molecule 1 (STIM1), a transmembrane protein with growth suppressor activity, contains an extracellular SAM domain modified by N-linked glycosylation. *Biochim. Biophys. Acta* **1596**, 131-137.
- Wisnoskey, B. J., Sinkins, W. G. and Schilling, W. P.** (2003). Activation of vanilloid receptor type I in the endoplasmic reticulum fails to activate store-operated Ca²⁺ entry. *Biochem. J.* **372**, 517-528.
- Wu, M. M., Buchanan, J., Luik, R. M. and Lewis, R. S.** (2006). Ca²⁺ store depletion causes STIM1 to accumulate in ER regions closely associated with the plasma membrane. *J. Cell Biol.* **174**, 803-813.
- Xu, P., Lu, J., Li, Z., Yu, X., Chen, L. and Xu, T.** (2006). Aggregation of STIM1 underneath the plasma membrane induces clustering of Orai1. *Biochem. Biophys. Res. Commun.* **350**, 969-976.
- Xu, X., Star, R. A., Tortorici, G. and Muallem, S.** (1994). Depletion of intracellular Ca²⁺ stores activates nitric-oxide synthase to generate cGMP and regulate Ca²⁺ influx. *J. Biol. Chem.* **269**, 12645-12653.
- Yao, Y., Ferrer-Montiel, A. V., Montal, M. and Tsien, R. Y.** (1999). Activation of store-operated Ca²⁺ current in *Xenopus* oocytes requires SNAP-25 but not a diffusible messenger. *Cell* **98**, 475-485.
- Yeromin, A. V., Zhang, S. L., Jiang, W., Yu, Y., Safrina, O. and Cahalan, M. D.** (2006). Molecular identification of the CRAC channel by altered ion selectivity in a mutant of Orai. *Nature* **443**, 226-229.
- Zeng, F., Xu, S. Z., Jackson, P. K., McHugh, D., Kumar, B., Fountain, S. J. and Beech, D. J.** (2004). Human TRPC5 channel activated by a multiplicity of signals in a single cell. *J. Physiol.* **559**, 739-750.
- Zhang, S. L., Yu, Y., Roos, J., Kozak, J. A., Deerinck, T. J., Ellisman, M. H., Stauderman, K. A. and Cahalan, M. D.** (2005). STIM1 is a Ca²⁺ sensor that activates CRAC channels and migrates from the Ca²⁺ store to the plasma membrane. *Nature* **437**, 902-905.
- Zhang, S. L., Yeromin, A. V., Zhang, X. H., Yu, Y., Safrina, O., Penna, A., Roos, J., Stauderman, K. A. and Cahalan, M. D.** (2006). Genome-wide RNAi screen of Ca²⁺ influx identifies genes that regulate Ca²⁺ release-activated Ca²⁺ channel activity. *Proc. Natl. Acad. Sci. USA* **103**, 9357-9362.
- Zhu, X., Jiang, M., Peyton, M., Boulay, G., Hurst, R., Stefani, E. and Birnbaumer, L.** (1996). *trp*, a novel mammalian gene family essential for agonist-activated capacitative Ca²⁺ entry. *Cell* **85**, 661-671.
- Zhu, X., Jiang, M. and Birnbaumer, L.** (1998). Receptor-activated Ca²⁺ influx via human Trp3 stably expressed in human embryonic kidney (HEK)293 cells. Evidence for a non-capacitative calcium entry. *J. Biol. Chem.* **273**, 133-142.
- Zitt, C., Zobel, A., Obukhov, A. G., Harteneck, C., Kalkbrenner, F., Lückhoff, A. and Schultz, G.** (1996). Cloning and functional expression of a human Ca²⁺-permeable cation channel activated by calcium store depletion. *Neuron* **16**, 1189-1196.
- Zweifach, A. and Lewis, R. S.** (1993). Mitogen-regulated Ca²⁺ current of T lymphocytes is activated by depletion of intracellular Ca²⁺ stores. *Proc. Natl. Acad. Sci. USA* **90**, 6295-6299.
- Zweifach, A. and Lewis, R. S.** (1995). Slow calcium-dependent inactivation of depletion-activated calcium current. *J. Biol. Chem.* **270**, 14445-14451.
- Zweifach, A. and Lewis, R. S.** (1996). Calcium-dependent potentiation of store-operated calcium channels in T lymphocytes. *J. Gen. Physiol.* **107**, 597-610.