

Eukaryotic chemotaxis at a glance

Anna Bagorda and Carole A. Parent*

Laboratory of Cellular and Molecular Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda MD 20892-4256, USA

*Author for correspondence (e-mail: parentc@mail.nih.gov)

Journal of Cell Science 121, 2621-2624
Published by The Company of Biologists 2008
doi:10.1242/jcs.018077

Supplementary material available online at <http://jcs.biologists.org/cgi/content/full/121/16/2621/DC1>

Introduction


Chemotaxis is a fundamental process in which cells migrate directionally when they are exposed to external chemical

gradients. It is exhibited by a wide variety of cell types and involves distinct strategies that depend on the environmental conditions. For example, the migration of fibroblasts or keratinocytes within tissues is relatively slow as it relies on strong adhesion forces to the extracellular matrix. Conversely, fast moving and highly deformable leukocytes navigate to sites of inflammation using dynamic actin remodeling and minimal adhesion forces. In this Cell Science at a Glance article, we describe the signals that control directed migration for two cellular systems that are commonly accepted as paradigms for fast movement: the genetically tractable eukaryotic unicellular organism *Dictyostelium discoideum* and mammalian neutrophils (see supplementary material Movies 1 and 2). These two systems are exceptionally tuned to dynamically sense and respond to very shallow external chemoattractant gradients. Although

evolutionarily separated by millions of years, their chemotactic machinery is controlled by highly conserved signal transduction pathways, which – in many instances – were first identified in *Dictyostelium*. The spatial activation of these pathways ultimately leads to the redistribution of polymerized actin at the front of the cell for propulsion and the assembly of myosin II at the back for retraction.

Chemoattractants and their receptors

The first event that occurs during chemotaxis is the binding of a chemoattractant molecule to its specific receptor. In *Dictyostelium*, two main chemoattractant molecules promote chemotaxis: folic acid and cAMP (Mahadeo and Parent, 2006). Folic acid, a bacterial by-product, acts as a directional cue to attract *Dictyostelium* to their food



(See poster insert)

source during growth, as these organisms live as single cells that feed on bacteria. Under conditions of starvation, *Dictyostelium* enter a developmental program and lose their ability to sense folic acid. Within 4–5 hours, they then acquire the chemotactic machinery that is responsible for the detection, synthesis and degradation of cAMP, the most potent chemoattractant for *Dictyostelium* (Manahan et al., 2004; Saran et al., 2002). Concurrently, starved cells become highly chemotactic and migrate directionally to form aggregates that later differentiate into a multicellular structure that can resist harsh environmental conditions (Chisholm and Firtel, 2004).

Receptors for folic acid and cAMP belong to the G-protein-coupled receptor (GPCR) family, each member of which comprises an extracellular N-terminal domain followed by seven transmembrane helices and a C-terminal tail (Milligan and Kostenis, 2006). These receptors mediate most of their effects through heterotrimeric G proteins, although a subset of responses occur independently of G proteins (Brzostowski and Kimmel, 2001). Four distinct cAMP receptors (cAR1 to cAR4), which are expressed at different times during *Dictyostelium* development, have been cloned. cAR1 is expressed during early development, when the cells are highly chemotactic, and is linked to the G protein $G_{\alpha_2}\beta\gamma$ (Parent and Devreotes, 1996). Both cAR1 and G proteins are uniformly distributed around the cell periphery and remain this way during chemotaxis (Parent and Devreotes, 1999). The folic-acid receptor, which is linked to $G_{\alpha_4}\beta\gamma$, has yet to be cloned.

Leukocyte chemotaxis is induced by several chemoattractants and chemokines that signal through pertussis-toxin-sensitive G_i -coupled as well as G_{12} - and G_{13} -coupled GPCRs (Baggiolini, 2001; Niggli, 2003; Stephens et al., 2008). Classical chemoattractants include formylated peptides that are secreted by bacteria [such as N-formylmethionyl-leucyl-phenylalanine (fMLP)], products of the complement cascade (such as C5a) and phospholipid metabolites [such as leukotriene B₄ (LTB₄)], whereas the family of chemokines includes chemotactic mediators that are derived from a variety of cells. As first observed for cAR1 in *Dictyostelium*, C5a receptors remain uniformly distributed around the plasma

membrane in chemotaxing neutrophils (Niggli, 2003; Parent, 2004). Although an even distribution of receptors around the cell periphery provides optimal sensitivity to highly dynamic chemoattractant gradients, there are cases in which chemokine receptors redistribute to the front of migrating cells; such behavior characterizes CXC-chemokine receptor 4 (CXCR4) in hematopoietic progenitor cells and CC-chemokine receptor 5 (CCR5) in Jurkat cells (Manes et al., 2005).

Signal transduction pathways and their effectors

Binding of the chemoattractant to its receptor triggers the dissociation of the heterotrimeric G protein into α - and $\beta\gamma$ -subunits. In both *Dictyostelium* and mammalian neutrophils, the $\beta\gamma$ dimer acts as the main transducer of chemotactic signals and activates several downstream effectors, including adenylyl cyclase, guanylyl cyclase and phospholipase C (PLC) (Bagorda et al., 2006; Stephens et al., 2008). The $G\beta\gamma$ stimulation is locally transduced via the activation of phosphoinositide 3-kinase (PI3K) at the front of cells in a Ras-dependent manner, although $G\beta\gamma$ can directly regulate PI3K in neutrophils (Kolsch et al., 2008; Stephens et al., 2008). PI3K converts phosphatidylinositol (4,5)-bisphosphate (PtdIns(4,5) P_2) into phosphatidylinositol (3,4,5)-trisphosphate (PtdIns(3,4,5) P_3), thereby promoting a localized PtdIns(3,4,5) P_3 production. In *Dictyostelium*, the spatial restriction of the PtdIns(3,4,5) P_3 signal to the front is ensured by the presence of PTEN (the 3'-phosphatase) at the sides and back of cells (Parent, 2004). The PtdIns(3,4,5) P_3 that is locally generated and retained at the front of cells provides a binding site for a subset of PH-domain-containing proteins, which eventually translocate to the leading edge and act as nucleation factors to activate key effectors of migration (Iijima et al., 2002).

The *Dictyostelium* cytosolic regulator of adenylyl cyclase (CRAC) protein was the first PH-domain-containing protein that was shown to translocate specifically to the front of chemotaxing cells. CRAC, originally identified as regulator for adenylyl-cyclase activity, also regulates chemotaxis. Another PH-domain-containing protein, Akt/PKB (protein kinase B), also translocates to the cell front and controls chemotaxis and cell polarity (Bagorda et al., 2006). Interestingly, PI3K,

although necessary for the chemotaxis of *Dictyostelium* in shallow chemical gradients, appears to become dispensable in steeper gradient conditions. Indeed, *Dictyostelium* cells that lack the five class I PI3Ks as well as PTEN move surprisingly well in steep gradients, suggesting that other pathways are required to control chemotaxis. Recently, phospholipase A₂ (PLA₂) and guanylyl cyclase were proposed to represent candidate pathways to regulate chemotaxis in parallel with PI3K signaling (Insall and Andrew, 2007; Kolsch et al., 2008). In addition, components of the target of rapamycin complex 2 (TORC2) have been shown to control *Dictyostelium* chemotaxis (Sasaki and Firtel, 2006), although the cellular distribution of this complex remains to be determined.

The use of Akt/PKB-GFP fusions revealed that PtdIns(3,4,5) P_3 is also enriched at the leading edge of chemotaxing neutrophils (Iijima et al., 2002; Rickert et al., 2000). In addition, neutrophils that are harvested from mice that lack PI3K γ , the principal PI3K isoform that is activated following chemoattractant stimulation, show reduced chemotaxis, although this trait appears to be environment- and chemokine-dependent (Ferguson et al., 2007; Franca-Koh et al., 2007; Liu et al., 2007; Stephens et al., 2008). Interestingly, in contrast to *Dictyostelium*, neutrophils that have been isolated from mice that lack PTEN do not show significant chemotaxis defects. Instead, in these cells, the 5'-PtdIns(3,4,5) P_3 phosphatase SHIP1 regulates chemotaxis – *ship1*^{-/-} neutrophils behave much like PTEN-deficient *Dictyostelium* cells (Nishio et al., 2007). However, the cellular distribution of SHIP1 during neutrophil chemotaxis has yet to be determined. These findings provide evidence that PtdIns(3,4,5) P_3 pathways are important regulators during mouse neutrophil chemotaxis.

Studies that were performed using HL-60 cells, a pluripotent human hematopoietic cell line that can be differentiated into neutrophil-like cells, showed that the PI3K γ -dependent PtdIns(3,4,5) P_3 enrichment at the cell front is balanced by the local phosphatidylinositol phosphate kinase I β (PIPKI β)- and PIPKI γ 661-dependent PtdIns(4,5) P_2 production at the back of chemotaxing neutrophils (Lacalle et al., 2007; Lokuta et al., 2007). Such a balance appears to be crucial for human

neutrophil chemotaxis, as abrogation of either kinase impairs neutrophil polarization and directional migration. Furthermore, the signal transduction pathway that is activated during human neutrophil chemotaxis is dependent on the type of chemoattractants. Primary chemoattractants trigger a response that depends on p38 MAPK, whereas secondary chemoattractants act via PI3K (Heit et al., 2002). In addition, and similar to *Dictyostelium*, other phospholipid signaling pathways that are mediated by PLD and PLC have been shown to impact neutrophil chemotaxis, although the exact mechanism by which this occurs remains to be determined (Kolsch et al., 2008).

The cytoskeleton and its regulators

To migrate directionally in a chemoattractant gradient, major cytoskeleton rearrangements that promote F-actin polymerization at the front and actomyosin assembly at the back of cells must occur in a coordinated fashion. Such redistributions, which are mediated by members of the Rho-family small G proteins, ultimately promote leading-edge formation at the cell front and retraction at the cell back. Actin polymerization occurs through the generation of newly formed barbed ends by the Arp2/3 complex, the activity of which is controlled by adaptor proteins of the WASP (Wiskott-Aldrich syndrome protein) and SCAR/WAVE (suppressor of cAMP receptor; also known as WAVE) families (Vartiainen and Machesky, 2004). In both *Dictyostelium* and neutrophils, SCAR/WAVE complexes are localized at the front of cells and have been shown to locally bind Rac proteins, which go on to stimulate F-actin assembly during pseudopod extension (Ibarra et al., 2005; Weiner et al., 2006). In neutrophils, Rac proteins are activated by the PH-domain-containing Rac guanine-nucleotide-exchange factors (GEFs) (P-Rex1 and Vav1) as well as DOCK2 (a DOCK180 homologue that is predominantly expressed in hematopoietic cells), the activities of which are regulated by PI3K signaling (Andrews et al., 2007; Stephens et al., 2008). In *Dictyostelium*, a single RacGEF, RacGEF1, has been identified (Park et al., 2004), although DOCK180 homologues are present in the genome. Although actin reorganization at the front of both *Dictyostelium* and neutrophils has been suggested to enhance PI3K activity, generating positive-feedback loops to reinforce polarity and directional sensing,

the mechanisms by which this takes place, remain elusive (Kolsch et al., 2008; van Haastert and Devreotes, 2004).

The retraction that occurs on the trailing edge of chemotaxing neutrophils is mediated by RhoA, which is activated by $G\alpha_{12/13}$ – another G protein that has been shown to mediate chemotactic processes, and by PDZrhoGEF (PDZ-domain-containing RhoGEF), a RhoAGEF. RhoA, through its effector ROCK (RhoA- and Rho-associated kinase), regulates myosin-II-mediated contractility (Meili and Firtel, 2003; Wong et al., 2007). The RhoA-dependent actomyosin contraction is in turn regulated by Rac (Pestonjamas et al., 2006) and Cdc42, which is activated at the front of cells by the Cdc42 GEF PIX α , and PAK1 (Li et al., 2003). In this manner Rac and Cdc42 control polarity by acting locally at the front to promote actin polymerization, and distally at the back (via RhoA) to mediate retraction (Van Keymeulen et al., 2006). In *Dictyostelium*, in which no Rho or Cdc42 homologues have been found, cell retraction is controlled by the nucleotide cGMP, which is produced by soluble guanylyl cyclase (sGC) (van Haastert and Devreotes, 2004). High cGMP levels give rise to increased myosin II phosphorylation and assembly. Myosin II assembly is also controlled by PAKa, which localizes at the back of cells. PAKa, however, is phosphorylated by Akt/PKB in a PI3K-dependent fashion (Kolsch et al., 2008). These findings highlight the importance of front-to-back crosstalk during chemotactic signaling in both *Dictyostelium* and neutrophils (Ridley et al., 2003) – a process that is also evident for the activation of adenylyl cyclase in *Dictyostelium* (see below).

Signal relay and chemotaxis amplification

During chemotaxis, *Dictyostelium* and neutrophils relay the chemoattractant signal to neighboring cells to amplify the chemotactic response. In *Dictyostelium*, CRAC – in conjuncture with TORC2 – stimulates the adenylyl cyclase ACA, which leads to the production of cAMP that is rapidly secreted to stimulate neighboring cells. During chemotaxis, ACA is highly enriched at the back of cells. This distribution is essential for the ability of *Dictyostelium* cells to align in a head-to-tail manner and migrate in the form of chains during chemotaxis (Bagorda et al., 2006). It is proposed that the ACA that is

positioned at the back of cells provides a compartment from which cAMP is locally released to attract neighboring cells to migrate in chains.

Interestingly, in neutrophils, the release of purines (such as ATP) has been shown to occur from the leading edge of the cell (Chen et al., 2006). As a neutrophil moves towards a source of chemoattractant, ATP is released and, together with its derived adenosine, provides an autocrine feedback loop by activating transmembrane purinergic receptors, thereby amplifying chemotaxis. Mammalian neutrophils also release a variety of chemokines during migration towards sites of inflammation. For example, chemoattractants such as fMLP and C5a stimulate the production and secretion of LTB₄, as well as interleukin-8. Although not much is known about the mechanisms that guide the secretion of chemokines, there is increasing evidence that signal relay is responsible for the amplification of chemotaxis and is a key regulator of inflammatory responses.

Perspectives

In the past 10 years, incredible progress has been made in understanding how cells sense and respond to chemotactic gradients. The use of GFP technology has revolutionized the field and allowed scientists to get a glimpse of the intracellular working of live migrating cells (Van Haastert and Devreotes, 2004). Although the PI3K pathway is highly polarized in migrating cells and has a role in regulating chemotaxis, other parallel pathways are clearly involved. The challenge now resides in identifying additional pathways and, most importantly, in assessing how they work in concert to control polarity and chemotaxis. In this context, powerful *Dictyostelium* genetic screens will be invaluable and continue to provide key insight. However, as neutrophils are exposed to a complex array of chemotactic agents, more elaborate signaling cascades are surely operational in higher eukaryotes. Continued studies involving gene manipulation in mice as well as specific pharmacological inhibitors will certainly bring significant insight.

We apologize to our colleagues in the field for only citing recent reviews and articles. The authors thank the members of the Parent laboratory for insightful discussions and for their helpful comments on the manuscript. We are grateful to the Intramural Research Program of the National Institutes of Health, National Cancer Institute, Center for Cancer Research

for support. A.B. is the recipient of an Italian Cancer Federation Fellowship.

References

- Andrews, S., Stephens, L. R. and Hawkins, P. T. (2007). PI3K class IB pathway in neutrophils. *Sci. STKE* 2007, cm3.
- Baggiolini, M. (2001). Chemokines in pathology and medicine. *J. Intern. Med.* **250**, 91-104.
- Bagorda, A., Mihaylov, V. A. and Parent, C. A. (2006). Chemotaxis: moving forward and holding on to the past. *Thromb. Haemost.* **95**, 12-21.
- Brzostowski, J. A. and Kimmel, A. R. (2001). Signaling at zero G: G-protein-independent functions for 7-TM receptors. *Trends Biochem. Sci.* **26**, 291-297.
- Chen, Y., Corriden, R., Inoue, Y., Yip, L., Hashiguchi, N., Zinkernagel, A., Nizet, V., Insel, P. A. and Junger, W. G. (2006). ATP release guides neutrophil chemotaxis via P2Y2 and A3 receptors. *Science* **314**, 1792-1795.
- Chisholm, R. L. and Firtel, R. A. (2004). Insights into morphogenesis from a simple developmental system. *Nat. Rev. Mol. Cell Biol.* **5**, 531-541.
- Ferguson, G. J., Milne, L., Kulkarni, S., Sasaki, T., Walker, S., Andrews, S., Crabbe, T., Finan, P., Jones, G., Jackson, S. et al. (2007). PI(3)K has an important context-dependent role in neutrophil chemokinesis. *Nat. Cell Biol.* **9**, 86-91.
- Franca-Koh, J., Kamimura, Y. and Devreotes, P. N. (2007). Leading-edge research: PtdIns(3,4,5)P₃ and directed migration. *Nat. Cell Biol.* **9**, 15-17.
- Heit, B., Tavener, S., Raharjo, E. and Kubers, P. (2002). An intracellular signaling hierarchy determines direction of migration in opposing chemotactic gradients. *J. Cell Biol.* **159**, 91-102.
- Ibarra, N., Pollitt, A. and Insall, R. H. (2005). Regulation of actin assembly by SCAR/WAVE proteins. *Biochem. Soc. Trans.* **33**, 1243-1246.
- Iijima, M., Huang, Y. E. and Devreotes, P. (2002). Temporal and spatial regulation of chemotaxis. *Dev. Cell* **3**, 469-478.
- Insall, R. and Andrew, N. (2007). Chemotaxis in *Dictyostelium*: how to walk straight using parallel pathways. *Curr. Opin. Microbiol.* **10**, 578-581.
- Kolsch, V., Charest, P. G. and Firtel, R. A. (2008). The regulation of cell motility and chemotaxis by phospholipid signaling. *J. Cell Sci.* **121**, 551-559.
- Lacalle, R. A., Peregil, R. M., Albar, J. P., Merino, E., Martinez, A. C., Merida, I. and Manes, S. (2007). Type I phosphatidylinositol 4-phosphate 5-kinase controls neutrophil polarity and directional movement. *J. Cell Biol.* **179**, 1539-1553.
- Li, Z., Hannigan, M., Mo, Z., Liu, B., Lu, W., Wu, Y., Smrcka, A. V., Wu, G., Li, L., Liu, M. et al. (2003). Directional sensing requires G $\beta\gamma$ -mediated PAK1 and PIX α -dependent activation of Cdc42. *Cell* **114**, 215-227.
- Liu, L., Puri, K. D., Penninger, J. M. and Kubers, P. (2007). Leukocyte PI3K γ and PI3K δ have temporally distinct roles for leukocyte recruitment in vivo. *Blood* **110**, 1191-1198.
- Lokuta, M. A., Senetar, M. A., Bennin, D. A., Nuzzi, P. A., Chan, K. T., Ott, V. L. and Huttenlocher, A. (2007). Type I gamma PIP kinase is a novel uropod component that regulates rear retraction during neutrophil chemotaxis. *Mol. Biol. Cell* **18**, 5069-5080.
- Mahadeo, D. C. and Parent, C. A. (2006). Signal relay during the life cycle of *Dictyostelium*. *Curr. Top. Dev. Biol.* **73**, 115-140.
- Manahan, C. L., Iglesias, P. A., Long, Y. and Devreotes, P. N. (2004). Chemoattractant signaling in *Dictyostelium discoideum*. *Annu. Rev. Cell Dev. Biol.* **20**, 223-253.
- Manes, S., Gomez-Mouton, C., Lacalle, R. A., Jimenez-Baranda, S., Mira, E. and Martinez, A. C. (2005). Mastering time and space: immune cell polarization and chemotaxis. *Semin. Immunol.* **17**, 77-86.
- Meili, R. and Firtel, R. A. (2003). Two poles and a compass. *Cell* **114**, 153-156.
- Milligan, G. and Kostenis, E. (2006). Heterotrimeric G-proteins: a short history. *Br. J. Pharmacol.* **147**, Suppl. 1, S46-S55.
- Niggli, V. (2003). Signaling to migration in neutrophils: importance of localized pathways. *Int. J. Biochem. Cell Biol.* **35**, 1619-1638.
- Nishio, M., Watanabe, K., Sasaki, J., Taya, C., Takasuga, S., Iizuka, R., Balla, T., Yamazaki, M., Watanabe, H., Itoh, R. et al. (2007). Control of cell polarity and motility by the PtdIns(3,4,5)P₃ phosphatase SHIP1. *Nat. Cell Biol.* **9**, 36-44.
- Parent, C. A. (2004). Making all the right moves: chemotaxis in neutrophils and *Dictyostelium*. *Curr. Opin. Cell Biol.* **16**, 4-13.
- Parent, C. A. and Devreotes, P. N. (1996). Molecular genetics of signal transduction in *Dictyostelium*. *Ann. Rev. Biochem.* **65**, 411-440.
- Parent, C. A. and Devreotes, P. N. (1999). A cell's sense of direction. *Science* **284**, 765-770.
- Park, K. C., Rivero, F., Meili, R., Lee, S., Apone, F. and Firtel, R. A. (2004). Rac regulation of chemotaxis and morphogenesis in *Dictyostelium*. *EMBO J.* **23**, 4177-4189.
- Pestonjamasp, K. N., Forster, C., Sun, C., Gardiner, E. M., Bohl, B., Weiner, O., Bokoch, G. M. and Glogauer, M. (2006). Rac1 links leading edge and uropod events through Rho and myosin activation during chemotaxis. *Blood* **108**, 2814-2820.
- Rickert, P., Weiner, O. D., Wang, F., Bourne, H. R. and Servant, G. (2000). Leukocytes navigate by compass: roles of PI3K γ and its lipid products. *Trends Cell Biol.* **10**, 466-473.
- Ridley, A. J., Schwartz, M. A., Burridge, K., Firtel, R. A., Ginsberg, M. H., Borisy, G., Parsons, J. T. and Horwitz, A. R. (2003). Cell migration: integrating signals from front to back. *Science* **302**, 1704-1709.
- Saran, S., Meima, M. E., Alvarez-Curto, E., Weening, K. E., Rozen, D. E. and Schaap, P. (2002). cAMP signaling in *Dictyostelium*. Complexity of cAMP synthesis, degradation and detection. *J. Muscle Res. Cell Motil.* **23**, 793-802.
- Sasaki, A. T. and Firtel, R. A. (2006). Regulation of chemotaxis by the orchestrated activation of Ras, PI3K, and TOR. *Eur. J. Cell Biol.* **85**, 873-895.
- Stephens, L., Milne, L. and Hawkins, P. (2008). Moving towards a better understanding of chemotaxis. *Curr. Biol.* **18**, R485-R494.
- van Haastert, P. J. and Devreotes, P. N. (2004). Chemotaxis: signaling the way forward. *Nat. Rev. Mol. Cell Biol.* **5**, 626-634.
- Van Keymeulen, A., Wong, K., Knight, Z. A., Govaerts, C., Hahn, K. M., Shokat, K. M. and Bourne, H. R. (2006). To stabilize neutrophil polarity, PIP3 and Cdc42 augment RhoA activity at the back as well as signals at the front. *J. Cell Biol.* **174**, 437-445.
- Vartiainen, M. K. and Machesky, L. M. (2004). The WASP-Arp2/3 pathway: genetic insights. *Curr. Opin. Cell Biol.* **16**, 174-181.
- Weiner, O. D., Rentel, M. C., Ott, A., Brown, G. E., Jedrychowski, M., Yaffe, M. B., Gygi, S. P., Cantley, L. C., Bourne, H. R. and Kirschner, M. W. (2006). Hem-1 complexes are essential for Rac activation, actin polymerization, and myosin regulation during neutrophil chemotaxis. *PLoS Biol.* **4**, e38.
- Wong, K., Van Keymeulen, A. and Bourne, H. R. (2007). PDZRhoGEF and myosin II localize RhoA activity to the back of polarizing neutrophil-like cells. *J. Cell Biol.* **179**, 1141-1148.

Cell Science at a Glance on the Web
Electronic copies of the poster insert are available in the online version of this article at jcs.biologists.org. The JPEG images can be downloaded for printing or used as slides.