

Proteins that bind A-type lamins: integrating isolated clues

Michael S. Zastrow, Sylvia Vlcek and Katherine L. Wilson*

Department of Cell Biology, Johns Hopkins University School of Medicine, 725 N. Wolfe Street, Baltimore, MD 21205, USA

*Author for correspondence (e-mail: klwilson@jhmi.edu)

Journal of Cell Science 117, 979-987 Published by The Company of Biologists 2004
doi:10.1242/jcs.01102

Summary

What do such diverse molecules as DNA, actin, retinoblastoma protein and protein kinase C α all have in common? They and additional partners bind 'A-type' lamins, which form stable filaments in animal cell nuclei. Mutations in A-type lamins cause a bewildering range of tissue-specific diseases, termed 'laminopathies', including Emery-Dreifuss muscular dystrophy and the devastating Hutchinson-Gilford progeria syndrome, which mimics premature aging. Considered individually and collectively, partners for A-type lamins form four loose groups: architectural partners, chromatin partners, gene-regulatory partners and signaling partners. We describe 16 partners in detail, summarize their binding sites in A-type

lamins, and sketch portraits of ternary complexes and functional pathways that might depend on lamins *in vivo*. On the basis of our limited current knowledge, we propose lamin-associated complexes with multiple components relevant to nuclear structure (e.g. emerin, nesprin 1 α , actin) or signaling and gene regulation (e.g. LAP2 α , retinoblastoma, E2F-DP heterodimers, genes) as 'food for thought'. Testing these ideas will deepen our understanding of nuclear function and human disease.

Key words: Laminopathy, Emerin, Emery-Dreifuss muscular dystrophy, Hutchinson-Gilford progeria syndrome, Nuclear envelope

Introduction


Lamins are intermediate filament proteins found only in the nuclei of multicellular eukaryotes. They form stable filaments at the nuclear inner membrane and stable structures in the nucleoplasm that might or might not be filamentous. Lamins are important for nuclear architecture; they provide mechanical strength, determine nuclear shape, and anchor and space the nuclear pore complexes. Perhaps more surprisingly, lamins are also essential for DNA replication and mRNA transcription (Goldman et al., 2002). Mammalian cells encode both A-type and B-type lamins, which are highly related but can be distinguished biochemically and functionally. The most crucial distinction is that B-type lamins, encoded by two human genes (*LMNB1* and *LMNB2*), are essential for viability of individual cells, whereas A-type lamins (encoded by *LMNA*) are dispensable. Loss of either lamin B1 or lamin B2 is lethal to dividing cells (Harborth et al., 2001), which is consistent with fundamental roles in replication and transcription. However, the basis for the dependence of metazoans on B-type lamins is still unknown. A-type lamins, although dispensable during development in mice, have key roles in differentiated tissues. Humans synthesize four A-type lamins (A, C, C2 and A Δ 10), which are generated by alternative mRNA splicing of the *LMNA* gene (Fisher et al., 1986; Furukawa et al., 1994; Machiels et al., 1996; McKeon et al., 1986). A-type lamins begin to be expressed in a tissue-specific manner during development, and cell-type-appropriate isoforms of A-type lamins continue to be expressed in most differentiated adult cells (Machiels et al., 1996; Rober et al., 1989). A-type lamins appeared late in metazoan evolution; they are not found in the nematode *Caenorhabditis elegans*, which has differentiated

cells and tissues, but are found in more complicated eukaryotes, including *Drosophila*, birds and vertebrates (Cohen et al., 2001; Zimek et al., 2003). Despite their clearly different roles, the conserved similarities between A- and B-type lamins suggest similar mechanisms of function in the nucleus.

Like other intermediate filament proteins, lamins consist of an N-terminal head domain, a central coiled-coil (rod) domain responsible for dimerization, and a large globular C-terminal tail. The tail includes an immunoglobulin (Ig)-fold domain, the backbone organization of which is unique to lamins, formed by residues 430-545 (Dhe-Paganon et al., 2002; Krimm et al., 2002). All lamins except lamin C and lamin C2 also contain a CAAX (Cys-aliphatic-aliphatic-any residue) motif at the C-terminus, which is farnesylated on Cys *in vivo* prior to proteolytic removal of the AAX sequence. Prelamin A is then carboxymethylated and re-cleaved to remove the modified Cys plus 14 additional residues (647-661), yielding mature lamin A (Fig. 1) (Moir et al., 1995). A site-specific protease named Zmpste24 is proposed to perform both cleavage events (Pendas et al., 2002). Lamin A is (so far) the only known substrate for Zmpste24 in mammals. Consistent with lamin A being a major substrate is the observation that loss of Zmpste24 causes muscular dystrophy and bone degeneration phenotypes in mice (Bergo et al., 2002; Pendas et al., 2002; see below) and mandibuloacral dysplasia in humans (Agarwal et al., 2003), disorders associated with compromised A-type lamin function.

The assembly mechanisms and structures of lamin filaments are reviewed elsewhere (Goldman et al., 2002; Stuurman et al., 1998). However, several properties are noteworthy. First, the abundance and organization of lamin filaments vary between

Fig. 1. Regions in lamins A and C to which partners bind. The structural domains (head, rod, tail) of prelamins A are shown. Exons 1-12 encoding residues 1-664 are numbered; the last residue encoded by each exon (except exon 11) is given above. The Ig-fold domain, which includes exons 8 and 9, is shaded gray. Residues 1-566 of lamin A and lamin C are identical. Unique lamin C tail residues 567-572 are produced by alternative mRNA splicing; thus the extreme C-terminal regions of lamins A and C may have distinct binding properties. The zigzag represents the farnesyl moiety on prelamins A; the farnesylated C-terminal peptide is normally removed by proteolytic cleavage after residue 646 (bold; dotted line) to generate mature lamin A. Colored bars indicate the region(s) in lamins A and C required for each named partner to bind, as detailed in Table 1 (e.g. actin can bind two different regions in the tail). For partners with question marks, the binding region in lamin A/C is unmapped. Based on current incomplete knowledge, interactions were loosely color-coded (top to bottom) as blue (architectural), orange (chromatin), yellow (gene regulation), pink (signaling) and green (unknown). Some partners (e.g. BAF) will continue to defy categorization until more is known about their functions.



species and cell types. In *Xenopus* oocyte nuclei, lamins form square networks of 10 nm filaments, all of which are tightly associated with the inner membrane (Stuurman et al., 1998). By contrast, in mammalian somatic nuclei, a significant number of lamins are found in the nuclear interior (Moir et al., 2000). The ultrastructure of interior lamins is unknown, except that they are unlikely to form 10 nm filaments. Second, although A-B heterodimers can form in vitro (Georgatos et al., 1988; Schirmer et al., 2001; Ye and Worman, 1995), there is growing evidence that A-type and B-type lamins form independent networks in vivo (Izumi et al., 2000; Moir et al., 2000; Haraguchi et al., 2001). Finally, and most interestingly from our current viewpoint, lamin polymers comprise a three-dimensional network that displays at least a million copies of each binding site in the exposed rod and tail domains of each lamin, per nucleus. (HeLa cell nuclei contain $\sim 10^6$ copies each of lamins A, B1 and C) (L. Gerace, personal communication). Most biochemical studies have measured partner binding to lamin dimers, not polymers; thus the corresponding interaction in vivo could have increased avidity owing to the large number of available sites on lamin polymers. It is also possible that lamins form a variety of distinct oligomers or polymers that have distinct 'docking' properties. The variety of human diseases linked to A-type lamins suggests that these sites, and

the partners that bind to them, are important for nuclear function.

Laminopathies

At least seven diseases are caused by mutations in *LMNA* (Burke and Stewart, 2002; Östlund and Worman, 2003). Each disease affects specific combinations of tissues such as the heart, skeletal muscle and tendons (Emery-Dreifuss muscular dystrophy; EDMD), skeletal muscle and heart (limb-girdle muscular dystrophy 1B), the heart only (dilated cardiomyopathy type 1A), neurons and muscle (Charcot-Marie-Tooth disorder type 2B1), adipocytes (Dunnigan-type familial partial lipodystrophy), or adipocytes and bone (mandibuloacral dysplasia). The most devastating laminopathy is Hutchinson-Gilford progeria syndrome (HGPS), which affects multiple tissues and resembles premature aging (De Sandre-Giovannoli et al., 2003; Eriksson et al., 2003; Chen et al., 2003). Patients who have this disease usually die in their teens. Whereas most laminopathies are caused by missense mutations spread throughout the molecule, the most frequent known cause of HGPS is a point mutation that creates a new splice donor site, resulting in the deletion of 50 residues (608-657), including the proteolytic cleavage site at residue 646.

Table 1. Current knowledge of binding regions in A-type lamins and each partner

Partner	Binding region in partner	Binding region in lamin A/C
Actin	ND*	Lamins A and C residues 461-536 (Zastrow et al., unpublished) Lamin A residues 563-646 (Sasseville and Langelier, 1998)
BAF	ND	ND
DNA	Minor groove	Lamins A and C residues 411-553 (Stierle et al., 2003)
E1B 19K	Residues 1-146	Lamins A and C residues 252-390 (Rao et al., 1997)
Emerin	Residues 70-170	Lamins A and C residues 384-566 (Lee et al., 2001; Sakaki et al., 2001)
Histones	ND	Lamin C residues 396-430 (Taniura et al., 1995)
Lamin B	ND	ND
LAP1	ND	ND
LAP2 α	Residues 616-693	Lamins A and C residues 319-566 (Dechat et al., 2000)
12(S)-LOX	ND	Lamin A residues 463-664 (Tang et al., 2000)
MOK2	Residues 1-173	Lamins A and C residues 243-387 (Dreuillet et al., 2002)
Narf	ND	Farnesylated prelamin A residues 389-664 (Barton and Worman, 1999)
Nesprin 1 α	Residues 476-1109	ND (Mislow et al., 2002a)
PKC α	Residues 200-217 Residues 660-672	Lamin A residues 500-664 (Martelli et al., 2002)
Rb	Residues 792-928 Residues 612-928	ND for lamin C (Markiewicz et al., 2002) Lamins A and C residues 247-355 (Ozaki et al., 1994)
SREBP	Residues 227-487	Lamin A residues 389-664 (Lloyd et al., 2002)

*ND, not determined.

Thus, HGPS patients are predicted to produce an internally deleted form of the farnesylated and carboxymethylated lamin A precursor. *Zmpste24*-null mice lack the processing protease and also have bone- and muscle-disorder phenotypes (Bergo et al., 2002). This suggests that severe laminopathy, if not progeria, might correlate with the presence of unprocessed lamin A precursors. Interestingly, a protein named nuclear prelamin A recognition factor (Narf), which is expressed at its highest levels in heart, skeletal muscle and brain, localizes to the nuclear envelope and intranuclear foci and binds specifically to farnesylated lamin A precursors (Barton and Worman, 1999). Thus, these forms of lamin A have the potential to saturate Narf, the function of which is unknown. When expressed exogenously, disease-associated lamins can disturb the organization of endogenous lamins (e.g. Raharjo et al., 2001; Capanni et al., 2003). In the few cases studied, laminopathy mutations can disrupt binding of lamins to specific partners (see below).

The first genetic model for laminopathy was the *LMNA*-knockout mouse; these mice appear normal at birth but develop severe muscular dystrophy and die within eight weeks (Sullivan et al., 1999). In the first reported human *LMNA*-null case, a baby possessing a homozygous nonsense mutation in *LMNA*, with no detectable A-type lamin proteins, died after premature birth, exhibiting severe joint contractures, muscular dystrophy, fibrosis and absence of muscle fibers in the diaphragm (Muchir et al., 2003). This premature birth and death cannot be attributed exclusively to loss of A-type lamins, because of parental consanguinity and the consequent potential for homozygous defects in other genes. Nonetheless, the resemblance to laminopathy suggests that A-type lamins, although not essential for cell viability, are essential for human life.

Mice have been constructed to express only the L530P-mutated forms of lamins A and C, which causes EDMD in humans (Mounkes et al., 2003). A side-effect of the targeting strategy was that the 'knocked-in' L530P allele also changed the mRNA splicing pattern to yield two predicted protein products: mature A-type lamins that lack exon 9, and a C-

terminally-truncated lamin A that has 19 extra (intron-encoded) internal residues (see supplementary data in Mounkes et al., 2003). Neither mutation is expected to disrupt C-terminal processing of lamins, but this possibility has not been ruled out. These mice have 'progeria-like' pathologies of bone and skin, and die prematurely (Mounkes et al., 2003). Loss of human *Zmpste24* gene activity causes severe mandibuloacral dysplasia with progeroid appearance and general lipodystrophy (Agarwal et al., 2003), which is altogether less severe than HGPS. Thus, large deletions in the lamin A/C tail (i.e. loss of 50 residues in HGPS, or loss of exon 9 residues in L530P-knock-in mice) might correlate with progeria, and must certainly disrupt lamin organization and interactions. The effects of 'progeric' mutations on the assembly and partner-binding activities of A-type lamins will be important to test.

Many plausible disease mechanisms have been proposed. Some models emphasize possible disruption of the mechanical properties of lamin filaments, which control the shape, stiffness and structural integrity of nuclei. Other models view lamins as sites ('scaffolds') for the assembly of other proteins that regulate transcription, cell fate or apoptosis (Burke and Stewart, 2002; Goldman et al., 2002). Both models are probably correct. We have taken a fresh look at the published binding partners for A-type lamins (Table 1, Fig. 1). Some bind A-type lamins in two-hybrid assays and in vitro, but their in vivo significance remains untested. In other cases, the interactions have also been studied in vivo. Many partners were familiar to us, but others were not. They form four loose groups: architectural partners, chromatin partners, gene-regulatory partners and signaling partners. Clearly, some partners fit multiple categories. What emerges, however, is that a rich variety of protein complexes and biochemical pathways could depend on lamins in vivo.

Architectural partners

We loosely defined architectural partners as those that (1) form

stable complexes during interphase (on the basis of their resistance to chemical extraction or their immobility in FRAP experiments), (2) bind tightly to A-type lamins *in vitro*, or (3) have known structural roles in other contexts (e.g. actin). Such partners might connect lamins to the nuclear envelope, chromatin or other subnuclear structures (Burke and Stewart, 2002; Cohen et al., 2001). All integral membrane proteins that bind to lamins fit this definition, including lamina-associated polypeptide 1 (LAP1) and emerin (Foisner and Gerace, 1993; Östlund et al., 1999). LAP1 has an unusually large luminal domain (Foisner and Gerace, 1993; Martin et al., 1995) and is expressed as multiple isoforms but has not been further characterized. More information is available for emerin, which is relatively immobile in the nuclear inner membrane (Östlund et al., 1999) and binds lamin A with an affinity of 40 nM *in vitro* (Holaska et al., 2003) (reviewed by Bengtsson and Wilson, 2004). Interestingly, emerin also binds directly to barrier-to-autointegration factor (BAF), a DNA-binding chromatin protein, and might thereby anchor lamins and chromatin to the nuclear envelope during interphase (Lee et al., 2001). Other nuclear membrane proteins (e.g. LAP2 β) bind to B-type lamins, BAF, DNA and chromatin, extending this architectural theme (Cai et al., 2001; Foisner and Gerace, 1993; Furukawa, 1999; Shumaker et al., 2001). Emerin and LAP2 β both belong to the LEM-domain family of nuclear proteins (Lin et al., 2000). The defining feature of LEM-domain proteins is their LEM domain: a folded motif of at least 40 residues that binds directly to BAF (Zheng et al., 2000; Cai et al., 2001). The LEM-domain family in vertebrates also includes at least six splice forms of LAP2 (Berger et al., 1996), MAN1 (which binds Smad transcription factors and regulates development) (Osada et al., 2003; Raju et al., 2003) and a smaller MAN1-related protein provisionally named Lem2 (Lee and Wilson, 2004), which is conserved in metazoans and essential for viability in *C. elegans* embryos (Lee et al., 2000; Liu et al., 2003). Lem2 (also known as NET-25) was identified in a proteomic study that also revealed more than 60 additional potential nuclear membrane proteins (Schirmer et al., 2003). We assume that many of these novel proteins bind A-type lamins. Further work will undoubtedly increase our appreciation of the complexity of the nuclear lamina network and possibly also extend its range of known functions.

The best-studied architectural partner for A-type lamins in the nuclear interior is LAP2 α , which has separate binding sites for BAF, A-type lamins and chromatin (Dechat et al., 1998; Dechat et al., 2000; Vlcek et al., 1999). LAP2 α binds tightly to A-type lamins during interphase (Dechat et al., 2000), but has dynamic architectural roles during nuclear envelope assembly (Vlcek et al., 2002). It is proposed to tether A-type lamins to intranuclear sites and to cooperate with lamins in organizing chromatin (Foisner, 2003). LAP2 α (and other LEM-domain proteins) also binds transcriptional regulators (see below), establishing a theme in which 'architectural' partners for lamins also influence gene regulation, directly or indirectly.

We do not yet know whether B-type lamins are direct architectural partners for A-type lamins *in vivo*. A- and B-type lamins have distinct assembly pathways *in vivo* (Moir et al., 2000), which could mean they form distinct networks. However, B-type lamins might 'set the stage' for the assembly of A-type lamins. A-type lamins are known to be architectural

partners for each other, since lamin C depends on lamin A for its localization and assembly in the nucleus (Vaughan et al., 2001).

Actin, actin-related proteins and numerous actin-binding proteins (including a nuclear-specific isoform of myosin I) are present in the nucleus, where their functions are now slowly emerging (Pederson and Aebi, 2002). There is currently no evidence for long actin filaments (F-actin) in the nucleus. However, actin can also form a multitude of special dimers, short protofilaments and tubular, flat or branched oligomers that would suit the chromatin-dominated nuclear space (Pederson and Aebi, 2002). Interestingly, nuclear actin polymers adopt a unique conformation that is recognized by specific antibodies (Milankov and De Boni, 1993). Actin binds directly to two regions in the lamin A/C tail: residues 461-536 (in the Ig-fold domain) (M.S.Z. and K.L.W., unpublished) and residues 563-646 (Sasseville and Langelier, 1998). A-type lamin filaments might thus bind to actin polymers in the nucleus. The possibility that actin polymers are architectural partners for lamin filaments deserves serious attention.

Nesprin 1 α , a 131 kDa nuclear membrane protein, binds directly to A-type lamins and emerin (Mislow et al., 2002a; Mislow et al., 2002b). All nesprin-family proteins (also known as Syne, Myne and NUANCE) have multiple spectrin repeat (SR) domains, and many also have an actin-binding domain (Zhang et al., 2002; Zhang et al., 2001; Starr and Han, 2003). Human nesprins are encoded by two genes, which yield multiple protein isoforms through alternative mRNA splicing and use of internal transcription initiation sites. These isoforms range in mass from huge (~1 MDa for nesprin 1) to modest (e.g. 131 kDa for nesprin 1 α) (Mislow et al., 2002a; Zhang et al., 2002; Zhang et al., 2001). Nesprins are anchored to specific membranes, such as the Golgi complex or the outer or inner membrane of the nuclear envelope (Gough et al., 2003; Mislow et al., 2002b; Zhang et al., 2001). Nesprin 1 α was proposed to anchor lamin A and emerin at the inner nuclear membrane. However, this model was overthrown by the finding that nesprin 1 α and emerin both mislocalize to the endoplasmic reticulum (ER) in human fibroblasts that lack A-type lamins (Muchir et al., 2003). Transient expression of either lamin A or lamin C is sufficient to relocalize both membrane proteins to the nuclear envelope. These results suggest an architectural hierarchy based on the integrity of A-type lamins.

Chromatin partners

Lamins bind DNA directly but nonspecifically *in vitro*, by contacting the minor groove (Luderus et al., 1994). The DNA-binding region is identical in lamin A and lamin C, and includes both the Ig-fold domain and the nuclear localization signal in the tail (Stierle et al., 2003). The affinity of lamin A for DNA is 420 nM (Stierle et al., 2003). The R482W missense mutation, which causes lipodystrophy, reduces this affinity for DNA by seven-fold (to 2.95 μ M), and also weakens binding of lamin A to the adipocyte differentiation factor sterol response element-binding protein (SREBP; see below).

DNA is wound around core histone octamers to form nucleosomes, the fundamental unit of chromatin structure. The human lamin A/C tail binds to mixtures of core histones with an apparent affinity of 300 nM (Taniura et al., 1995), which is similar to its affinity for naked DNA. Since histones and DNA

are abundant in the nucleus, given such an affinity A-type lamins are probably always close to chromatin. This would allow lamins to serve as scaffolds for multiprotein complexes associated with chromatin.

BAF is essential for cell viability in *C. elegans* (Zheng et al., 2000) and *Drosophila* (Furukawa et al., 2003). It binds to lamin A with 1 μ M affinity in vitro (Holaska et al., 2003) and also binds to LEM-domain proteins (Furukawa, 1999; Lee et al., 2001; Shumaker et al., 2001; Liu et al., 2003). However, BAF also qualifies as a 'chromatin' partner for lamins because it binds nonspecifically to double-stranded DNA (Zheng et al., 2000; Shumaker et al., 2001) and co-localizes with chromatin in certain cell types (e.g. *Drosophila* embryos) (Furukawa et al., 2003). Live-cell studies using GFP-fused BAF show direct binding to emerin at the nuclear envelope but also show that GFP-BAF is highly mobile during interphase (Shimi et al., 2004). In *Drosophila* embryos, loss of BAF causes chromatin and nuclear envelope defects (Furukawa et al., 2003). Additional mitotic defects seen in these embryos (e.g. lack of cyclin expression) might reflect a role for BAF in gene expression (see below). Nonetheless, excess BAF present during nuclear assembly can profoundly alter chromatin structure in *Xenopus* egg extracts (Segura-Totten et al., 2002), confirming its importance as an architectural partner but not revealing the mechanism. We can only speculate that BAF serves as a chromatin partner for lamin A by rapidly interlinking chromatin, LEM-domain proteins and lamins.

Gene-regulatory partners

Lamin A/C filaments are thought to provide scaffolds for protein complexes that regulate gene expression (Cohen et al., 2001). Hypophosphorylated retinoblastoma protein (Rb) binds specifically to A-type lamins in vitro (Mancini et al., 1994; Markiewicz et al., 2002; Ozaki et al., 1994). Rb is a transcriptional regulator that has central roles in cell-cycle control. Hypo-phosphorylated Rb binds to E2F-DP heterodimers and blocks E2F-dependent gene expression through a variety of mechanisms, including the recruitment of histone deacetylase complexes (HDACs) (Chau and Wang, 2003). Rb also regulates expression of apoptosis protease-activating factor 1 (Apaf1) and thus apoptosis, which is proposed to contribute to muscle loss in EDMD patients (Bonne et al., 2003). During G1 phase, hypophosphorylated (repressive) Rb is anchored to the nucleoskeleton by its 'pocket-C' domain, which binds directly to A-type lamins and LAP2 α (Mancini et al., 1994; Markiewicz et al., 2002). Overexpression of LAP2 α interferes with cell-cycle progression and leads to apoptosis (Vlcek et al., 2002), which supports 'scaffolding' models in which Rb activity depends on its attachment to lamins and LAP2 α (Fig. 2). Such complexes might be weakened by disease-causing mutations, because Rb and LAP2 α interact with the rod and rod/tail domains, respectively, of lamins A and C (Fig. 1). Biochemical studies using a different repressor of E2F activity, germ-cell-less (GCL), independently support scaffolding models, showing that GCL, emerin and lamin A form stable tertiary complexes in vitro (Holaska et al., 2003). Indeed, stable, lamin-based scaffolds might be useful to tether the enormous (1-2 MDa) chromatin-remodeling machines

recruited by Rb and other regulators (Neely and Workman, 2002) (Fig. 2).

SREBP1a and SREBP1c are additional lamin-binding proteins. Encoded by alternatively spliced mRNAs, they are both basic helix-loop-helix leucine zipper transcription factors that, when activated, are released from the ER by proteolytic cleavage and move into the nucleus. Once in the nucleus, they activate genes required for cholesterol biosynthesis and lipogenesis (reviewed by Horton, 2002) and promote adipocyte differentiation (Kim and Spiegelman, 1996). SREBP proteins bind to the Ig-fold domain of the lamin A/C tail, as do many other partners (Fig. 1). Interestingly, lipodystrophy-causing mutations in A-type lamins reduce SREBP binding by 25-40% (Lloyd et al., 2002). It will be important to determine which other partners (in addition to SREBP and DNA) are affected by lipodystrophy-causing mutations.


MOK2 is a DNA-binding transcriptional repressor that interacts with lamins A and C. It represses genes activated by cone-rod homeobox protein (Crx) transcription factors by competing with them for binding sites (Dreuillet et al., 2002). MOK2 also binds to RNA in vitro, and might thus influence RNA processing (Arranz et al., 1997). Like Rb, MOK2 binds to the coil 2 region of A-type lamins (Dreuillet et al., 2002) (Fig. 1). It therefore seems that A-type lamins have binding site(s) for transcriptional repressors not only in their tail domain, but also in the rod domain. Optimal access to rod domain sites might require a specific oligomeric (dimer, tetramer or octamer) or assembly (filamentous) state.

Another gene-regulatory partner is BAF (see above). BAF binds directly to several different homeodomain transcription activators, including Crx, and represses Crx-dependent gene expression in retinal cells (Wang et al., 2002) (reviewed by Segura-Totten and Wilson, 2004). Thus, BAF and MOK2 both repress CRX-activated genes, but by different mechanisms (Fig. 2). It is curious that two out of four identified lamin-binding repressors influence Crx-regulated genes, but this might simply reflect the low number of proteins studied so far.

The BAF-binding partner emerin is proposed to be a fifth gene-regulatory partner for A-type lamins. Other LEM-domain proteins (e.g. LAP2 β and MAN1) that bind lamin B inhibit gene expression in vivo (Liu et al., 2003; Nili et al., 2001; Osada et al., 2003). Emerin has not yet been tested for such activity in vivo. However, it does form a complex with lamin A and the transcriptional repressor GCL in vitro (Holaska et al., 2003). Emerin is very special among known binding partners for lamin A/C, because loss of emerin causes the same disease – EDMD – as many dominant mutations in A-type lamins (Bione et al., 1994). Perhaps in future other laminopathies will be analogously 'paired' to loss of a specific lamin-binding protein(s). Meanwhile the wide expression of emerin in human tissues, and functional overlap between emerin and other LEM-domain proteins make it difficult to pinpoint which function(s) of emerin are critical to the EDMD disease mechanism (Bengtsson and Wilson, 2004). Further molecular and in vivo dissection of emerin functions is needed to clarify its role in lamin A/C complexes, which might combine roles in nuclear architecture (Fig. 2) and gene regulation.

Various gene regulators thus bind A-type lamins and/or emerin, and this supports the idea that dysregulation of gene expression is an important contributing factor in laminopathies.

Fig. 2. Speculative cartoon to 'integrate isolated clues', and stimulate thought. A few interactions described for A-type lamins in the text are illustrated here, speculatively, in the context of lamin filaments near the nuclear inner membrane (IM) or associated with chromatin. Integral membrane proteins nesprin 1 α (Nesprin) and emerin (EM) are embedded at the IM. Many other lamin-binding IM proteins are not depicted. NPC, nuclear pore complex. NS, nucleosome. OM, nuclear outer membrane. (Top left) Emerin, representing all LEM-domain proteins, binds BAF. BAF and MOK bind lamins *in vitro* and inhibit Crx-dependent gene activation *in vivo*, implicating both proteins in lamin-dependent gene-regulation events. Not depicted are potential gene-regulatory complexes involving nuclear membrane proteins LAP2 β (Nili et al., 2001; Foisner, 2003) or lamin B receptor (Östlund and Worman, 2003).



Emerin has many direct binding partners including nesprin-1 α , lamins, actin (not depicted), BAF and several transcription regulators (not shown) (Bengtsson and Wilson, 2004). The number of distinct oligomeric complexes that include emerin is not known. (Top right) Lamin-binding enzyme 12(S)-lipoxygenase [12(S)-LOX] converts arachidonic acid (AA) to 12(S)-hydroxyeicosatetraenoic acid [12(S)-HETE]. We speculate that this reaction occurs in a signaling complex near the inner membrane, where 12(S)-HETE then activates lamin-bound protein kinase C α (PKC α), leading to phosphorylation of unknown target proteins. (Bottom center) Pairwise interactions, summarized in the text, suggest that one or multiple oligomeric gene-regulation complexes might be formed by LAP2 α , A-type lamins, retinoblastoma (Rb), E2F/DP heterodimers (gene-specific activators), chromatin-silencing histone deacetylase complex (HDAC), BAF and additional unidentified chromatin partners for LAP2 α (Vlcek et al., 2002).

Furthermore, studies with LEM-domain proteins (emerin, LAP2 α and LAP2 β) and transcriptional regulators (GCL, BAF and Rb) also provide the best current evidence that lamins scaffold multiprotein complexes (Holaska et al., 2003; Markiewicz et al., 2002; Nili et al., 2001). Interestingly, the gene regulators known to bind A-type lamins are all repressors. Gene silencing is certainly key to cell differentiation, but it is premature to assume that all gene regulators act repressively when complexed with A-type lamins.

Signaling partners

Signal transduction pathways regulate many activities in the nucleus. Three signaling proteins are known to bind A-type lamins *in vivo*; lamin filaments might therefore provide a structural basis for signaling. Protein kinase C α (PKC α) – a serine-threonine kinase activated by many signals, including diacylglycerol (DAG) and 12(S)-hydroxyeicosatetraenoic acid [12(S)-HETE] – binds to the C-terminal half of the lamin A/C tail (Martelli et al., 2002) (Fig. 1). The lamin-binding and substrate-binding regions of PKC α are distinct. In response to

elevated DAG levels, PKC α translocates to the nucleus and interacts with A-type lamins, which suggests that signal transduction by PKC α might involve its attachment to a lamin scaffold (Martelli et al., 2000; Ron and Kazanietz, 1999) (Fig. 2).

12(S)-lipoxygenase [12(S)-LOX] also binds to the lamin A/C tail, perhaps overlapping the binding site for PKC α (Tang et al., 2000) (Fig. 1). Interestingly, 12(S)-LOX is part of a lipid-signaling pathway that converts arachidonic acid to 12(S)-HETE, which activates PKC α in prostate tumor cells (Liu et al., 1994). The 12(S)-LOX protein is detected in both the cytoplasm and nucleus in western blots of cell fractions (Tang et al., 2000). We speculate that PKC α might translocate to the nucleus and co-dock with 12(S)-LOX on lamin A/C filaments (Fig. 2). Alternatively, nuclear-localized PKC α might be activated in a 12(S)-LOX-dependent manner. The knowledge that PKC α and 12(S)-LOX are both direct partners for lamin A can be used to test the hypothesis that lamin scaffolds are important for PKC α -mediated signal transduction.

The third signaling partner, E1B 19K, is an adenovirus early protein (Cuconati and White, 2002). E1B 19K binds directly to lamin A in two-hybrid assays, and cofractionates with

lamins in vivo. In adenovirus-infected cells, E1B 19K, which shares a low level of sequence similarity with Bcl-2, localizes to the ER and nuclear membranes, and blocks apoptosis in a lamin-dependent manner (Rao et al., 1997). E1B 19K also interacts with (and potentially inactivates) a death-promoting repressor named Btf (Kasof et al., 1999), which binds to emerlin in vitro (Haraguchi et al., 2004). Does E1B 19K bind lamin A in order to co-assemble with Btf on emerlin-lamin complexes, or might it displace Btf? By considering each strand in this web of interactions, we may be able to understand how each partner 'uses' lamin A.

Perspectives

There are at least a million copies of each binding site on A-type lamins per nucleus. This repetition should ensure that all binding partners have access to lamin structures. To understand the functions of A-type lamins, we must now search for new binding partners, particularly those expressed in human tissues affected by laminopathy. However, it would be a mistake to neglect known partners, which provide tools to take this field to the next level. Old-fashioned (but still powerful) biochemical analysis, specific functional assays both in vitro and in vivo, and real-time analysis in living cells will all be needed if we are to understand how and why these complexes interact with lamins and what goes wrong in disease. Known partners for A-type lamins can also be mined for structural information to facilitate discovery of new partners. For example, partners whose binding sites on A-type lamins overlap (e.g. MOK2, Rb and E1B 19K in Fig. 1) might share a lamin-binding motif. Such motifs might allow investigators to identify potentially disease-relevant lamin-binding proteins rapidly in databases. We are hopeful that further work on lamin A/C partners, both known and novel, will lead to a better understanding of nuclear lamins and clinical treatments for diseases in which they are implicated.

References

- Agarwal, A. K., Fryns, J.-P., Auchus, R. J. and Garg, A. (2003). Zinc metalloproteinase, *ZMPSTE24*, is mutated in mandibuloacral dysplasia. *Hum. Mol. Genet.* **12**, 1995-2001.
- Arranz, V., Harper, F., Florentin, Y., Puvion, E., Kress, M. and Ernoult-Lange, M. (1997). Human and mouse MOK2 proteins are associated with nuclear ribonucleoprotein components and bind specifically to RNA and DNA through their zinc finger domains. *Mol. Cell. Biol.* **17**, 2116-2126.
- Barton, R. M. and Worman, H. J. (1999). Prenylated prelamin A interacts with Narf, a novel nuclear protein. *J. Biol. Chem.* **274**, 30008-30018.
- Bengtsson, L. and Wilson, K. L. (2004). Multiple and surprising new functions for emerlin, a nuclear membrane protein. *Curr. Opin. Cell Biol.* **16**, 73-79.
- Berger, R., Theodor, L., Shoham, J., Gokkel, E., Brok-Simoni, F., Avraham, K. B., Copeland, N. G., Jenkins, N. A., Rechavi, G. and Simon, A. J. (1996). The characterization and localization of the mouse thymopoietin/lamina-associated polypeptide 2 gene and its alternatively spliced products. *Genome Res.* **6**, 361-370.
- Bergo, M. O., Gavino, B., Ross, J., Schmidt, W. K., Hong, C., Kendall, L. V., Mohr, A., Meta, M., Genant, H., Jiang, Y. et al. (2002). Zmpste24 deficiency in mice causes spontaneous bone fractures, muscle weakness, and a prelamin A processing defect. *Proc. Natl. Acad. Sci. USA* **99**, 13049-13054.
- Bione, S., Maestrini, E., Rivella, S., Mancini, M., Regis, S., Romeo, G. and Toniolo, D. (1994). Identification of a novel X-linked gene responsible for Emery-Dreifuss muscular dystrophy. *Nat. Genet.* **8**, 323-327.
- Bonne, G., Yaou, R. B., Beroud, C., Boriani, G., Brown, S., de Visser, M., Duboc, D., Ellis, J., Hausmanowa-Petrusewicz, I., Lattanzi, G. et al. (2003). 108th ENMC International Workshop, 3rd Workshop of the MYO-CLUSTER project: EUROMEN, 7th International Emery-Dreifuss Muscular Dystrophy (EDMD) Workshop, 13-15 September 2002, Naarden, The Netherlands. *Neuromuscul. Disord.* **13**, 508-515.
- Burke, B. and Stewart, C. (2002). Life at the edge: the nuclear envelope and human disease. *Nat. Rev. Mol. Cell. Biol.* **3**, 575-585.
- Cai, M., Huang, Y., Ghirlardo, R., Wilson, K. L., Craigie, R. and Clore, G. M. (2001). Solution structure of the constant region of nuclear envelope protein LAP2 reveals two LEM-domain structures: one binds BAF and the other binds DNA. *EMBO J.* **20**, 4399-4407.
- Capanni, C., Cenni, V., Mattioli, E., Sabatelli, P., Ognibene, A., Columbaro, M., Parnaik, V. K., Wehnert, M., Maraldi, N. M., Squarzone, S. and Lattanzi, G. (2003). Failure of lamin A/C to functionally assemble in R482L mutated partial lipodystrophy fibroblasts: altered intermolecular interaction with emerlin and implications for gene transcription. *Exp. Cell Res.* **291**, 122-134.
- Chau, B. N. and Wang, J. Y. J. (2003). Coordinated regulation of life and death by RB. *Nat. Rev. Cancer* **3**, 130-138.
- Chen, L., Lee, L., Kudlow, B. A., Dos Santos, H. G., Sletvold, O., Shafeghati, Y., Botha, E. G., Garg, A., Hanson, N. B., Martin, G. M. et al. (2003). LMNA mutations in atypical Werner's syndrome. *Lancet* **362**, 440-445.
- Cohen, M., Lee, K. K., Wilson, K. L. and Gruenbaum, Y. (2001). Transcriptional repression, apoptosis, human disease and the functional evolution of the nuclear lamina. *Trends Biochem. Sci.* **26**, 41-47.
- Cuconati, A. and White, E. (2002). Viral homologs of BCL-2: role of apoptosis in the regulation of virus infection. *Genes Dev.* **16**, 2465-2478.
- De Sandre-Giovannoli, A., Bernard, R., Cau, P., Navarro, C., Amiel, J., Boccaccio, I., Lyonnet, S., Stewart, C. L., Munnich, A., Le Merrer, M. and Levy, N. (2003). Lamin A truncation in Hutchinson-Gilford progeria. *Science* **300**, 2055.
- Dechat, T., Gotzmann, J., Stockinger, A., Harris, C. A., Talle, M. A., Siekierka, J. J. and Foisner, R. (1998). Detergent-salt resistance of LAP2 α in interphase nuclei and phosphorylation-dependent association with chromosomes early in nuclear assembly implies functions in nuclear structure dynamics. *EMBO J.* **17**, 4887-4902.
- Dechat, T., Korbei, B., Vaughan, O. A., Vitek, S., Hutchison, C. J. and Foisner, R. (2000). Lamina-associated polypeptide 2 α binds intranuclear A-type lamins. *J. Cell Sci.* **113**, 3473-3484.
- Dhe-Paganon, S., Werner, E. D., Chi, Y. I. and Shoelson, S. E. (2002). Structure of the globular tail of nuclear lamin. *J. Biol. Chem.* **277**, 17381-17384.
- Dreuillet, C., Tillit, J., Kress, M. and Ernoult-Lange, M. (2002). In vivo and in vitro interaction between human transcription factor MOK2 and nuclear lamin A/C. *Nucleic Acids Res.* **30**, 4634-4642.
- Eriksson, M., Brown, W. T., Gordon, L. B., Glynn, M. W., Singer, J., Scott, L., Erdos, M. R., Robbins, C. M., Moses, T. Y., Berglund, P. et al. (2003). Recurrent de novo point mutations in lamin A cause Hutchinson-Gilford progeria syndrome. *Nature* **423**, 293-298.
- Fisher, D. Z., Chaudhary, N. and Blobel, G. (1986). cDNA sequencing of nuclear lamins A and C reveals primary and secondary structural homology to intermediate filament proteins. *Proc. Natl. Acad. Sci. USA* **83**, 6450-6454.
- Foisner, R. and Gerace, L. (1993). Integral membrane proteins of the nuclear envelope interact with lamins and chromosomes, and binding is modulated by mitotic phosphorylation. *Cell* **73**, 1267-1279.
- Foisner, R. (2003). Cell cycle dynamics of the nuclear envelope. *Scientific World Journal* **3**, 1-20.
- Furukawa, K., Inagaki, H. and Hotta, Y. (1994). Identification and cloning of an mRNA coding for a germ cell-specific A-type lamin in mice. *Exp. Cell Res.* **212**, 426-430.
- Furukawa, K. (1999). LAP2 binding protein 1 (L2BP1/BAF) is a candidate mediator of LAP2-chromatin interaction. *J. Cell Sci.* **112**, 2485-2492.
- Furukawa, K., Sugiyama, S., Osouda, S., Goto, H., Inagaki, M., Horigome, T., Omata, S., McConnell, M., Fisher, P. A. and Nishida, Y. (2003). Barrier-to-autointegration factor plays crucial roles in cell cycle progression and nuclear organization in *Drosophila*. *J. Cell Sci.* **116**, 3811-3823.
- Georgatos, S. D., Stournaras, C. and Blobel, G. (1988). Heterotypic and homotypic associations between the nuclear lamins: site-specificity and control by phosphorylation. *Proc. Natl. Acad. Sci. USA* **85**, 4325-4329.
- Goldman, R. D., Gruenbaum, Y., Moir, R. D., Shumaker, D. K. and Spann, T. P. (2002). Nuclear lamins: building blocks of nuclear architecture. *Genes Dev.* **16**, 533-547.
- Gough, L. L., Fan, J., Chu, S., Winnick, S. and Beck, K. A. (2003). Golgi localization of Syne-1. *Mol. Biol. Cell* **14**, 2410-2424.
- Haraguchi, T., Koujin, T., Segura-Toffen, M., Lee, K. K., Matsuoka, Y., Yoneda, Y., Wilson, K. L. and Hiraoka, Y. (2001). BAF is required for

- emerin assembly into the reforming nuclear envelope. *J. Cell Sci.* **114**, 4575-4585.
- Haraguchi, T., Holaska, J. M., Yamane, M., Koujin, T., Hashiguchi, N., Mori, C., Wilson, K. L. and Hiraoka, Y.** (2004). Emerin binding to Btf, a death-promoting transcriptional repressor, is disrupted by a missense mutation that causes Emery-Dreifuss muscular dystrophy. *Eur. J. Biochem.* (in press).
- Harborth, J., Elbashir, S. M., Bechert, K., Tuschl, T. and Weber, K.** (2001). Identification of essential genes in cultured mammalian cells using small interfering RNAs. *J. Cell Sci.* **114**, 4557-4565.
- Holaska, J. M., Lee, K. K., Kowalski, A. K. and Wilson, K. L.** (2003). Transcriptional repressor germ cell-less (GCL) and barrier to autointegration factor (BAF) compete for binding to emerin in vitro. *J. Biol. Chem.* **278**, 6969-6975.
- Horton, J. D.** (2002). Sterol regulatory element-binding proteins: transcriptional activators of lipid synthesis. *Biochem. Soc. Trans.* **30**, 1091-1095.
- Izumi, M., Vaughan, O. A., Hutchison, C. J. and Gilbert, D. M.** (2000). Head and/or CaaX domain deletions of lamin proteins disrupt preformed lamin A and C but not lamin B structure in mammalian cells. *Mol. Biol. Cell* **11**, 4323-4337.
- Kasof, G. M., Goyal, L. and White, E.** (1999). Btf, a novel death-promoting transcriptional repressor that interacts with Bcl-2-related proteins. *Mol. Cell. Biol.* **19**, 4390-4404.
- Kim, J. B. and Spiegelman, B. M.** (1996). ADD1/SREBP1 promotes adipocyte differentiation and gene expression linked to fatty acid metabolism. *Genes. Dev.* **10**, 1096-1107.
- Krimm, I., Ostlund, C., Gilquin, B., Couprie, J., Hossenlopp, P., Mornon, J. P., Bonne, G., Courvalin, J. C., Worman, H. J. and Zinn-Justin, S.** (2002). The Ig-like structure of the C-terminal domain of lamin A/C, mutated in muscular dystrophies, cardiomyopathy, and partial lipodystrophy. *Structure (Camb.)* **10**, 811-823.
- Lee, K. K., Gruenbaum, Y., Spann, P., Liu, J. and Wilson, K. L.** (2000). *C. elegans* nuclear envelope proteins emerin, MAN1, lamin, and nucleoporins reveal unique timing of nuclear envelope breakdown during mitosis. *Mol. Biol. Cell* **11**, 3089-3099.
- Lee, K. K., Haraguchi, T., Lee, R. S., Koujin, T., Hiraoka, Y. and Wilson, K. L.** (2001). Distinct functional domains in emerin bind lamin A and DNA-bridging protein BAF. *J. Cell Sci.* **114**, 4567-4573.
- Lee, K. K. and Wilson, K. L.** (2004). All in the family: evidence for three new LEM-domain proteins Lem3, Lem2 (NET-25) and Lem1 in the human genome. In *Communication and Gene Regulation at the Nuclear Envelope* (ed. D. Evans, J. Bryant and C. J. Hutchison). Oxford: BIOS Scientific Publishers (in press).
- Lin, F., Blake, D. L., Callebaut, I., Skerjanc, I. S., Holmer, L., McBurney, M. W., Paulin-Levasseur, M. and Worman, H. J.** (2000). MAN1, an inner nuclear membrane protein that shares the LEM domain with lamina-associated polypeptide 2 and emerin. *J. Biol. Chem.* **275**, 4840-4847.
- Liu, B., Maher, R. J., Hannun, Y. A., Porter, A. T. and Honn, K. V.** (1994). 12(S)-HETE enhancement of prostate tumor cell invasion: selective role of PKC alpha. *J. Natl. Cancer Inst.* **86**, 1145-1151.
- Liu, J., Lee, K. K., Segura-Totten, M., Neufeld, E., Wilson, K. L. and Gruenbaum, Y.** (2003). MAN1 and emerin have overlapping function(s) essential for chromosome segregation and cell division in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* **100**, 4598-4603.
- Lloyd, D. J., Trembath, R. C. and Shackleton, S.** (2002). A novel interaction between lamin A and SREBP1: implications for partial lipodystrophy and other laminopathies. *Hum. Mol. Genet.* **11**, 769-777.
- Luderus, M. E., den Blaauwen, J. L., de Smit, O. J., Compton, D. A. and van Driel, R.** (1994). Binding of matrix attachment regions to lamin polymers involves single-stranded regions and the minor groove. *Mol. Cell. Biol.* **14**, 6297-6305.
- Machiels, B. M., Zorenc, A. H., Endert, J. M., Kuijpers, H. J., van Eys, G. J., Ramaekers, F. C. and Broers, J. L.** (1996). An alternative splicing product of the lamin A/C gene lacks exon 10. *J. Biol. Chem.* **271**, 9249-9253.
- Mancini, M. A., Shan, B., Nickerson, J. A., Penman, S. and Lee, W. H.** (1994). The retinoblastoma gene product is a cell cycle-dependent, nuclear matrix-associated protein. *Proc. Natl. Acad. Sci. USA* **91**, 418-422.
- Markiewicz, E., Dechat, T., Foisner, R., Quinlan, R. A. and Hutchison, C. J.** (2002). Lamin A/C binding protein LAP2 α is required for nuclear anchorage of retinoblastoma protein. *Mol. Biol. Cell* **13**, 4401-4413.
- Martelli, A. M., Tabellini, G., Bortul, R., Manzoli, L., Bareggi, R., Baldini, G., Grill, V., Zweyer, M., Narducci, P. and Cocco, L.** (2000). Enhanced nuclear diacylglycerol kinase activity in response to a mitogenic stimulation of quiescent Swiss 3T3 cells with insulin-like growth factor I. *Cancer Res.* **60**, 815-821.
- Martelli, A. M., Bortul, R., Tabellini, G., Faenza, I., Cappellini, A., Bareggi, R., Manzoli, L. and Cocco, L.** (2002). Molecular characterization of protein kinase C-alpha binding to lamin A. *J. Cell Biochem.* **86**, 320-330.
- Martin, L., Crimando, C. and Gerace, L.** (1995). cDNA cloning and characterization of lamina-associated polypeptide 1C (LAP1C), an integral protein of the inner nuclear membrane. *J. Biol. Chem.* **270**, 8822-8828.
- McKeon, F. D., Kirschner, M. W. and Caput, D.** (1986). Homologies in both primary and secondary structure between nuclear envelope and intermediate filament proteins. *Nature* **319**, 463-468.
- Milankov, K. and De Boni, U.** (1993). Cytochemical localization of actin and myosin aggregates in interphase nuclei in situ. *Exp. Cell Res.* **209**, 189-199.
- Mislow, J. M. K., Holaska, J. M., Kim, M. S., Lee, K. K., Segura-Totten, M., Wilson, K. L. and McNally, E. M.** (2002a). Nesprin-1 α self-associates and binds directly to emerin and lamin A in vitro. *FEBS Lett.* **525**, 135-140.
- Mislow, J. M. K., Kim, M. S., Davis, D. B. and McNally, E. M.** (2002b). Myne-1, a spectrin repeat transmembrane protein of the myocyte inner nuclear membrane, interacts with lamin A/C. *J. Cell Sci.* **115**, 61-70.
- Moir, R. D., Spann, T. P. and Goldman, R. D.** (1995). The dynamic properties and possible functions of nuclear lamins. *Int. Rev. Cytol.* **162B**, 141-182.
- Moir, R. D., Yoon, M., Khuon, S. and Goldman, R. D.** (2000). Nuclear lamins A and B1: different pathways of assembly during nuclear envelope formation in living cells. *J. Cell Biol.* **151**, 1155-1168.
- Mounkes, L. C., Kozlov, S., Hernandez, L., Sullivan, T. and Stewart, C. L.** (2003). A progeroid syndrome in mice is caused by defects in A-type lamins. *Nature* **423**, 298-301.
- Muchir, A., van Engelen, B. G., Lammens, M., Mislow, J. M., McNally, E., Schwartz, K. and Bonne, G.** (2003). Nuclear envelope alterations in fibroblasts from LGMD1B patients carrying nonsense Y259X heterozygous or homozygous mutation in lamin A/C gene. *Exp. Cell Res.* **291**, 352-362.
- Neely, K. E. and Workman, J. L.** (2002). Histone acetylation and chromatin remodeling: which comes first? *Mol. Genet. Metab.* **76**, 1-5.
- Nili, E., Cojocar, G. S., Kalma, Y., Ginsberg, D., Copeland, N. G., Gilbert, D. J., Jenkins, N. A., Berger, R., Shakkai, S., Amariglio, N. et al.** (2001). Nuclear membrane protein LAP2 β mediates transcriptional repression alone and together with its binding partner GCL (germ-cell-less). *J. Cell Sci.* **114**, 3297-3307.
- Osada, S., Ohmori, S. Y. and Taira, M.** (2003). XMAN1, an inner nuclear membrane protein, antagonizes BMP signaling by interacting with Smad1 in *Xenopus* embryos. *Development* **130**, 1783-1794.
- Östlund, C., Ellenberg, J., Hallberg, E., Lippincott-Schwartz, J. and Worman, H. J.** (1999). Intracellular trafficking of emerin, the Emery-Dreifuss muscular dystrophy protein. *J. Cell Sci.* **112**, 1709-1719.
- Östlund, C. and Worman, H. J.** (2003). Nuclear envelope proteins and neuromuscular diseases. *Muscle Nerve* **27**, 393-406.
- Ozaki, T., Saijo, M., Murakami, K., Enomoto, H., Taya, Y. and Sakiyama, S.** (1994). Complex formation between lamin A and the retinoblastoma gene product: identification of the domain on lamin A required for its interaction. *Oncogene* **9**, 2649-2653.
- Pederson, T. and Aebi, U.** (2002). Actin in the nucleus: what form and what for? *J. Struct. Biol.* **140**, 3-9.
- Pendas, A. M., Zhou, Z., Cadinanos, J., Freije, J. M., Wang, J., Hultenby, K., Astudillo, A., Wernerson, A., Rodriguez, F., Tryggvason, K. and Lopez-Otin, C.** (2002). Defective prelamin A processing and muscular and adipocyte alterations in Zmpste24 metalloproteinase-deficient mice. *Nat. Genet.* **31**, 94-99.
- Raharjo, W. H., Enarson, P., Sullivan, T., Stewart, C. L. and Burke, B.** (2001). Nuclear envelope defects associated with LMNA mutations causing dilated cardiomyopathy and Emery-Dreifuss muscular dystrophy. *J. Cell Sci.* **114**, 4447-4457.
- Raju, G. P., Dimova, N., Klein, P. S. and Huang, H. C.** (2003). SANE, a novel LEM domain protein, regulates bone morphogenetic protein signaling through interaction with Smad1. *J. Biol. Chem.* **278**, 428-437.
- Rao, L., Modha, D. and White, E.** (1997). The E1B 19K protein associates with lamins in vivo and its proper localization is required for inhibition of apoptosis. *Oncogene* **15**, 1587-1597.
- Robber, R. A., Weber, K. and Osborn, M.** (1989). Differential timing of nuclear lamin A/C expression in the various organs of the mouse embryo and the young animal: a developmental study. *Development* **105**, 365-378.

- Ron, D. and Kazanietz, M. G. (1999). New insights into the regulation of protein kinase C and novel phorbol ester receptors. *FASEB J.* **13**, 1658-1676.
- Sakaki, M., Koike, H., Takahashi, N., Sasagawa, N., Tomioka, S., Arahata, K. and Ishiura, S. (2001). Interaction between emerin and nuclear lamins. *J. Biochem.* **129**, 321-327.
- Sasseville, A. M. and Langelier, Y. (1998). In vitro interaction of the carboxy-terminal domain of lamin A with actin. *FEBS Lett.* **425**, 485-489.
- Schirmer, E. C., Florens, L., Guan, T., Yates, J. R., III and Gerace, L. (2003). Nuclear membrane proteins with potential disease links found by subtractive proteomics. *Science* **301**, 1380-1382.
- Segura-Totten, M. and Wilson, K. L. (2004). Barrier to autointegration factor (BAF): roles in chromatin structure, gene regulation, nuclear assembly and HIV-1 infection. *Trends Cell Biol.* (in press).
- Segura-Totten, M., Kowalski, A. K., Craigie, R. and Wilson, K. L. (2002). Barrier-to-autointegration factor: major roles in chromatin decondensation and nuclear assembly. *J. Cell Biol.* **158**, 475-485.
- Shimi, T., Koujin, T., Segura-Totten, M., Wilson, K. L., Hiraoka, H. and Haraguchi, T. (2004). Dynamic interaction between BAF and emerin revealed by FRAP, FLIP and FRET analysis in living HeLa cells. *J. Struct. Biol.* (in press).
- Shumaker, D. K., Lee, K. K., Tanhehco, Y. C., Craigie, R. and Wilson, K. L. (2001). LAP2 binds to BAF-DNA complexes: requirement for the LEM domain and modulation by variable regions. *EMBO J.* **20**, 1754-1764.
- Starr, D. A. and Han, M. (2003). ANChors away: an actin based mechanism of nuclear positioning. *J. Cell Sci.* **116**, 211-216.
- Stierle, V., Couprie, J., Ostlund, C., Krimm, I., Zinn-Justin, S., Hossenlopp, P., Worman, H. J., Courvalin, J. C. and Duband-Goulet, I. (2003). The carboxyl-terminal region common to lamins A and C contains a DNA binding domain. *Biochemistry* **42**, 4819-4828.
- Stuurman, N., Heins, S. and Aebi, U. (1998). Nuclear lamins: their structure, assembly, and interactions. *J. Struct. Biol.* **122**, 42-66.
- Sullivan, T., Escalante-Alcalde, D., Bhatt, H., Anver, M., Bhat, N., Nagashima, K., Stewart, C. L. and Burke, B. (1999). Loss of A-type lamin expression compromises nuclear envelope integrity leading to muscular dystrophy. *J. Cell Biol.* **147**, 913-920.
- Tang, K., Finley, R. L., Nie, D. and Honn, K. V. (2000). Identification of 12-lipoxygenase interaction with cellular proteins by yeast two-hybrid screening. *Biochemistry* **39**, 3185-3191.
- Taniura, H., Glass, C. and Gerace, L. (1995). A chromatin binding site in the tail domain of nuclear lamins that interacts with core histones. *J. Cell Biol.* **131**, 33-44.
- Vaughan, A., Alvarez-Reyes, M., Bridger, J. M., Broers, J. L., Ramaekers, F. C., Wehnert, M., Morris, G. E., Whitfield, W. G. F. and Hutchison, C. J. (2001). Both emerin and lamin C depend on lamin A for localization at the nuclear envelope. *J. Cell Sci.* **114**, 2577-2590.
- Vlcek, S., Just, H., Dechat, T. and Foisner, R. (1999). Functional diversity of LAP2 α and LAP2 β in postmitotic chromosome association is caused by an α -specific nuclear targeting. *EMBO J.* **18**, 6370-6384.
- Vlcek, S., Korbei, B. and Foisner, R. (2002). Distinct functions of the unique C terminus of LAP2 α in cell proliferation and nuclear assembly. *J. Biol. Chem.* **277**, 18898-18907.
- Wang, X., Xu, S., Rivolta, C., Li, L. Y., Peng, G. H., Swain, P. K., Sung, C. H., Swaroop, A., Berson, E. L., Dryja, T. P. and Chen, S. (2002). Barrier to autointegration factor interacts with the cone-rod homeobox and represses its transactivation function. *J. Biol. Chem.* **277**, 43288-43300.
- Ye, Q. and Worman, H. J. (1995). Protein-protein interactions between human nuclear lamins expressed in yeast. *Exp. Cell Res.* **219**, 292-298.
- Zhang, Q., Skepper, J. N., Yang, F., Davies, J. D., Hegyi, L., Roberts, R. G., Weissberg, P. L., Ellis, J. A. and Shanahan, C. M. (2001). Nesprins: a novel family of spectrin-repeat-containing proteins that localize to the nuclear membrane in multiple tissues. *J. Cell Sci.* **114**, 4485-4498.
- Zhang, Q., Ragnauth, C., Greener, M. J., Shanahan, C. M. and Roberts, R. G. (2002). The nesprins are giant actin-binding proteins, orthologous to *Drosophila melanogaster* muscle protein MSP-300. *Genomics* **80**, 473-481.
- Zheng, R., Ghirlando, R., Lee, M. S., Mizuuchi, K., Krause, M. and Craigie, R. (2000). Barrier-to-autointegration factor (BAF) bridges DNA in a discrete, higher-order nucleoprotein complex. *Proc. Natl. Acad. Sci. USA* **97**, 8997-9002.
- Zimek, A., Stick, R. and Weber, K. (2003). Genes coding for intermediate filament proteins: common features and unexpected differences in the genomes of humans and the teleost fish *Fugu rubripes*. *J. Cell Sci.* **116**, 2295-2302.