

# The insulin-PI3K/TOR pathway induces a HIF-dependent transcriptional response in *Drosophila* by promoting nuclear localization of HIF- $\alpha$ /Sima

Andrés Dekanty<sup>1</sup>, Sofía Lavista-Llanos<sup>1</sup>, Maximiliano Irisarri<sup>1</sup>, Sean Oldham<sup>2</sup> and Pablo Wappner<sup>1,\*</sup>

<sup>1</sup>Instituto Leloir and IIB, FCEyN-Universidad de Buenos Aires, CONICET, Patricias Argentinas 435, Buenos Aires (1405), Argentina

<sup>2</sup>The Burnham Institute, La Jolla, CA 92037, USA

\*Author for correspondence (e-mail: pwappner@leloir.org.ar)

Accepted 16 August 2005

Journal of Cell Science 118, 5431-5441 Published by The Company of Biologists 2005

doi:10.1242/jcs.02648

## Summary

The hypoxia-inducible factor (HIF) is a heterodimeric transcription factor composed of a constitutively expressed HIF- $\beta$  subunit and an oxygen-regulated HIF- $\alpha$  subunit. We have previously defined a hypoxia-inducible transcriptional response in *Drosophila melanogaster* that is homologous to the mammalian HIF-dependent response. In *Drosophila*, the bHLH-PAS proteins Similar (Sima) and Tango (Tgo) are the functional homologues of the mammalian HIF- $\alpha$  and HIF- $\beta$  subunits, respectively. HIF- $\alpha$ /Sima is regulated by oxygen at several different levels that include protein stability and subcellular localization. We show here for the first time that insulin can activate HIF-dependent transcription, both in *Drosophila* S2 cells and in living *Drosophila* embryos. Using a pharmacological approach as well as RNA interference, we determined that the effect of insulin on HIF-dependent transcriptional induction is mediated by PI3K-AKT and TOR pathways. We demonstrate that stimulation of the transcriptional response involves upregulation of Sima protein but not *sima*

mRNA. Finally, we have analyzed in vivo the effect of the activation of the PI3K-AKT pathway on the subcellular localization of Sima protein. Overexpression of dAKT and dPDK1 in normoxic embryos provoked a major increase in Sima nuclear localization, mimicking the effect of a hypoxic treatment. A similar increase in Sima nuclear localization was observed in *dPTEN* homozygous mutant embryos, confirming that activation of the PI3K-AKT pathway promotes nuclear accumulation of Sima protein. We conclude that regulation of HIF- $\alpha$ /Sima by the PI3K-AKT-TOR pathway is a major conserved mode of regulation of the HIF-dependent transcriptional response in *Drosophila*.

Supplementary material available online at  
<http://jcs.biologists.org/cgi/content/full/118/23/5431/DC1>

Key words: Sima, Hypoxia-inducible factor (HIF), Nuclear localization, *Drosophila*, PI3K pathway.

## Introduction

The mammalian hypoxia-inducible factor-1 (HIF-1) is an  $\alpha/\beta$  heterodimeric transcription factor composed of two basic helix-loop-helix-PAS (bHLH-PAS) subunits (Wang et al., 1995). HIF-1 binds to HIF responsive elements (HREs) in the regulatory region of a wide array of hypoxia-regulated target genes, thereby inducing hypoxia-dependent transcription (Maxwell et al., 1993; Wang and Semenza, 1993). Over the last decade, substantial progress in understanding the mechanisms that underlie HIF-1 oxygen-dependent regulation has been reported (Bruick, 2003; Schofield and Ratcliffe, 2004). It is now clear that, whereas the  $\beta$  subunit is constitutive, the  $\alpha$  subunit is regulated by oxygen tension at several different levels that include control of protein half-life, transcriptional co-activator recruitment and regulation of subcellular localization. Regulation of HIF- $\alpha$  degradation depends on oxygen-dependent hydroxylation of two key prolyl residues (P402 and P564), that is catalyzed by specific prolyl-4-hydroxylases (Bruick and McKnight, 2001; Epstein et al., 2001; Ivan, 2001; Jaakkola, 2001). This post-translational modification enables interaction with the ubiquitination

machinery, thus promoting proteasomal degradation (Maxwell et al., 1999).

Transcriptional coactivator recruitment is regulated by a second oxygen-sensing system in which an asparaginyl hydroxylase, a different oxoglutarate and iron-dependent dioxygenase, promotes hydroxylation of a specific asparaginyl residue located within the HIF- $\alpha$  C-terminal domain. Hydroxylation of this asparagine prevents interaction with the transcriptional co-activator p300, resulting in the inhibition of transactivation (Dann et al., 2002; Elkins et al., 2002; Hewitson et al., 2002; Lando et al., 2002). A third regulatory mechanism, the control of subcellular localization determines that, in normoxia, HIF- $\alpha$  is mostly cytoplasmic, and in hypoxia it accumulates in the nuclear compartment (Kallio et al., 1998). Structure-function studies have led to the identification of a functional nuclear localization signal and of additional domains that are required for regulated nuclear localization (Kallio et al., 1998; Luo and Shibuya, 2001). Nevertheless, the cellular and molecular mechanisms controlling HIF- $\alpha$  oxygen-dependent subcellular localization are poorly understood.

In addition, it has been shown that several non-hypoxic

stimuli, such as nitric oxide, growth factors, hormones and cytokines, can modulate HIF-1-dependent transcription (Feldser et al., 1999; Fukuda et al., 2003; Kasuno et al., 2004). In particular, stimulation with insulin or with the insulin-like growth factor 1/2 (IGF1/2) can upregulate HIF- $\alpha$  protein levels, thereby inducing HIF-1 target genes (Feldser et al., 1999; Kietzmann et al., 2003; Zelzer et al., 1998). Since insulin can activate two parallel signaling pathways – the mitogen-activated protein kinase (MAPK) and the phosphoinositide 3-kinase (PI3K) pathway (Oldham and Hafen, 2003; Yenush et al., 1996; Miele et al., 2000) – several groups have investigated which of these two pathways is involved in HIF-1-dependent gene induction. Although this issue is still controversial, the PI3K pathway seems to be prevalent (Kietzmann et al., 2003; Roth et al., 2004; Treins et al., 2002). Activation of the PI3K-AKT pathway by insulin starts when the ligand binds to the insulin receptor (InR), which in turn interacts with InR substrate (IRS) adapter proteins, thereby activating PI3K. This kinase catalyzes the phosphorylation of phosphatidylinositol (4,5)-bisphosphate [PtdIns(4,5) $P_2$ ] to phosphatidylinositol (3,4,5)-trisphosphate [PtdIns(3,4,5) $P_3$ ] in a reaction antagonized by the phosphatase PTEN. PtdIns(3,4,5) $P_3$  promotes recruitment to the plasma membrane of two other kinases, AKT/PKB and PDK1, resulting in the full activation of AKT/PKB, which in turn phosphorylates downstream substrates (Yenush and White, 1997). It has been shown that PI3K-AKT-dependent increase of HIF- $\alpha$  protein levels is neither due to protein stabilization, nor involves the participation of prolyl hydroxylases or the von Hippel-Lindau (VHL) protein (Fukuda et al., 2002). Rather, augmented HRE activity is due to increased HIF- $\alpha$  translation, which is mediated by the target of rapamycin kinase (TOR) (Laughner et al., 2001; Treins et al., 2002; Zhong et al., 2000).

In *Drosophila melanogaster*, the InR-PI3K-AKT pathway is conserved (Fernandez et al., 1995; Garofalo and Rosen, 1988; LeRoith et al., 1981; Seecof and Dewhurst, 1974) and exerts several different functions including control of growth, anabolic metabolism and lifespan (Oldham and Hafen, 2003). We and others have recently defined a hypoxia-responsive system that is homologous to mammalian HIF (Bruick and McKnight, 2001; Gorr et al., 2004; Lavista-Llanos et al., 2002), in which the bHLH-PAS proteins Similar (Sima) and Tango (Tgo), are the *Drosophila* functional homologues of HIF- $\alpha$  and HIF- $\beta$ , respectively. In a hypoxic situation Sima and Tango were shown to mediate transcriptional induction of an HRE reporter both in cell culture and in vivo, as well as regulation of specific oxygen-regulated genes. The molecular mechanisms controlling the hypoxia-responsive machinery are largely conserved because oxygen tension determines the half-life of Sima protein, which is rapidly degraded in normoxia and stabilized in hypoxia. Normoxic Sima degradation depends on the activity of a conserved prolyl-4-hydroxylase that we have named Fatiga. In *fatiga* loss-of-function mutants Sima is stabilized, and high expression of the HRE reporter occurs constitutively in normoxia (Lavista-Llanos et al., 2002). Conservation of the cellular machinery that regulates the hypoxic response extends even further, because Sima subcellular localization also depends on oxygen tension. When rapid degradation was overwhelmed in vivo through the overexpression of Sima in transgenic fly lines, Sima accumulated in the cytoplasm in normoxia and became nuclear

in hypoxia (Lavista-Llanos et al., 2002). The mechanism controlling HIF- $\alpha$ /Sima oxygen-dependent subcellular localization is so far unclear.

Involvement of the insulin pathway in the induction of the HRE response in *Drosophila* has not yet been reported. Here, we show both in cell culture and in vivo that the PI3K-AKT signaling pathway can activate an HRE response in *Drosophila* that is mediated by the bHLH-PAS proteins Sima and Tango. HRE induction depends on TOR and its upstream regulator Rheb and involves upregulation of Sima protein levels as well as an increase of its nuclear localization.

## Materials and Methods

### Cell culture

Schneider's line (S2) cells were maintained at 25°C in Schneider *Drosophila* medium (Sigma) supplemented with 10% fetal bovine serum (Gibco), containing 50 units/ml penicillin and 50  $\mu$ g/ml streptomycin in 25 or 75 cm<sup>2</sup> T-flasks (Greiner). Cells were plated in 12-, 24- or 96-well cell culture dishes (Greiner), cultured for 3 days and then treated with 1-50  $\mu$ g/ml insulin (Gibco), deferroxamine (Sigma) or were exposed to hypoxia in a Forma Scientific 3131 incubator.

### Plasmids, transfections and luciferase assays

To generate the HRE-luciferase reporter plasmid (HRE-Luc), a dimerized 51-bp sequence derived from the murine lactate dehydrogenase-A enhancer (Lavista-Llanos et al., 2002) was inserted into *KpnI* and *XhoI* sites of the polylinker of a pGL3 vector (Promega), containing a *Drosophila* Hsp70 basal promoter (Darlington et al., 1998). *Drosophila* Sima and human AKT coding sequences were inserted into *EcoRI-XbaI* and *EcoRI-HindIII* sites, respectively, of a pAC/V5His plasmid (Invitrogen); human AKT cDNA was a gift from Marcelo Katzanietz (Department of Pharmacology, University of Philadelphia, PA).

For transfections, 10<sup>6</sup> cells were plated in a 12-well plate and cultured overnight. Transfections were performed with a calcium phosphate transfection kit (Invitrogen) with 5  $\mu$ g of total DNA using pcDNA3 plasmid as filler DNA when necessary. To generate a stable cell line bearing the HRE-Luc reporter, cells were co-transfected with the pGL3 plasmid and with a second plasmid encoding a blasticidine resistance gene (pBLAST) in a 19:1 ratio. Forty-eight hours after transfection, 20  $\mu$ g/ml of blasticidine (Invitrogen) were added to the cells and selection was carried out for 3 weeks. For transcription assays in the stable cell line treated with various stimuli, luciferase activity was measured, normalized to protein concentrations and expressed as fold-induction with respect to non-stimulated cells. For transient transfection experiments, expression vectors were co-transfected together with a *Renilla* luciferase plasmid (pAC/*Renilla* luciferase) to normalize transfection efficiency. Forty-eight hours after transfections cells were harvested, lysed and *firefly* and *Renilla* luciferase activities were determined. Normalized luciferase activity (i.e. *Renilla*:luciferase activity ratio) was expressed as fold-induction with respect to the transfection control (empty vector). Luciferase activity was measured using Steady-Glo or Dual-Glo luciferase reagents (Promega) by following instructions of the manufacturer. Protein quantification was performed by the Bradford method (BIORAD).

### RNA isolation, northern blots and RT-PCR

Total RNA from cells exposed to different treatments was isolated using the Trizol reagent (Invitrogen). Thirty micrograms of total RNA were separated on denaturing 6% formaldehyde 1% agarose gels and transferred in 10 $\times$  SCC buffer to Zeta-Probe GT Genomic

membranes (BIO-RAD) and hybridized with [<sup>32</sup>P]CTP-labeled *sima*, LDH or 18S rRNA probes; a storm 840 phosphorimager (Molecular Dynamics) was used to detect <sup>32</sup>P emission. Probes were synthesized using 25 ng of the corresponding PCR products and labeled with [<sup>32</sup>P]CTP using a Random Primer Labeling Kit (Invitrogen). PCR products used to prepare probes were: *Sima* (nucleotides 612-1312), 18S rRNA (complete sequence), LDH (nucleotides 157-1118).

For RT-PCR experiments reverse transcription was carried out with a superscript reverse transcriptase enzyme (Invitrogen), according to the instructions of the manufacturer, using oligo-dT as a primer. Semi-quantitative RT-PCR experiments were performed at different cycles of amplification using a hot-start DNA polymerase enzyme (*Taq*-platinum, Invitrogen) and the analysis was carried out in each case at the exponential phase of amplification. For normalization, *actin42A* was amplified along with the particular transcript of interest.

### RNA interference

Fragments from the *sima* transcript (nucleotides 4099-5058), *tango* (nucleotides 901-1413), *dPTEN* (nucleotides 205-912), *dAKT* (nucleotides 784-1510), *dPDK1* (nucleotides 1749-2261), *dTOR* (nucleotides 3694-4208), *dRheb* (nucleotides 455-645) and *dS6K* (nucleotides 376-1165) were amplified by reverse transcriptase (RT)-PCR using specific primers containing part of the T7 RNA-polymerase-binding site at the 5' end of each primer, and a second PCR using the complete sequence of the T7 RNA-polymerase-binding site (5'-GAATTAATACGACTCACTATAGGGAGA-3') as primer. PCR products were purified in Microspin G25 columns (Amersham) and 1 µg of these DNA products was used as a template to synthesize double-stranded (ds)RNA with the T7 Megascript kit (Ambion) following manufacturer's instructions. The synthesized RNA was precipitated with LiCl, resuspended in water and annealed by heating at 65°C for 30 minutes followed by slow cooling until reaching room temperature. RNA interference (RNAi) experiments in S2 cells were performed as previously described (Clemens et al., 2000). Briefly, S2 cells were plated at a density of 10<sup>6</sup> cells/ml in 96- or 24-well plates in serum-free medium, and 1 to 5 µg of dsRNA were added. After 1 hour incubation, serum-containing medium was added and cells were cultured for 3 days at 25°C. Then, cells were stimulated with insulin, or exposed to hypoxia or the iron chelator deferoxamine (DFO) for luciferase reporter assays. To confirm silencing after the RNAi treatment, cells were harvested and endogenous mRNA levels were analysed by semi-quantitative RT-PCR.

### Immunofluorescence and β-galactosidase assays

Immunofluorescence was performed as previously described (Lavista-Llanos et al., 2002). Briefly, embryos from different developmental stages were bleach-dechorinated and fixed with 3.7% formaldehyde for 20 minutes; fixed embryos were blocked for 2 hours in PBS containing 1% BSA and 0.1% Triton X-100 (PT) and then incubated overnight with the primary antibody. After washing, embryos were incubated with the secondary antibody for 2 hours in PT containing 10% normal goat serum. Analysis of subcellular localization was performed in an Olympus BX60 microscope or in a Carl Zeiss LSM5 Pascal confocal microscope. For X-Gal stainings, embryos were bleach-dechorinated and fixed with 0.5% glutaraldehyde for 20 minutes; fixed embryos were washed three times in PT and incubated with the β-galactosidase synthetic substrate X-gal.

To quantify the subcellular localization of *Sima*, at least two independent experiments were carried out and 30 or more embryos from each analyzed developmental stage were recorded. The subcellular localization was scored in three categories (as described in the Results section) and data are represented as histograms showing the percentage of embryos in each category. The  $\chi^2$ -test was applied to compare different samples.

### SDS-PAGE and immunoblotting

Twenty-five micrograms of nuclear protein extracts were prepared as described in Gorr et al. (Gorr et al., 2004), separated in 6% SDS-polyacrylamide gel electrophoresis (PAGE) and blotted onto PVDF membranes (BIO-RAD). Thereafter, membranes were blocked for 1 hour at room temperature with 5% nonfat milk in TBS with 0.1% Tween 20 (TBS-T) and then incubated overnight with rat anti-*Sima* antiserum (1:1000) or mouse anti-Hsp70 antibody (1:1000) in 5% nonfat milk in TBS-T (Bacon et al., 1998). Membranes were washed in TBS-T and incubated for 1 hour at room temperature with a peroxidase-conjugated anti-rat or anti-mouse secondary antibody (Jackson) diluted 1:5000 in 5% nonfat milk in TBS-T. Immunoblots were developed with the ECL detection reagent (Cell Signalling).

### Fly stocks and genetics

The following fly strains were used: *PTEN*<sup>2L117</sup> (Oldham et al., 2000), *UAS-dAKT*; *PDK1*<sup>EP0857</sup> (Rintelen et al., 2001), *UAS-dPTEN* (Huang et al., 1999) *UAS-Sima* and *LDH-LacZ* (Lavista-Llanos et al., 2002); *engrailed-Gal4* (*en-GAL4*, Bloomington Stock Center).

Embryos collected in overnight egg-laying agar plates were kept in normoxia or exposed to hypoxia for 4 hours; then embryos were recovered, fixed and immunostained. Hypoxia was applied by regulating the proportions of oxygen and nitrogen in a Forma Scientific 3131 incubator at 25°C.

In *Sima* subcellular localization experiments, ectopic expression of *Sima* was performed by crossing *en-GAL4* and *UAS-Sima* fly strains. To study the effect of the mutation *dPTEN*<sup>2L117</sup> on *Sima* subcellular localization, *dPTEN*<sup>2L117</sup> *UAS-Sima* and *dPTEN*<sup>2L117</sup> *en-GAL4* recombinant chromosomes were generated by meiotic recombination. Briefly, heterozygous *yw*; *dPTEN*<sup>2L117</sup>/*Cyo* flies were crossed with *yw*; *UAS-Sima*/*Cyo* or *yw*; *en-GAL4* flies and non-*Cyo* females from the F1 were crossed with *yw*; *Xa*/*Cyo* males. Males bearing *UAS-Sima* or *en-GAL4* elements, identified by red eyes, were tested for lethality by crossing them with *yw*; *dPTEN*<sup>2L117</sup>/*Cyo* females. Once established, the recombinant strains *yw*; *dPTEN*<sup>2L117</sup> *UAS-Sima*/*Cyo* males were crossed with *yw*; *dPTEN*<sup>2L117</sup> *en-GAL4*/*Cyo* females and embryos from the offspring were immunostained. To restore *PTEN* expression in the *PTEN* mutant background, a strain with the following genotype was generated *yw*; *dPTEN*<sup>2L117</sup> *UAS-Sima*/*Cyo*; *UAS-dPTEN* and males from this strain were crossed to *yw*; *dPTEN*<sup>2L117</sup> *en-Gal4*/*Cyo* females and analyzed as above. To study whether overexpression of *dAKT* and *dPDK1* affect *Sima* subcellular localization, recombinant chromosomes bearing *UAS-Sima* and *UAS-dAKT* were generated by meiotic recombination as above. Males bearing *UAS-Sima* and *UAS-dAKT* elements were identified by strong red eyes and used to establish individual strains; the presence of both *UAS* constructs was confirmed by RNA in-situ hybridization. Subcellular localization of *Sima* was studied in crosses with *en-GAL4* flies and processed for anti-*Sima* immunostaining as above.

## Results


### Hypoxic induction of an HRE response in *Drosophila* S2 cells

In a previous report, we have shown that a dimerized 51-bp sequence derived from the murine lactate dehydrogenase-A (LDH-A) enhancer can drive hypoxia-dependent transcription of a *Gal4* reporter element in transgenic fly lines (Lavista-Llanos et al., 2002). To generate a cell culture system that enables pharmacological intervention, we generated an S2 cell line bearing an HRE reporter. The dimerized regulatory sequence of the LDH gene was cloned upstream of a firefly luciferase reporter gene bearing a fly Hsp70 minimal promoter (HRE-Luc) in a pGL3 plasmid (Fig. 1A). With this construct, stable

*Drosophila* S2 cell lines were generated and challenged with diverse hypoxic treatments. As depicted in Fig. 1B, exposure of the cells to 1% O<sub>2</sub> for 20 hours, elicited a 23-fold increase of luciferase activity, 2% O<sub>2</sub> caused a 9-fold increase and 3% O<sub>2</sub> led to 1.5-fold increase. These results show that the LDH regulatory element can drive hypoxia-dose-dependent transcription in *Drosophila* S2 cells, as it does in transgenic embryos (Lavista-Llanos et al., 2002). In mammalian cell culture systems, iron chelation was reported to mimic the effect of hypoxia, because prolyl hydroxylases involved in oxygen sensing have an absolute requirement of iron. In agreement with this notion, addition to the cells of the iron chelator DFO provoked a consistent 6-fold increase in luciferase activity (Fig. 1B), confirming that the HRE-Luc reporter drives a bona fide transcriptional response to hypoxia in the transfected S2 cell line. Exposure of the cells to 2% oxygen for 16 hours caused an increase in the expression of endogenous *dLDH* mRNA, which was detected by northern blot (Fig. 1C) and semi-quantitative RT-PCR (not shown). Induction of this hypoxia-responsive gene parallels induction of the HRE-Luc reporter, demonstrating that this element can report an endogenous physiological response to hypoxia. To confirm that HRE-Luc hypoxic induction is driven by the *Drosophila* HIF-1 homologue, we treated the cells with *sima* or *tango* dsRNA and, as expected, silencing of *sima* or *tango* expression caused complete abrogation of luciferase induction by hypoxia or DFO (Fig. 1D). Consistent with these results, transfection of an expression vector encoding Sima was sufficient to elicit a strong induction of the HRE-Luc reporter in a dose-dependent manner (Fig. 1E). We conclude from these experiments that, induction of the HRE-Luc reporter in the stable cell line depends on the *Drosophila* HIF-1 functional homologue, which is composed by the bHLH-PAS subunits Sima and Tgo.

### Insulin can induce an HRE response in S2 cells through the bHLH-PAS proteins Sima and Tango


Having characterized the behavior of the HRE-Luc reporter in the stably transfected *Drosophila* S2 cell line, we sought to investigate whether insulin can trigger the HRE response in this setting. Upon treating the cell culture with 50 µg/ml insulin, expression of the HRE-Luc reporter was increased about 20-fold, attaining levels similar to those observed upon hypoxia and higher levels to those recorded upon DFO treatment (Fig. 2A). Moreover, insulin induction of



**Fig. 1.** Hypoxic induction of an HRE-response in a stably transfected *Drosophila* S2 cell line. (A) Schematic representation of the HIF-responsive firefly luciferase reporter gene used in this study (HRE-Luc), in which transcription depends on a dimerized 51 bp sequence from the murine lactate dehydrogenase enhancer that contains two HIF responsive elements (HREs) and one cyclic AMP responsive element (CRE). (B) Effect of hypoxia on HRE-Luc activity in S2 cells. Cells were exposed to different oxygen levels for 20 hours or stimulated with 100 µM DFO. Luciferase activity was determined and normalized to protein concentration in the extracts. Induction of the reporter augmented when oxygen levels decrease. (C) Variations in *dLDH* mRNA levels. To measure the effect of hypoxia on endogenous *dLDH* gene expression, cells were exposed to 2% O<sub>2</sub> for 16 hours. Total RNA was then prepared and analyzed by northern blot. LDH mRNA is strongly induced by hypoxia (upper panel). 18S rRNA that was used as loading control remains constant (lower panel). (D) Role of Sima and Tango in HRE-Luc induction. Cells were incubated with dsRNA directed against *GFP* (control), *sima* or *tango* and 3 days later, the culture was treated or not with DFO (100 µM) or was exposed to 1% O<sub>2</sub> for 20 hours. Cells were then lysed and luciferase activity was determined. Hypoxia or DFO-dependent induction of the reporter was specifically abrogated by *sima* or *tango* RNAi treatment. (E) Effect of exogenous Sima. Cells were transiently transfected with an empty vector (pAC) or with a plasmid expressing Sima protein (pAC-Sima), together with 100 ng of a vector encoding Renilla luciferase (pRL). Twenty-four hours after transfection, cells were lysed and firefly or Renilla luciferase activity was determined. Induction of the HRE-Luc reporter was proportional to the transfected amount of *sima*-encoding plasmid. Normalized luciferase activity (firefly/Renilla ratio) is expressed as fold-induction with respect to the activity upon transfection of an empty vector. In all experiments involving luciferase determinations, data represent the mean±s.e.m. of three independent experiments performed in triplicate and expressed as fold-induction with respect to untreated cells.

the reporter was dose-dependent (Fig. 2B), suggesting that graded activation of an insulin-dependent pathway can render proportional induction of the HRE response. Northern blot analysis revealed that *dLDH* mRNA levels were also upregulated upon insulin treatment (Fig. 2C), suggesting that

**Fig. 2.** Insulin induces an HRE-response in S2 cells. (A) Effect of insulin on HRE-Luc reporter induction. S2 cells stably transfected with HRE-Luc were cultured for 3 days and then exposed or not to hypoxia (1% O<sub>2</sub>), DFO (100 μM) or insulin (50 μg/ml) for 20 hours. Luciferase activity was determined and normalized to protein concentration. HRE-Luc was induced by insulin, and attained levels similar to those upon exposure to hypoxia and higher than those observed upon DFO treatment. (B) Treatment with different doses of insulin. Cells were stimulated with 1-50 μg/ml insulin for 20 hours, lysed and luciferase activity was determined and normalized to protein concentration. HRE-Luc was induced as a function of insulin concentration. (C) Endogenous *dLDH* mRNA levels upon insulin treatment. To assess the effect of insulin on the expression of the endogenous *dLDH* gene, cells were treated for 4, 8, or 16 hours with insulin (10 μg/ml) and then total RNA was prepared and analyzed by northern blot using a specific *dLDH* probe (upper panel) or 18S rRNA as a control (lower panel); expression of *dLDH* mRNA increased as a function of the time of exposure to insulin. (D) Role of Sima and Tango in insulin-dependent induction of HRE-Luc. Cells were incubated in the absence (control) or presence of dsRNA directed against *GFP*, *sim* or *tango* and 3 days later, they were treated or not with 10 μg/ml insulin for 20 hours. Cells were then lysed and luciferase activity determined. Induction of the HRE reporter by insulin was totally abrogated by *sim* or *tango* dsRNA.



both stimuli, hypoxia and insulin, can induce common target genes in *Drosophila*. To assess the identity of the transcription factor involved in insulin-dependent induction of HRE-dependent transcription, we treated the cells with *sim* or *tango* RNAi. Major inhibition of HRE-Luc induction by insulin was observed in both cases, whereas a control *GFP* dsRNA did not have any effect (Fig. 2D). These results indicate that insulin-dependent induction of the HRE response is mediated by the Sima-Tango heterodimer.

#### Activation of the insulin-elicited HRE response depends on the PI3K-AKT pathway

It has been reported that activation of the *Drosophila* InR can trigger the induction of MAPK and PI3K signaling pathways (Oldham et al., 2002; Olivier et al., 1993). To assess which pathway transduces insulin-dependent induction of the HRE response in *Drosophila*, we treated the cells with LY294002, a specific inhibitor of PI3K. Insulin-triggered induction was strongly suppressed in a dose-dependent manner (Fig. 3A), suggesting that the PI3K signaling pathway is crucially involved in the transduction of the signal that results in HRE induction. Consistent with this, LY294002 treatment significantly reduced *dLDH* mRNA induction by insulin, confirming involvement of the PI3K-AKT pathway (Fig. 3B). By contrast, treatment with U0126, a specific inhibitor of MAPK kinase (MEK), did not have an effect on insulin-dependent HRE-Luc induction (data not shown), suggesting that the MAPK pathway does not contribute significantly to HRE induction by insulin. To gather additional evidence for a role of the PI3K-AKT pathway in the HRE response, we

silenced the expression of *dAKT* or *dPDK1* by RNAi treatment. Complete silencing of *dAKT* and *dPDK1* was attained upon addition of dsRNAs (data not shown) and, concomitantly, strong suppression of insulin-dependent induction of the HRE reporter was observed (Fig. 3C). These results are consistent with the notion that *dAKT* and *dPDK1* are both required for this transcriptional response. Further confirming that activation of the PI3K-AKT pathway can induce an HRE response, transfection of an expression vector encoding AKT elicited a fivefold induction of the HRE-Luc reporter (Fig. 3D). To obtain further evidence of an involvement of the PI3K-AKT pathway in the HRE response, we altered *dPTEN* levels. *PTEN* is a negative regulator of the PI3K-AKT pathway and *Drosophila PTEN* (*dPTEN*) mutants were shown to upregulate PtdIns(3,4,5)P<sub>3</sub> levels, leading to constitutive activation of the PI3K-AKT pathway (Oldham et al., 2002; Stocker et al., 2002). Consistent with this, *dPTEN* RNAi treatment caused a 5 fold induction of the HRE response in the absence of insulin stimulation (Fig. 3E), further demonstrating that activation of the PI3K-AKT pathway can elicit an HRE transcriptional response in S2 cells. We next wanted to test whether augmented activity of the PI3K signaling pathway can also induce an HRE response in vivo. To this end, we analyzed the expression of the hypoxia-responsive reporter *LDH-LacZ* in transgenic embryos that were mutant for *dPTEN* (*PTEN<sup>2LL17</sup>*). As previously described (Lavista-Llanos et al., 2002), the *LDH-LacZ* reporter element was silent in wild-type normoxic embryos (Fig. 4A) and induced upon hypoxic exposure in a characteristic expression pattern that corresponds to some of the developing tracheal branches (Fig. 4B). Interestingly, in *PTEN<sup>2LL17</sup>* homozygous mutant embryos the *LDH-LacZ*

reporter was clearly activated in normoxia (Fig. 4C), mimicking the expression pattern of wild-type embryos exposed to hypoxia (Fig. 4B,C). These results suggest that activation of the PI3K-AKT pathway can induce the HRE response not only in cell culture but also in vivo.


#### Activation of the PI3K-AKT pathway provokes upregulation of Sima protein but not of its mRNA

To gain insights into the molecular mechanism that mediate induction of insulin-dependent HRE response, we analyzed whether Sima protein levels were modified upon insulin stimulation. As depicted in Fig. 5A, western blot analysis revealed that Sima protein accumulated in insulin-treated cells, attaining levels comparable to those of cells exposed to hypoxia (2% O<sub>2</sub> for 16 hours). By contrast, northern blot analysis showed that *sima* mRNA levels remained constant after exposure to insulin (Fig. 5B), indicating that Sima protein levels are increased through a post-transcriptional mechanism. To analyze whether in *Drosophila* S2 cells insulin-dependent accumulation of Sima depends on TOR, we studied the effect of rapamycin on HRE-Luc induction. Indeed, rapamycin treatment provoked a strong inhibition of the HRE response (Fig. 5C) and, consistent with this, treatment of the cells with dsRNA for *TOR*, for its upstream regulator *Rheb* or for S6 kinase (*S6K*), which is known to be activated by TOR (Oldham and Hafen, 2003; Li et al., 2004; Saucedo et al., 2003; Stocker


et al., 2003; Zhang et al., 2003) led to complete abrogation of insulin-dependent induction of HRE-Luc (Fig. 5D). These results suggest that the insulin-PI3K pathway can modulate Sima protein levels through the activation of TOR.

#### Sima nucleo-cytoplasmic localization is developmentally modulated and depends on oxygen levels in a dose-dependent manner

We have shown above that insulin-PI3K stimulation leads to upregulation of Sima protein levels. Therefore, we sought to assess whether this pathway can regulate the subcellular localization of Sima. For comparison, it was important to first study the effect of oxygen levels on Sima localization in detail. In a previous report, we showed that it was possible to override rapid Sima degradation by overexpressing the protein in *Drosophila* transgenic embryos through an *engrailed*-Gal4 driver, and determined that Sima is mostly cytoplasmic in normoxia and accumulates preferentially in the nucleus in hypoxia (Lavista-Llanos et al., 2002). To better characterize this regulation, we used the same overexpression strategy (Brand and Perrimon, 1993) and for quantitative purposes, because Sima subcellular localization is not an all-or-none phenomenon, thirty or more embryos of each developmental stage were analyzed. Each observation was classified into one of the following three arbitrary categories of Sima subcellular localization: (a) cytoplasmic; (b) ubiquitous and (c) nuclear (Fig. 6A) (see supplementary material Tables 1-5, for statistical analysis). With this criterion, we first studied whether subcellular localization is influenced by developmental



**Fig. 3.** HRE-Luc induction in S2 cells is mediated by the activation of the PI3K-AKT pathway. (A) Effect of a PI3K inhibitor on the insulin-dependent HRE response. To assess the effect of PI3K inhibition on the insulin-stimulated HRE response, cells were treated for 20 hours with insulin (10 µg/ml) in the presence of LY294002 (◆) or the vehicle (DMSO) (△). LY294002 was added 1 hour prior to insulin stimulation and luciferase activity was determined and normalized to protein concentration. LY294002 provoked a dose-dependent inhibition of the HRE response. (B) Effect of LY294002 on *dLDH* mRNA induction. To study the effect of PI3K inhibition on *dLDH* mRNA increase after insulin stimulation, S2 cells were treated for 16 hours with insulin (10 µg/ml) in the presence or not of 10 µM LY294002. Total RNA was prepared and analyzed by northern blot using a specific *dLDH* probe (upper panel) or 18S rRNA as a control (lower panel). Treatment with LY294002 provoked a clear reduction of *dLDH* mRNA levels. (C) Role of dAKT and dPDK1 in HRE-Luc induction by insulin. Cells were incubated in the presence or absence of dsRNA directed against *GFP*, *dAKT* or *dPDK1*. After 3 days, the cells were treated or not with insulin 10 µg/ml for 20 hours, lysed and luciferase activity was determined. *dAKT* and *dPDK1* dsRNAs caused strong inhibition of HRE-Luc stimulation by insulin. (D) Effect of exogenous AKT on HRE-Luc expression. Cells were transiently transfected with a plasmid expressing AKT (pAC-AKT, 250 ng), together with 100 ng of a plasmid expressing Renilla luciferase (pRL). Twenty-four hours after transfection, cells were lysed and luciferase activity was determined. Results are shown as the ratio of firefly to Renilla luciferase activities; transfection of AKT led to strong induction of the HRE reporter. (E) Effect of dPTEN silencing. Cells were incubated in the presence or absence of dsRNA directed against *GFP* or *dPTEN*, lysed and luciferase activity was determined. dPTEN dsRNA treatment provoked strong induction of the HRE response.



**Fig. 4.** *PTEN*<sup>2L117</sup> homozygous embryos show constitutive induction of the transgenic HRE reporter LDH-*LacZ*. (A) Normoxic wild-type embryos. *Drosophila* wild-type embryos maintained in normoxia do not express the hypoxia-inducible reporter LDH-*LacZ*, as revealed by X-gal staining. (B) Hypoxic wild-type embryos. Embryos were exposed to 3% O<sub>2</sub> for 4 hours, fixed and stained with X-gal. Induction of the LDH-*LacZ* reporter was observed in an expression pattern that is coincident with some of the tracheal branches (arrow) (Lavista-Llanos et al., 2002). (C) Effect of *PTEN* loss-of-function on HRE induction in vivo. In *PTEN*<sup>2L117</sup> homozygous embryos the LDH-*LacZ* reporter was induced in normoxia (arrow) in a pattern that paralleled induction of the reporter in hypoxic wild-type embryos (compare with B).


parameters in normoxia. As depicted in Fig. 6B, in normoxia (21% O<sub>2</sub>) Sima was completely cytoplasmic at embryonic stages 11-14 but later in embryogenesis, at stage 16, a proportion of the embryos showed ubiquitous (61.8%) or nuclear (14.7%) localization, indicating that the subcellular


localization of Sima is developmentally modulated. Next, we addressed whether localization depends on graded hypoxia and performed the analysis at different oxygen levels throughout development. We observed that, indeed at all stages, Sima became gradually more nuclear when oxygen levels decreased (Fig. 6B) and, confirming developmental modulation, at embryonic stage 11 the protein tended to be more cytoplasmic and became gradually more nuclear as embryogenesis progressed (Fig. 6B). We therefore conclude that the subcellular localization of Sima depends on oxygen levels in a dose-dependent manner and that this regulation is under developmental control.

#### Activation of the PI3K pathway promotes Sima nuclear localization

To test whether Sima subcellular localization also depends on the PI3K-AKT pathway, we initially analyzed the effect of ectopic expression of dAKT. In transgenic embryos that overexpress dAKT under the control of the *engrailed*-Gal4 driver in normoxia, Sima localized in the nucleus in a much larger proportion than in wild-type embryos all throughout development (Fig. 6C). When dAKT was overexpressed together with dPDK1, an additional enhancement of Sima nuclear localization was observed (Fig. 6C), further suggesting that activation of the pathway can induce the HRE response at this level. Moreover, in *PTEN*<sup>2L117</sup> homozygous mutant embryos, Sima nuclear localization was also increased, and overexpression of dPTEN in a mutant *PTEN*<sup>2L117</sup> background completely reverted Sima localization to wild-type levels (Fig. 6C). Altogether, these results strongly suggest that, activation of the PI3K-AKT pathway promotes nuclear localization of HIF $\alpha$ /Sima, thereby contributing to the induction of an HRE-dependent transcriptional response in *Drosophila* cells.

**Fig. 5.** Insulin treatment provokes an increase in Sima protein levels; involvement of TOR in the HRE response. (A) Sima protein levels upon insulin treatment. S2 cells were treated with insulin (10  $\mu$ g/ml) (I) or exposed to hypoxia (2% O<sub>2</sub>) for 16 hours (H) and Sima protein levels in nuclear extracts were analyzed by western blot (upper panel), hsp70 (lower panel) was used as a loading control. Sima protein levels were strongly upregulated upon insulin treatment, attaining levels similar to those caused by exposure to 2% O<sub>2</sub> during 16 hours (results of a representative experiment ( $n=3$ ) are shown). (B) *sim*a mRNA upon insulin treatment. Total RNA from cells treated or not with insulin (10  $\mu$ g/ml) was analysed by northern blot using a specific *sim*a probe (upper panel), S18 rRNA was used as a loading control (lower panel). *sim*a mRNA levels did not change upon insulin treatment. (C) Effect of TOR on insulin-dependent induction of HRE-Luc. Cells were treated or not with insulin (10  $\mu$ g/ml) for 20 hours in the presence of rapamycin (200 nM). Luciferase activity was determined and normalized to protein concentration; rapamycin provoked clear inhibition of the of HRE-Luc induction. (D) Silencing of *dTOR*, *dRheb* or *dS6K*. Cells were incubated in the presence of dsRNA directed against *GFP*, *dTOR*, *dRheb* or *dS6K* and 3 days later the cells were treated or not with 10  $\mu$ g/ml insulin for 20 hours, then lysed and luciferase activity was determined; *dTOR*, *dRheb* and *dS6K* dsRNAs provoked strong inhibition of HRE induction.





**Fig. 6.** Hypoxia or activation of the PI3K-AKT pathway promote nuclear localization of Sima in vivo. (A) Definition of three categories of Sima subcellular localization. Sima and a  $\beta$ -galactosidase ( $\beta$ -gal) variant bearing a nuclear localization signal were simultaneously expressed in UAS transgenic embryos under control of an *engrailed-Gal4* driver. Embryos were fixed, subjected to double  $\alpha$ -Sima (red)/ $\alpha$ - $\beta$ gal (green) immunofluorescence and observed at the confocal microscope. Three arbitrary categories of subcellular localization were defined: 'Cytoplasmic', where more than 90% of the cells in the embryo exhibited cytoplasmic localization; 'Nuclear', where more than 90% of the cells showed nuclear localization and 'Ubiquitous', where less than 90% of the cells showed cytoplasmic or nuclear localization. (B) Sima subcellular localization throughout development at different oxygen concentrations (white, Cytoplasmic; gray, Ubiquitous; black, Nuclear). Overnight egg-laying plates were exposed to different oxygen concentrations for 4 hours and then fixed and immunostained. In normoxia, at embryonic stages 11-14, Sima protein was exclusively cytoplasmic, whereas later in embryogenesis Sima became gradually more nuclear. As oxygen concentration decreased Sima protein became progressively more nuclear. (C) Effect of the activation of the PI3K-AKT pathway on Sima subcellular localization (white, Cytoplasmic; gray, Ubiquitous; black, Nuclear). Overnight egg-laying plates with normoxic embryos of the indicated genotype were fixed and immunostained. *engrailed*-driven overexpression of dAKT led to an increase in Sima nuclear localization at stages 15-16. Introduction of an EP element overexpressing dPDK1 along with dAKT enhanced Sima nuclear localization in comparison with embryos overexpressing dAKT alone. *dPTEN*<sup>2L117</sup> homozygous mutant embryos also showed increased nuclear localization; confirming that loss of dPTEN was indeed the cause of increased nuclear localization of Sima, ectopic expression of dPTEN in the *dPTEN*<sup>2L117</sup> homozygous mutant background rescued wild-type localization of Sima.

## Discussion

### Regulation of Sima through the PI3K-AKT pathway

We have used a combination of biochemical and genetic strategies to show both in cell culture and in vivo that activation of the PI3K-AKT pathway promotes HIF-dependent transcription in *Drosophila* cells. We demonstrated that insulin stimulation or exposure to hypoxia can induce common target genes, and that such transcriptional response depends on the *Drosophila* HIF $\alpha$  and HIF $\beta$  homologues Sima and Tango. We provided evidence that insulin-stimulated HRE response is transduced by the PI3K-AKT pathway and, furthermore, that the effect depends on TOR and involves an increase in Sima protein levels, whereas mRNA levels are not affected. Our results are in good agreement with the reported effect of the PI3K-AKT/TOR pathway on mammalian HIF, because in several cell lines activation of this pathway led to an increase in HIF protein levels (Zundel et al., 2000) caused by augmented translation (Fukuda et al., 2002; Laughner et al., 2001; Treins et al., 2002) or stabilization of the protein (Hudson et al., 2002).

In addition, we have shown that the subcellular localization of Sima depends on oxygen tension in a dose-dependent

manner and that activation of the PI3K-AKT pathway also causes a major increase in Sima nuclear localization. This regulatory mechanism might represent another conserved aspect of HIF regulation, because one recent report suggests that HIF- $\alpha$  accumulates in the nucleus of retinal epithelial cells upon IGF-1 $\alpha$  treatment (Treins et al., 2005). The molecular bases of HIF $\alpha$  nuclear accumulation upon hypoxia (Kallio et al., 1998) or PI3K activation are so far unclear. Nucleo-cytoplasmic localization of many transcription factors results from a steady-state equilibrium between nuclear import and nuclear export, and accumulation in one or the other compartment depends on the relative rate of import versus export (Fabbro and Henderson, 2003; Swameye et al., 2003). Whether HIF nuclear accumulation upon hypoxia or growth factor stimulation depends on regulated nuclear import or regulated nuclear export remains to be determined.

### Conservation of HIF regulation in evolution

We have previously shown that all major cellular features of oxygen-dependent regulation of HIF proteins are conserved in *Drosophila* (Lavista-Llanos et al., 2002) and, thus, activation of the HRE response in *Drosophila* by the PI3K-AKT and TOR pathways extends even further the notion of a conserved HIF system in evolution. The functional significance of the regulation exerted by PI3K-AKT and TOR pathways over the HRE response was previously discussed in the context of cancer biology (Brugarolas and Kaelin, 2004; Hudson et al.,



2002; Jiang et al., 2001; Zhong et al., 2000; Zundel et al., 2000), because the loss of PTEN or the tuberous sclerosis complex 1 (TSC1) and TSC2 proteins is frequently associated to human tumors (Di Cristofano and Pandolfi, 2000; Montagne et al., 2001; Trotman et al., 2003). What is the functional meaning of PI3K-AKT pathway regulation of the HRE response in normal cells? The PI3K-AKT and TOR pathways are regulated in part by growth factors and other endocrine signals and thus, endocrine control of the HRE response is conserved among animal species that have diverged 700 million years ago. It seems reasonable to postulate that the main physiological role of HRE induction by the PI3K-AKT pathway is the stimulation of glycolysis, but a function in the regulation of animal body size and growth control is another interesting possibility.

#### Role of the PI3K-AKT and TOR pathways in growth control and metabolism

A cardinal function of the PI3K-AKT and TOR pathways throughout evolution is to regulate growth, and to determine the final size of developing organs and whole organisms (Garofalo, 2002; Oldham and Hafen, 2003). Genetic studies in *Drosophila* have shown that a reduction of the activity of the PI3K-AKT pathway results in flies with a reduced body size, bearing smaller cells (Bohni et al., 1999; Goberdhan et al., 1999; Huang et al., 1999; Rintelen et al., 2001). Likewise, a reduction in TOR signaling provokes growth decrease (Oldham et al., 2000; Zhang et al., 2000) and, conversely, over-activation of TOR signaling due to loss-of-function of its negative regulators TSC1 and/or TSC2, leads to an increase in cell and body size (Gao and Pan, 2001; Potter et al., 2001). The effect of TOR on cell growth was reported to be mediated at least in part by S6K, a kinase that phosphorylates the ribosomal protein S6, leading to translational activation (Gingras et al., 2004; Miron et al., 2003).


Besides its role in growth control, the insulin-PI3K-AKT pathway has been traditionally implicated in the regulation of circulating glucose levels and anabolic metabolism (Saltiel and Kahn, 2001). It was recently demonstrated that the cellular bases of glucose sensing and regulation of serum glucose are conserved between mammals and *Drosophila* (Kim and Rulifson, 2004; Rulifson et al., 2002), and it has been proposed that PI3K-AKT signaling in conjunction with the TOR pathway coordinates growth according to environmental conditions and the nutritional status of the organism (Britton et al., 2002). The mechanism involved in this coordination is still unclear (Jacinto and Hall, 2003; Oldham and Hafen, 2003). Oxygen tension is one environmental factor that has been shown to modulate growth in *Drosophila*, because hypoxic flies have a reduced body size (Frazier et al., 2001). We recently provided a mechanistic explanation to this phenomenon, by showing that overexpression of Sima protein causes a reduction in cell size in an autonomous manner (Centanin et al., 2005). Consistent with this, it has previously been shown that hypoxia provokes a reduction in *Drosophila* TOR pathway activity and that such reduction results from hyperactivation of the TSC1-TSC2 tumor suppressor complex (Reiling and Hafen, 2004). In a simultaneous publication, Brugarolas et al. reported similar results in mammalian cells, implying that TOR regulation by hypoxia (Arsham et al., 2003)

is an evolutionary conserved feature (Brugarolas et al., 2004). Furthermore, both studies (Brugarolas et al., 2004; Reiling and Hafen, 2004) demonstrated that hypoxia-dependent TSC1-TSC2 stimulation and growth inhibition are mediated by the product of a HIF/Sima-inducible gene called *scylla* in *Drosophila* and *RTP801/REDD1* in mammals.

#### Sima/HIF as a modulator of animal growth

The results presented here establish a direct link between pathways largely implicated in growth regulation (PI3K-AKT and TOR) and the hypoxia-responsive machinery (HIF/Sima). A mechanistic explanation for hypoxia-dependent inhibition of growth is depicted in Fig. 7. In hypoxia, HIF prolyl hydroxylase (Hph)/Fatiga activity is reduced, resulting in HIF/Sima stabilization and induction of an HRE response. One of the genes induced by hypoxia is *scylla/RTP801/REDD1*, that in turn activates TSC1-TSC2. Then, stimulation of the TSC complex provokes reduction of TOR activity and decreased S6K phosphorylation, resulting in growth inhibition (Brugarolas et al., 2004; Reiling and Hafen, 2004). According to this model, PI3K-TOR activation of HIF- $\alpha$ /Sima might generate a negative feedback loop to limit or downregulate growth; in this scenario, low oxygen levels are expected to enhance Sima-dependent inhibition of growth.

Frei et al. have recently reported that mitochondrial dysfunction inhibits Hph/Fatiga activity, thereby triggering the transcriptional response to hypoxia, and also that it concomitantly provokes growth defects (Frei et al., 2005). They propose that Hph/Fatiga operates as an integration node between oxygen levels and growth regulation (Frei and Edgar, 2004; Frei et al., 2005). Our results have shown that the effect of Hph/Fatiga on growth regulation is conveyed at least in part



**Fig. 7.** Model of insulin signaling through HIF/Sima. The activated InR can signal through AKT to prevent TSC1-TSC2 from inhibiting TOR, thereby activating S6K leading to growth stimulation. In addition, TOR can activate Sima/HIF-1 that induces transcription of several target genes, including *scylla/RTP801* that in turn stimulates TSC1-TSC2, resulting in a negative feedback loop that reduces TOR activity. Activated AKT can stimulate nuclear localization of Sima/HIF-1 through an unknown mechanism (dashed line), contributing to the negative regulation of TOR. Low oxygen levels can inhibit growth by reducing the activity of Hph/Fatiga, which provokes stabilization of HIF- $\alpha$ /Sima and, in consequence, induction of *scylla/RTP801*, thereby reducing TOR activity.

by Sima (Centanin et al., 2005). Further studies will reveal whether Hph/Fatiga also plays Sima-independent roles in cell and organ size determination.

We wish to thank Thomas Gorr for advice on the western blots, Juan Jose Fanara for help with statistical analysis, the *Drosophila* Bloomington Stock Center for fly stocks, members of Wappner lab for discussion and Graciela Boccaccio for critical reading of the manuscript. This work was supported by Wellcome Trust Grant 070161/Z/03/Z, Universidad de Buenos Aires grant X411 and ANPCyT 01-10839. A.D. is a fellow of ANPCyT and P.W. is a career investigator of CONICET.

## References

- Arsham, A. M., Howell, J. J. and Simon, M. C. (2003). A novel hypoxia-inducible factor-independent hypoxic response regulating mammalian target of rapamycin and its targets. *J. Biol. Chem.* **278**, 29655-29660.
- Bacon, N. C., Wappner, P., O'Rourke, J. F., Bartlett, S. M., Shilo, B., Pugh, C. W. and Ratcliffe, P. J. (1998). Regulation of the *Drosophila* bHLH-PAS protein Sima by hypoxia: functional evidence for homology with mammalian HIF-1 alpha. *Biochem. Biophys. Res. Commun.* **249**, 811-816.
- Bohni, R., Riesgo-Escovar, J., Oldham, S., Brogiolo, W., Stocker, H., Andruss, B. F., Beckingham, K. and Hafen, E. (1999). Autonomous control of cell and organ size by CHICO, a *Drosophila* homolog of vertebrate IRS1-4. *Cell* **97**, 865-875.
- Brand, A. H. and Perrimon, N. (1993). Targeted gene expression as means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-415.
- Britton, J. S., Lockwood, W. K., Li, L., Cohen, S. M. and Edgar, B. A. (2002). *Drosophila*'s insulin/PI3-kinase pathway coordinates cellular metabolism with nutritional conditions. *Dev. Cell* **2**, 239-249.
- Brugarolas, J. and Kaelin, W. G., Jr (2004). Dysregulation of HIF and VEGF is a unifying feature of the familial hamartoma syndromes. *Cancer Cell* **6**, 7-10.
- Brugarolas, J., Lei, K., Hurley, R. L., Manning, B. D., Reiling, J. H., Hafen, E., Witters, L. A., Ellisen, L. W. and Kaelin, W. G., Jr (2004). Regulation of mTOR function in response to hypoxia by REDD1 and the TSC1/TSC2 tumor suppressor complex. *Genes Dev.* **18**, 2893-2904.
- Bruick, R. K. (2003). Oxygen sensing in the hypoxic response pathway: regulation of the hypoxia-inducible transcription factor. *Genes Dev.* **17**, 2614-2623.
- Bruick, R. K. and McKnight, S. L. (2001). A conserved family of prolyl-4-hydroxylases that modify HIF. *Science* **294**, 1337-1340.
- Centanin, L., Ratcliffe, P. J. and Wappner, P. (2005). Reversion of lethality and growth defects in Fatiga oxygen-sensor mutant flies by loss of Hypoxia-Inducible Factor-alpha/Sima. *EMBO Rep.* (in press).
- Clemens, J. C., Worby, C. A., Simonson-Leff, N., Muda, M., Maehama, T., Hemmings, B. A. and Dixon, J. E. (2000). Use of double-stranded RNA interference in *Drosophila* cell lines to dissect signal transduction pathways. *Proc. Natl. Acad. Sci. USA* **97**, 6499-6503.
- Dann, C. E., 3rd, Bruick, R. K. and Deisenhofer, J. (2002). Structure of factor-inhibiting hypoxia-inducible factor 1. An asparaginyl hydroxylase involved in the hypoxic response pathway. *Proc. Natl. Acad. Sci. USA* **99**, 15351-15356.
- Darlington, T. K., Wager-Smith, K., Ceriani, M. F., Staknis, D., Gekakis, N., Steeves, T. D., Weitz, C. J., Takahashi, J. S. and Kay, S. A. (1998). Closing the circadian loop: CLOCK-induced transcription of its own inhibitors per and tim. *Science* **280**, 1599-1603.
- Di Cristofano, A. and Pandolfi, P. P. (2000). The multiple roles of PTEN in tumor suppression. *Cell* **100**, 387-390.
- Elkins, J. M., Hewitson, K. S., McNeill, L. A., Seibel, J. F., Schlemminger, I., Pugh, C. W., Ratcliffe, P. J. and Schofield, C. J. (2002). Structure of factor inhibiting HIF reveals mechanism of oxidative modification of HIF-1alpha. *J. Biol. Chem.* **278**, 1802-1806.
- Epstein, A. C., Gleadle, J. M., McNeill, L. A., Hewitson, K. S., O'Rourke, J., Mole, D. R., Mukherji, M., Metzen, E., Wilson, M. I., Dhanda, A. et al. (2001). *C. elegans* EGL-9 and Mammalian Homologs Define a Family of Dioxygenases that Regulate HIF by Prolyl Hydroxylation. *Cell* **107**, 43-54.
- Fabbro, M. and Henderson, B. R. (2003). Regulation of tumor suppressors by nuclear-cytoplasmic shuttling. *Exp. Cell Res.* **282**, 59-69.
- Feldser, D., Agani, F., Iyer, N. V., Pak, B., Ferreira, G. and Semenza, G. L. (1999). Reciprocal positive regulation of hypoxia-inducible factor 1alpha and insulin-like growth factor 2. *Cancer Res.* **59**, 3915-3918.
- Fernandez, R., Tabarini, D., Azpiazu, N., Frasch, M. and Schlessinger, J. (1995). The *Drosophila* insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential. *EMBO J.* **14**, 3373-3384.
- Frazier, M. R., Woods, H. A. and Harrison, J. F. (2001). Interactive effects of rearing temperature and oxygen on the development of *Drosophila melanogaster*. *Physiol. Biochem. Zool.* **74**, 641-650.
- Frei, C. and Edgar, B. A. (2004). *Drosophila* cyclin D/Cdk4 requires Hif-1 prolyl hydroxylase to drive cell growth. *Dev. Cell* **6**, 241-251.
- Frei, C., Galloni, M., Hafen, E. and Edgar, B. A. (2005). The *Drosophila* mitochondrial ribosomal protein mRpL12 is required for Cyclin D/Cdk4-driven growth. *EMBO J.* **24**, 623-634.
- Fukuda, R., Hirota, K., Fan, F., Jung, Y. D., Ellis, L. M. and Semenza, G. L. (2002). Insulin-like growth factor 1 induces hypoxia-inducible factor 1-mediated vascular endothelial growth factor expression, which is dependent on MAP kinase and phosphatidylinositol 3-kinase signaling in colon cancer cells. *J. Biol. Chem.* **277**, 38205-38211.
- Fukuda, R., Kelly, B. and Semenza, G. L. (2003). Vascular endothelial growth factor gene expression in colon cancer cells exposed to prostaglandin E2 is mediated by hypoxia-inducible factor 1. *Cancer Res.* **63**, 2330-2334.
- Gao, X. and Pan, D. (2001). TSC1 and TSC2 tumor suppressors antagonize insulin signaling in cell growth. *Genes Dev.* **15**, 1383-1392.
- Garofalo, R. S. (2002). Genetic analysis of insulin signaling in *Drosophila*. *Trends Endocrinol. Metab.* **13**, 156-162.
- Garofalo, R. S. and Rosen, O. M. (1988). Tissue localization of *Drosophila melanogaster* insulin receptor transcripts during development. *Mol. Cell. Biol.* **8**, 1638-1647.
- Gingras, A. C., Raught, B. and Sonenberg, N. (2004). mTOR signaling to translation. *Curr. Top. Microbiol. Immunol.* **279**, 169-197.
- Goberdhan, D. C., Paricio, N., Goodman, E. C., Mlodzik, M. and Wilson, C. (1999). *Drosophila* tumor suppressor PTEN controls cell size and number by antagonizing the Chico/PI3-kinase signaling pathway. *Genes Dev.* **13**, 3244-3258.
- Gorr, T. A., Tomita, T., Wappner, P. and Bunn, H. F. (2004). Regulation of *Drosophila* hypoxia-inducible factor (HIF) activity in SL2 cells: identification of a hypoxia-induced variant isoform of the HIFalpha homolog gene similar. *J. Biol. Chem.* **279**, 36048-36058.
- Hewitson, K. S., McNeill, L. A., Riordan, M. V., Tian, Y. M., Bullock, A. N., Welford, R. W., Elkins, J. M., Oldham, N. J., Bhattacharya, S., Gleadle, J. M. et al. (2002). Hypoxia-inducible factor (HIF) asparagine hydroxylase is identical to factor inhibiting HIF (FIH) and is related to the cupin structural family. *J. Biol. Chem.* **277**, 26351-26355.
- Huang, H., Potter, C. J., Tao, W., Li, D. M., Brogiolo, W., Hafen, E., Sun, H. and Xu, T. (1999). PTEN affects cell size, cell proliferation and apoptosis during *Drosophila* eye development. *Development* **126**, 5365-5372.
- Hudson, C. C., Liu, M., Chiang, G. G., Otterness, D. M., Loomis, D. C., Kaper, F., Giaccia, A. J. and Abraham, R. T. (2002). Regulation of hypoxia-inducible factor 1alpha expression and function by the mammalian target of rapamycin. *Mol. Cell. Biol.* **22**, 7004-7014.
- Ivan, M., K. K., Yang, H., Kim, W., Valiando, J., Ohh, M., Salic, A., Asara, J. M., Lane, W. S., Kaelin, W. S., Jr (2001). HIFalpha targeted for VHL-mediated destruction by proline hydroxylation: implications for oxygen sensing. *Science* **292**, 464-468.
- Jaakkola, P. M. D., Tian, Y. M., Wilson, M. I., Gielbert, J., Gaskell, S. J., Kriegsheim, A. V., Heberstreit, H. F., Mukherji, M., Schofield, C. J., Maxwell, P. H. et al. (2001). Targeting of HIF-alpha to the Von Hippel Lindau ubiquitylation complex by O2-regulated prolyl hydroxylation. *Science* **292**, 468-472.
- Jacinto, E. and Hall, M. N. (2003). Tor signalling in bugs, brain and brawn. *Nat. Rev. Mol. Cell. Biol.* **4**, 117-126.
- Jiang, B. H., Jiang, G., Zheng, J. Z., Lu, Z., Hunter, T. and Vogt, P. K. (2001). Phosphatidylinositol 3-kinase signaling controls levels of hypoxia-inducible factor 1. *Cell Growth Diff.* **12**, 363-369.
- Kallio, P. J., Okamoto, K., O'Brien, S., Carrero, P., Makino, Y., Tanaka, H. and Poellinger, L. (1998). Signal transduction in hypoxic cells: inducible nuclear translocation and recruitment of the CBP/p300 coactivator by the hypoxia-inducible factor-1alpha. *EMBO J.* **17**, 6573-6586.
- Kasuno, K., Takabuchi, S., Fukuda, K., Kizaka-Kondoh, S., Yodoi, J., Adachi, T., Semenza, G. L. and Hirota, K. (2004). Nitric oxide induces hypoxia-inducible factor 1 activation that is dependent on MAPK and phosphatidylinositol 3-kinase signaling. *J. Biol. Chem.* **279**, 2550-2558.

- Kietzmann, T., Samoylenko, A., Roth, U. and Jungermann, K. (2003). Hypoxia-inducible factor-1 and hypoxia response elements mediate the induction of plasminogen activator inhibitor-1 gene expression by insulin in primary rat hepatocytes. *Blood* **101**, 907-914.
- Kim, S. K. and Rulifson, E. J. (2004). Conserved mechanisms of glucose sensing and regulation by *Drosophila* corpora cardiaca cells. *Nature* **431**, 316-320.
- Lando, D., Peet, D. J., Gorman, J. J., Whelan, D. A., Whitelaw, M. L. and Bruick, R. K. (2002). FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional activity of hypoxia-inducible factor. *Genes Dev.* **16**, 1466-1471.
- Laughner, E., Taghavi, P., Chiles, K., Mahon, P. C. and Semenza, G. L. (2001). HER2 (neu) signaling increases the rate of hypoxia-inducible factor 1 $\alpha$  (HIF-1 $\alpha$ ) synthesis: novel mechanism for HIF-1-mediated vascular endothelial growth factor expression. *Mol. Cell. Biol.* **21**, 3995-4004.
- Lavista-Llanos, S., Centanin, L., Irisarri, M., Russo, D. M., Gleadle, J. M., Bocca, S. N., Muzzopappa, M., Ratcliffe, P. J. and Wappner, P. (2002). Control of the hypoxic response in *Drosophila* melanogaster by the basic helix-loop-helix PAS protein similar. *Mol. Cell. Biol.* **22**, 6842-6853.
- LeRoith, D., Lesniak, M. A. and Roth, J. (1981). Insulin in insects and annelids. *Diabetes* **30**, 70-76.
- Li, Y., Inoki, K. and Guan, K. L. (2004). Biochemical and functional characterizations of small GTPase Rheb and TSC2 GAP activity. *Mol. Cell. Biol.* **24**, 7965-7975.
- Luo, J. C. and Shibuya, M. (2001). A variant of nuclear localization signal of bipartite-type is required for the nuclear translocation of hypoxia inducible factors (1 $\alpha$ , 2 $\alpha$  and 3 $\alpha$ ). *Oncogene* **20**, 1435-1444.
- Maxwell, P. H., Pugh, C. W. and Ratcliffe, P. J. (1993). Inducible operation of the erythropoietin 3' enhancer in multiple cell lines: evidence for a widespread oxygen-sensing mechanism. *Proc. Natl. Acad. Sci. USA* **90**, 2423-2427.
- Maxwell, P. H., Wiesener, M. S., Chang, G. W., Clifford, S. C., Vaux, E. C., Cockman, M. E., Wykoff, C. C., Pugh, C. W., Maher, E. R. and Ratcliffe, P. J. (1999). The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis [see comments]. *Nature* **399**, 271-275.
- Miele, C., Rochford, J. J., Filippa, N., Giorgetti-Peraldi, S. and Van Obberghen, E. (2000). Insulin and insulin-like growth factor-I induce vascular endothelial growth factor mRNA expression via different signaling pathways. *J. Biol. Chem.* **275**, 21695-21702.
- Miron, M., Lasko, P. and Sonenberg, N. (2003). Signaling from Akt to FRAP/TOR targets both 4E-BP and S6K in *Drosophila* melanogaster. *Mol. Cell. Biol.* **23**, 9117-9126.
- Montagne, J., Radimerski, T. and Thomas, G. (2001). Insulin signaling: lessons from the *Drosophila* tuberous sclerosis complex, a tumor suppressor. *Sci. STKE* **2001**, PE36.
- Oldham, S. and Hafen, E. (2003). Insulin/IGF and target of rapamycin signaling: a TOR de force in growth control. *Trends Cell. Biol.* **13**, 79-85.
- Oldham, S., Montagne, J., Radimerski, T., Thomas, G. and Hafen, E. (2000). Genetic and biochemical characterization of dTOR, the *Drosophila* homolog of the target of rapamycin. *Genes Dev.* **14**, 2689-2694.
- Oldham, S., Stocker, H., Laffargue, M., Wittwer, F., Wymann, M. and Hafen, E. (2002). The *Drosophila* insulin/IGF receptor controls growth and size by modulating PtdIns(3) levels. *Development* **129**, 4103-4109.
- Olivier, J. P., Raabe, T., Henkemeyer, M., Dickson, B., Mbamalu, G., Margolis, B., Schlessinger, J., Hafen, E. and Pawson, T. (1993). A *Drosophila* SH2-SH3 adaptor protein implicated in coupling the sevenless tyrosine kinase to an activator of Ras guanine nucleotide exchange, Sos. *Cell* **73**, 179-191.
- Potter, C. J., Huang, H. and Xu, T. (2001). *Drosophila* Tsc1 functions with Tsc2 to antagonize insulin signaling in regulating cell growth, cell proliferation, and organ size. *Cell* **105**, 357-368.
- Reiling, J. H. and Hafen, E. (2004). The hypoxia-induced paralogs Scylla and Charybdis inhibit growth by down-regulating S6K activity upstream of TSC in *Drosophila*. *Genes Dev.* **18**, 2879-2892.
- Rintelen, F., Stocker, H., Thomas, G. and Hafen, E. (2001). PDK1 regulates growth through Akt and S6K in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **98**, 15020-15025.
- Roth, U., Curth, K., Unterman, T. G. and Kietzmann, T. (2004). The transcription factors HIF-1 and HNF-4 and the coactivator p300 are involved in insulin-regulated glucokinase gene expression via the phosphatidylinositol 3-kinase/protein kinase B pathway. *J. Biol. Chem.* **279**, 2623-2631.
- Rulifson, E. J., Kim, S. K. and Nusse, R. (2002). Ablation of insulin-producing neurons in flies: growth and diabetic phenotypes. *Science* **296**, 1118-1120.
- Saltiel, A. R. and Kahn, C. R. (2001). Insulin signalling and the regulation of glucose and lipid metabolism. *Nature* **414**, 799-806.
- Saucedo, L. J., Gao, X., Chiarelli, D. A., Li, L., Pan, D. and Edgar, B. A. (2003). Rheb promotes cell growth as a component of the insulin/TOR signalling network. *Nat. Cell Biol.* **5**, 566-571.
- Schofield, C. J. and Ratcliffe, P. J. (2004). Oxygen sensing by HIF hydroxylases. *Nat. Rev. Mol. Cell. Biol.* **5**, 343-354.
- Seecof, R. L. and Dewhurst, S. (1974). Insulin is a *Drosophila* hormone and acts to enhance the differentiation of embryonic *Drosophila* cells. *Cell Diff.* **3**, 63-70.
- Stocker, H., Andjelkovic, M., Oldham, S., Laffargue, M., Wymann, M. P., Hemmings, B. A. and Hafen, E. (2002). Living with lethal PIP3 levels: viability of flies lacking PTEN restored by a PH domain mutation in Akt/PKB. *Science* **295**, 2088-2091.
- Stocker, H., Radimerski, T., Schindelhof, B., Wittwer, F., Belawat, P., Daram, P., Breuer, S., Thomas, G. and Hafen, E. (2003). Rheb is an essential regulator of S6K in controlling cell growth in *Drosophila*. *Nat. Cell. Biol.* **5**, 559-565.
- Swameye, I., Muller, T. G., Timmer, J., Sandra, O. and Klingmuller, U. (2003). Identification of nucleocytoplasmic cycling as a remote sensor in cellular signaling by databased modeling. *Proc. Natl. Acad. Sci. USA* **100**, 1028-1033.
- Treins, C., Giorgetti-Peraldi, S., Murdaca, J., Semenza, G. L. and Van Obberghen, E. (2002). Insulin stimulates hypoxia-inducible factor 1 through a phosphatidylinositol 3-kinase/target of rapamycin-dependent signaling pathway. *J. Biol. Chem.* **277**, 27975-27981.
- Treins, C., Giorgetti-Peraldi, S., Murdaca, J., Monthouel-Kartmann, M. N. and Van Obberghen, E. (2005). Regulation of HIF-1 activity and expression of HIF hydroxylases in response to IGF-1. *Mol. Endocrinol.* **19**, 1304-1317.
- Trotman, L. C., Niki, M., Dotan, Z. A., Koutcher, J. A., Di Cristofano, A., Xiao, A., Khoo, A. S., Roy-Burman, P., Greenberg, N. M., Van Dyke, T. et al. (2003). Pten dose dictates cancer progression in the prostate. *PLoS Biol.* **1**, E59.
- Wang, G. L. and Semenza, G. L. (1993). General involvement of hypoxia-inducible factor 1 in transcriptional response to hypoxia. *Proc. Natl. Acad. Sci. USA* **90**, 4304-4308.
- Wang, G. L., Jiang, B. H., Rue, E. A. and Semenza, G. L. (1995). Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O<sub>2</sub> tension. *Proc. Natl. Acad. Sci. USA* **92**, 5510-5514.
- Yenush, L. and White, M. F. (1997). The IRS-signalling system during insulin and cytokine action. *BioEssays* **19**, 491-500.
- Yenush, L., Fernandez, R., Myers, M. G., Jr, Grammer, T. C., Sun, X. J., Blenis, J., Pierce, J. H., Schlessinger, J. and White, M. F. (1996). The *Drosophila* insulin receptor activates multiple signaling pathways but requires insulin receptor substrate proteins for DNA synthesis. *Mol. Cell. Biol.* **16**, 2509-2517.
- Zelzer, E., Levy, Y., Kahana, C., Shilo, B. Z., Rubinstein, M. and Cohen, B. (1998). Insulin induces transcription of target genes through the hypoxia-inducible factor HIF-1 $\alpha$ /ARNT. *EMBO J.* **17**, 5085-5094.
- Zhang, H., Stallock, J. P., Ng, J. C., Reinhard, C. and Neufeld, T. P. (2000). Regulation of cellular growth by the *Drosophila* target of rapamycin dTOR. *Genes Dev.* **14**, 2712-2724.
- Zhang, Y., Gao, X., Saucedo, L. J., Ru, B., Edgar, B. A. and Pan, D. (2003). Rheb is a direct target of the tuberous sclerosis tumour suppressor proteins. *Nat. Cell Biol.* **5**, 578-581.
- Zhong, H., Chiles, K., Feldser, D., Laughner, E., Hanrahan, C., Georgescu, M. M., Simons, J. W. and Semenza, G. L. (2000). Modulation of hypoxia-inducible factor 1 $\alpha$  expression by the epidermal growth factor/phosphatidylinositol 3-kinase/PTEN/AKT/FRAP pathway in human prostate cancer cells: implications for tumor angiogenesis and therapeutics. *Cancer Res.* **60**, 1541-1545.
- Zundel, W., Schindler, C., Haas-Kogan, D., Koong, A., Kaper, F., Chen, E., Gottschalk, A. R., Ryan, H. E., Johnson, R. S., Jefferson, A. B. et al. (2000). Loss of PTEN facilitates HIF-1-mediated gene expression. *Genes Dev.* **14**, 391-396.

## Statistical analysis of variations of HIF/Sima subcellular localization.

**Table 1. Variations in development.** Chi-square ( $\chi^2$ ) test to analyze developmental variations of HIF/Sima subcellular localization at different oxygen concentrations. Proportions of cytoplasmic, ubiquitous and nuclear categories of subcellular localization are compared between embryonic stages 11-12, 13-14, 15 and 16.

% O <sub>2</sub>	<i>P</i>	$\chi^2$ ; df=6	Statistically significant ( $\alpha=0.05$ )
21	<10 <sup>-4</sup>	87.3	+
5	<10 <sup>-4</sup>	88.3	+
3	<10 <sup>-4</sup>	124.9	+
1	<10 <sup>-4</sup>	38.7	+

$\chi^2$  values. *P*, P-value. df, degrees of freedom.

**Table 2. Effect of oxygen levels.** Chi-square ( $\chi^2$ ) test to analyze oxygen-dependent variations of HIF/Sima subcellular localization at different embryonic stages. Proportions of cytoplasmic, ubiquitous and nuclear categories of subcellular localization are compared between 21% O<sub>2</sub>, 5% O<sub>2</sub>, 3% O<sub>2</sub> and 1% O<sub>2</sub>.

Stages	<i>P</i>	$\chi^2$ ; df=6	Statistically significant ( $\alpha=0.05$ )
11-12	<10 <sup>-4</sup>	234.9	+
13-14	<10 <sup>-4</sup>	81.3	+
15	<10 <sup>-4</sup>	88.2	+
16	<10 <sup>-4</sup>	93.9	+

$\chi^2$ , chi-square values. *P*, P-value. df, degrees of freedom.

**Table 3. Effect of the over-expression of dAKT.** Chi-square ( $\chi^2$ ) test to analyze differences of HIF/Sima subcellular localization between wild type embryos and embryos over-expressing dAKT. Proportions of cytoplasmic, ubiquitous and nuclear categories of subcellular localization are compared at different embryonic stages.

Stages	<i>P</i>	$\chi^2$ ; df=2	Statistically significant ( $\alpha=0.05$ )
11-12	0.09	4.8	-
13-14	0.2	3.2	-
15	<10 <sup>-4</sup>	23.6	+
16	<10 <sup>-4</sup>	21.1	+

$\chi^2$ , chi-square values. *P*, P-value. df, degrees of freedom.

**Table 4: Effect of the overexpression of dAKT and dPDK1.** Chi-square ( $\chi^2$ ) test to analyze differences of HIF/Sima subcellular localization between embryos over-expressing dAKT and embryos over-expressing both dAKT and dPDK1. Proportions of cytoplasmic, ubiquitous and nuclear categories of subcellular localization are compared at different embryonic stages.

Stages	<i>P</i>	$\chi^2$ ; df=2	Statistically significant ( $\alpha=0.05$ )
11-12	0.08	5	-
13-14	<10 <sup>-4</sup>	35.6	+
15	0.01	7.9	+
16	0.3	2.4	-

$\chi^2$ , chi-square values. *P*, P-value. df, degrees of freedom.

**Table 5: Effect of PTEN loss of function.** Chi-square ( $\chi^2$ ) test to analyze differences of HIF/Sima subcellular localization between wild type embryos and  $PTEN^{2L117}$  homozygous mutant embryos. Proportions of cytoplasmic, ubiquitous and nuclear categories of subcellular localization are compared at different embryonic stages.

Stages	<i>P</i>	$\chi^2$ ; df=2	Statistically significant ( $\alpha=0.05$ )
11-12	0.07	5	-
13-14	$2 \cdot 10^{-4}$	16.7	+
15	$<10^{-4}$	24.6	+
16	$<10^{-4}$	30.7	+

$\chi^2$ , chi-square values. *P*, P-value. df, degrees of freedom.