

# Bre1p-mediated histone H2B ubiquitylation regulates apoptosis in *Saccharomyces cerevisiae*

David Walter, Anja Matter and Birthe Fahrenkrog\*‡

M. E. Müller Institute for Structural Biology, Biozentrum, University of Basel, CH-4056 Basel, Switzerland

\*Present address: Institute for Molecular Biology and Medicine, Université Libre de Bruxelles, 6041 Gosselies, Belgium

‡Author for correspondence (bfahrenk@ulb.ac.be)

Accepted 22 March 2010

Journal of Cell Science 123, 1931–1939

© 2010. Published by The Company of Biologists Ltd

doi:10.1242/jcs.065938

## Summary

*BRE1* encodes an E3 ubiquitin protein ligase that is required for the ubiquitylation of histone H2B at lysine 123 (K123). Ubiquitylation of this histone residue is involved in a variety of cellular processes including gene activation and gene silencing. Abolishing histone H2B ubiquitylation also confers X-ray sensitivity and abrogates checkpoint activation after DNA damage. Here we show that *Saccharomyces cerevisiae* Bre1p exhibits anti-apoptotic activity in yeast and that this is linked to histone H2B ubiquitylation. We found that enhanced levels of Bre1p protect from hydrogen-peroxide-induced cell death, whereas deletion of *BRE1* enhances cell death. Moreover, cells lacking Bre1p show reduced lifespan during chronological ageing, a physiological apoptotic condition in yeast. Importantly, the resistance against apoptosis is conferred by histone H2B ubiquitylation mediated by the E3 ligase activity of Bre1p. Furthermore, we found that the death of  $\Delta bre1$  cells depends on the yeast caspase Yca1p, because  $\Delta bre1$  cells exhibit increased caspase activity when compared with wild-type cells, and deletion of *YCA1* leads to reduced apoptosis sensitivity of cells lacking Bre1p.

**Key words:** Apoptosis, Histone H2B, *S. cerevisiae*, Ubiquitin, YCA1

## Introduction

Apoptosis is a form of programmed cell death that has a central role in development and cellular homeostasis in higher eukaryotes. Knowledge of apoptotic regulation is particularly important for medical research, since apoptotic misregulation is implicated in many human diseases, such as Alzheimer's disease and Huntington's disease, immunodeficiency and cancer (Fadeel and Orrenius, 2005). Recent studies have established yeast as a model to study the mechanisms of apoptotic regulation. Defects in distinct cellular processes, such as actin dynamics (Gourlay et al., 2004), vesicular fusion (Madeo et al., 1997), DNA replication (Weinberger et al., 2005), histone chaperone activity (Yamaki et al., 2001) or histone deubiquitylation (Bettiga et al., 2004) can trigger apoptotic cell death in *Saccharomyces cerevisiae* and an apoptotic-like phenotype has also been demonstrated in yeast cells treated with various agents including hydrogen peroxide, acetic acid and pheromone (Ludovico et al., 2001; Madeo et al., 1999; Severin and Hyman, 2002). Notably, the yeast apoptotic machinery has functional orthologues of key mammalian apoptotic regulators, including the metacaspase Yca1p (Madeo et al., 2002), the apoptosis-inducing factor AIF (Wissing et al., 2004), the endonuclease EndoG (Buttner et al., 2007), the serine protease HtrA2/Omi (Fahrenkrog et al., 2004) and the inhibitor-of-apoptosis protein Bir1p (Walter et al., 2006). In addition, yeast apoptosis has been linked to cellular events such as mitochondrial fragmentation (Fannjiang et al., 2004), cytochrome c release (Ludovico et al., 2002), ageing (Herker et al., 2004; Laun et al., 2001) and phosphorylation of histone H2B (Ahn et al., 2005; Ahn et al., 2006).

Rapid protein modifications allow the cell to promptly adapt to environmental changes by different cellular responses, including apoptosis. The post-translational modification by covalent attachment of ubiquitin is one of the major biochemical

mechanisms that regulate apoptosis (Lee and Peter, 2003). Ubiquitylation controls the level of proteins by targeting them for proteasomal degradation. Members of the inhibitor-of-apoptosis protein (IAP) family are targeted for degradation, but they also contain a RING domain with ubiquitylating activity, by which they are able to mark other proteins such as caspases for degradation (Wilson et al., 2002). Additionally, monoubiquitylation and nonclassical polyubiquitylation of components of the apoptotic pathway regulate apoptosis on a molecular level, beyond the degradation effects (Huang et al., 2000; Lee et al., 2002; Mimnaugh et al., 2001).

Histone proteins are well-known substrates for numerous covalent post-translational modifications and these modifications regulate a number of cellular processes including apoptosis (Ahn et al., 2005; Ahn et al., 2006; Cheung et al., 2003). Histone H2B is monoubiquitylated at Lys123 by the ubiquitin conjugase Rad6p and the E3 ligase Bre1p (Hwang et al., 2003; Robzyk et al., 2000; Wood et al., 2003b). *BRE1* disruption or lysine-to-arginine substitution at residue 123 of histone H2B (H2B-K123R), results in a complex phenotype that includes failures in gene activation (Henry et al., 2003; Kao et al., 2004; Wyce et al., 2007; Xiao et al., 2005) and lack of telomeric silencing (Briggs et al., 2001; Dover et al., 2002; Mutiu et al., 2007; Sun and Allis, 2002). Moreover, *Caenorhabditis elegans BRE1* was identified in a screen for anti-apoptotic proteins (Lettre et al., 2004). Only recently, histone H2B ubiquitylation has been implicated in DNA repair and checkpoint activation after DNA damage (Game et al., 2006; Giannattasio et al., 2005). As the DNA damage response machinery is closely linked to apoptosis in yeast and higher eukaryotes (Burhans et al., 2003) a relationship between histone H2B ubiquitylation and apoptosis might exist. This possibility is supported by the finding that loss of the ubiquitin-specific protease *UBP10*, which is involved in cleaving the ubiquitin moiety from

histone H2B, causes caspase activation and apoptosis in yeast (Bettiga et al., 2004).

We aimed to gain insight into the role of histone H2B ubiquitylation in apoptosis and found that *S. cerevisiae* Bre1p exhibits anti-apoptotic activity in yeast and that this is linked to histone H2B ubiquitylation and the metacaspase Yca1p.

## Results

### Bre1p protects against hydrogen-peroxide-induced cell death in budding yeast


The E3 ubiquitin ligase Bre1p is required for histone H2B ubiquitylation (Hwang et al., 2003; Robzyk et al., 2000; Wood et al., 2003b), which in turn is implicated in transcriptional regulation and DNA repair. Moreover, H2B ubiquitylation appears to have a role in apoptosis regulation because the loss of the ubiquitin-specific protease *UBP10*, which is involved in cleaving the ubiquitin moiety from histone H2B, causes caspase activation and apoptosis in yeast (Bettiga et al., 2004). Notably, the *C. elegans* homologue of the *S. cerevisiae* Bre1p was identified as a regulator of germ-cell apoptosis in worms (Lettre et al., 2004), further supporting the importance of H2B ubiquitylation in apoptosis regulation.

To explore a role for Bre1p-mediated histone H2B ubiquitylation in yeast apoptosis, wild-type cells, cells lacking *BRE1* ( $\Delta bre1$ ) and cells constitutively overexpressing a protein-A-tagged Bre1p fusion protein (ProtA-Bre1p) under the control of the *NOPI* promoter were exposed to 0.6 mM hydrogen peroxide ( $H_2O_2$ ) to induce apoptosis. After 8 hours of incubation, cell survival was determined by clonogenicity and cells were tested for apoptotic markers, such as DNA single-strand breaks and reactive oxygen species (ROS), which are causally linked to yeast apoptosis. To do so, cells were stained with dihydroethidium (DHE) to visualise accumulation of ROS and TUNEL labelling was used to detect single-stranded


DNA breaks. As shown in Fig. 1A, yeast cells lacking *BRE1* exhibited an increased sensitivity to  $H_2O_2$  ( $31\pm4\%$  cell viability) compared with wild-type (wt) cells ( $53\pm6\%$  cell viability). This increase in sensitivity to  $H_2O_2$  of  $\Delta bre1$  cells was accompanied by enhanced ROS production ( $54\pm4\%$  of  $\Delta bre1$  versus  $25\pm2\%$  of wild-type cells were DHE positive; Fig. 1B,C) and by an increase in apoptotic DNA fragmentation and TUNEL-positive cells compared with wild-type cells ( $25\pm3\%$  versus  $14\pm1\%$  TUNEL-positive cells; Fig. 1B,D). By contrast, cells overexpressing ProtA-Bre1p showed resistance to  $H_2O_2$  ( $78\pm7\%$  cell viability; Fig. 1A) and a decrease in ROS accumulation ( $15\pm1\%$  DHE-positive cells; Fig. 1B,C) and DNA fragmentation ( $5\pm1\%$  TUNEL-positive cells; Fig. 1B,D). Taken together, our data indicate that Bre1p exhibits anti-apoptotic activity.

### Disruption of *BRE1* causes an early onset of cell death during chronological ageing

Chronological ageing defines an ageing process of post-mitotic yeast cells that triggers apoptosis (Herker et al., 2004). Therefore, we next investigated whether or not Bre1p is involved in chronological ageing. To do so, we determined the chronological lifespan of cells lacking *BRE1* and found that these cells showed an early onset of age-induced cell death when compared with wild-type cells (Fig. 2A). After 2 days in culture, cells lacking *BRE1* showed survival rates of  $23\pm4\%$  compared with  $75\pm5\%$  in wild-type cells (Fig. 2A). When, after 2 days in culture, these yeast cells were tested for apoptotic markers,  $\Delta bre1$  cells showed typical hallmarks of apoptosis, such as the production of ROS as detected by DHE staining ( $77\pm3\%$  of  $\Delta bre1$  versus  $16\pm4\%$  of wild-type cells were DHE positive; Fig. 2B,C) and an increase in apoptotic DNA fragmentation as detected by TUNEL labelling ( $26\pm4\%$  of  $\Delta bre1$  versus  $6\pm1\%$  of wild-type cells were TUNEL positive; Fig.



**Fig. 1. *S. cerevisiae* Bre1p confers resistance to apoptosis induced by  $H_2O_2$ .** (A) Wild-type (wt) and  $\Delta bre1$  yeast cells harbouring the empty vector control, and ProtA-Bre1p-overexpressing cells were treated with 0.6 mM  $H_2O_2$  for 8 hours and survival was determined by clonogenicity. Data represent means  $\pm$  s.d. ( $n=6$ ; \*\*\* $P<0.001$ ). (B) ROS accumulation and DNA fragmentation in wild-type (wt) and  $\Delta bre1$  cells harbouring the vector control or overexpressing ProtA-Bre1p was determined by DHE staining and TUNEL staining, respectively. (C) DHE-positive and (D) TUNEL-positive cells were quantified by manually counting at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ).



**Fig. 2. Disruption of *BRE1* causes early onset of apoptosis during chronological ageing.** (A) Survival of wild-type (wt) and  $\Delta bre1$  cells determined by clonogenicity during chronological ageing. Data represent means  $\pm$  s.d. ( $n=9$ ). (B) ROS accumulation and DNA fragmentation in wild-type and  $\Delta bre1$  cells after 2 days in culture determined by DHE staining and TUNEL staining, respectively. (C) DHE-positive cells were quantified after 2 days in culture using flow cytometry. In each experiment, 10,000 cells were evaluated. Data represent means  $\pm$  s.d. ( $n=3$ ). (D) TUNEL-positive cells were quantified by manual counting of at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ).

2B,D). These data further support the notion that Bre1p acts as inhibitor of apoptosis in yeast. Constitutive overexpression of ProtA-Bre1p, however, did not significantly influence chronological lifespan of yeast cells (data not shown).

### The E3 ligase activity of Bre1p is required for its anti-apoptotic properties

A hallmark of Bre1p is a C-terminal C3HC4 (RING) zinc-finger domain (Hwang et al., 2003). RING domains are typically found in E3 ubiquitin ligases and frequently mediate the interaction with the E2 ubiquitin-activating enzyme (Deshaies and Joazeiro, 2009). The RING domains are therefore crucial for catalysing the transfer of ubiquitin from the E2 to the substrate. Accordingly, the RING domain of Bre1p confers E3 ubiquitin ligase activity, which is required for the ubiquitylation of histone H2B (Hwang et al., 2003; Wood et al., 2003a). To test whether the E3 ligase activity is required to grant resistance to age-induced apoptosis, we generated two different RING-domain mutants. First, two conserved cysteine residues (C648 and C651) within the RING domain were mutated to glycine and second, leucine L650 was mutated to glutamate (L650E). Although the mutation of the cysteines might affect the overall folding of the RING domain, L650 probably mediates the interaction of Bre1p with its E2 ligase, but does not affect the zinc coordination and stability of the protein. We next complemented  $\Delta bre1$  and  $\Delta bre1$  cells expressing H2B-GFP ( $\Delta bre1$  HTB1-GFP),

respectively, with plasmid-borne ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) and tested the functionality of the ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) as a measure of their ability to ubiquitylate histone H2B.


Monoubiquitylation of histone H2B can be detected in wild-type cells harbouring a functional GFP-tagged allele of the *HTB1* gene (encoding histone H2B) as a slower-migrating form upon SDS-PAGE and immunoblotting of whole-cell extracts with anti-GFP antibody. The ubiquitylated species was absent in  $\Delta bre1$  cells harbouring a GFP-tagged allele of *HTB1* (Fig. 3A), which is consistent with a previous study (Wood et al., 2003a). However,  $\Delta bre1$  HTB1-GFP cells complemented with ProtA-Bre1p displayed no defect in H2B monoubiquitylation, whereas  $\Delta bre1$  HTB1-GFP cells complemented with ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E), respectively, lacked ubiquitylation of histone H2B similarly to  $\Delta bre1$  HTB1-GFP cells (Fig. 3A). We conclude that the ProtA-Bre1p fusion protein is functional and that Bre1p requires the conserved cysteines C648, C651 and leucine 650 for its E3 ligase activity.

To explore the contribution of the E3 ligase activity of Bre1p to apoptosis resistance, we analysed the survival of  $\Delta bre1$  cells complemented with ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) during chronological ageing. *BRE1*-null cells complemented with ProtA-Bre1p showed no significant difference in cell survival when compared with wild-type cells ( $89 \pm 5\%$  cell viability versus  $92 \pm 5\%$  cell viability after 2 days in culture; Fig. 3A). Consistently,  $\Delta bre1$  cells complemented with ProtA-Bre1p and wild-type cells showed similar amounts of apoptotic markers with  $\sim 16\%$  DHE-positive cells and 6-7% TUNEL-positive cells, respectively (Fig. 3D-F). By contrast,  $\Delta bre1$  cells complemented with ProtA-Bre1p(C648G, C651G) or ProtA-Bre1p(L650E) showed an early onset of cell death during chronological ageing similarly to  $\Delta bre1$  cells ( $21 \pm 2\%$  and  $23 \pm 1\%$  cell viability versus  $23 \pm 4\%$  cell viability after 2 days in culture; Fig. 3B).  $79 \pm 6\%$  of  $\Delta bre1$  cells complemented with Bre1p(C648G, C651G) showed ROS accumulation and  $25 \pm 3\%$  were TUNEL positive, similarly to  $\Delta bre1$  cells ( $77 \pm 3\%$  ROS-positive,  $26 \pm 4\%$  TUNEL-positive cells; Fig. 3D-F).

Consistent with the chronological ageing experiments, plasmid-borne ProtA-Bre1p, but not ProtA-Bre1p(C648G, C651G) or ProtA-Bre1p(L650E) rescued  $\Delta bre1$  cells from  $H_2O_2$ -induced cell death (Fig. 3C).  $H_2O_2$ -treated cells lacking *BRE1* that were complemented with ProtA-Bre1p showed a slightly better cell survival than wild-type cells ( $67 \pm 5\%$  cell viability versus  $54 \pm 6\%$  cell viability; Fig. 3C) and fewer apoptotic markers ( $18 \pm 5\%$  versus  $25 \pm 2\%$  DHE-positive cells as well as  $8 \pm 3\%$  versus  $14 \pm 1\%$  TUNEL-positive cells; Fig. 3D,G,H). However,  $\Delta bre1$  cells complemented with ProtA-Bre1p(C648G, C651G) or ProtA-Bre1p(L650E) showed  $H_2O_2$  sensitivity that was similar to  $\Delta bre1$  cells ( $35 \pm 2\%$  and  $33 \pm 3\%$  cell viability versus  $31 \pm 4\%$  cell viability; Fig. 3C).  $48 \pm 3\%$  of  $\Delta bre1$  cells complemented with Bre1p(C648G, C651G) showed ROS accumulation and  $26 \pm 2\%$  were TUNEL positive, similarly to  $\Delta bre1$  cells ( $54 \pm 4\%$  ROS positive,  $25 \pm 3\%$  TUNEL positive; Fig. 3D,G,H). We conclude that the E3 ligase activity of Bre1p is required for its ability to confer resistance to apoptosis.

### Bre1p confers apoptosis resistance by histone H2B ubiquitylation

Bre1p targets Lys123 in histone H2B for ubiquitylation. We therefore asked whether the ubiquitylation site of histone H2B is




**Fig. 3. The E3 ligase activity of Bre1p is required for apoptosis inhibition.**

(A) Functionality of ProtA-Bre1p, ProtA-Bre1p(C648,651G) and ProtA-Bre1p(L650E) was tested as a measurement of the ability to ubiquitylate histone H2B. Wild-type (wt),  $\Delta bre1$  and  $\Delta bre1$  cells complemented with ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) and harbouring a functional GFP-tagged allele of the HTB1 gene (encoding histone H2B) were grown in synthetic complete medium (SC) overnight. Whole-cell lysates were separated on a 12% acrylamide gel and the blot was probed with GFP (Dianova, clone: MA1-26343; Hamburg, Germany) and protein-A antibody. (B) Survival of wild-type,  $\Delta bre1$ ,  $\Delta bre1$  cells complemented with ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) was determined by clonogenicity during chronological ageing. Data represent means  $\pm$  s.d. ( $n=9$ ). (C)  $\Delta bre1$  cells complemented with ProtA-Bre1p, ProtA-Bre1p(C648G, C651G) and ProtA-Bre1p(L650E) were treated with 0.6 mM H<sub>2</sub>O<sub>2</sub> for 8 hours and survival was determined by clonogenicity. Data represent means  $\pm$  s.d. ( $n=9$ ). (D) ROS accumulation and DNA fragmentation in  $\Delta bre1$  cells complemented with ProtA-Bre1p and ProtA-Bre1p(C648G, C651G) after 2 days in culture and after H<sub>2</sub>O<sub>2</sub> treatment determined by DHE staining and TUNEL staining, respectively. (E) DHE-positive cells during chronological ageing were quantified after 2 days in culture using flow cytometry. In each experiment, 10,000 cells were evaluated. Data represent means  $\pm$  s.d. ( $n=3$ ). (F) TUNEL-positive cells during chronological ageing were quantified after 2 days in culture by manual counting at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ). (G) DHE-positive cells were quantified after H<sub>2</sub>O<sub>2</sub> treatment by manually counting at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ). (H) TUNEL-positive cells were quantified after H<sub>2</sub>O<sub>2</sub> treatment by manually counting at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ).

required for the anti-apoptotic property of Bre1p. To address this question, we analysed the chronological lifespan of the yeast strain *FLAG-htb1K123R*, which expresses a FLAG-tagged histone H2B variant containing a lysine-to-arginine substitution at Lys123 and therefore fails to be ubiquitylated (Sun and Allis, 2002). We found that these cells showed an early onset of cell death during chronological ageing, similarly to  $\Delta bre1$  *FLAG-HTB1* cells that lack *BRE1* and express FLAG-tagged wild type histone H2B (36 $\pm$ 7% cell viability versus 32 $\pm$ 5% cell viability after 3 days in culture; Fig. 4A).

We next determined whether the lack of Bre1p and the lack of histone H2B ubiquitylation affect the same pathway leading to an early onset of cell death during chronological ageing. To do so, we disrupted *BRE1* in *FLAG-htb1K123R* cells and analysed the chronological lifespan of the resulting double mutant  $\Delta bre1$  *FLAG-htb1K123R*. An additive phenotype for the double mutant

is expected if the two mutations affect the chronological lifespan of yeast independently. However, the double mutant strain  $\Delta bre1$  *FLAG-htb1K123R* showed no further decrease in survival during chronological ageing when compared with either single mutant (35 $\pm$ 5% cell viability versus 36 $\pm$ 7% and 32 $\pm$ 5% cell viability, respectively, after 3 days in culture; Fig. 4A), indicating that both mutations affect the same pathway. Moreover, these data suggest that Bre1p confers resistance to age-induced apoptosis by mediating histone H2B ubiquitylation. Likewise, *FLAG-htb1-K123R* mutant cells showed H<sub>2</sub>O<sub>2</sub> sensitivity that was similar to that in  $\Delta bre1$  *FLAG-HTB1* cells (66 $\pm$ 7% cell viability versus 69 $\pm$ 7% cell viability; Fig. 4B), whereas the double mutant strain  $\Delta bre1$  *FLAG-htb1K123R* exhibited no further decrease in survival when compared with either single mutant (70 $\pm$ 6% cell viability versus 66 $\pm$ 7% and 68 $\pm$ 7% cell viability, respectively; Fig. 4B). Together, our data indicate that Bre1p




**Fig. 4. Histone H2B ubiquitylation confers apoptosis resistance.**

(A) Survival determined by clonogenicity of wild-type cells (*BRE1 FLAG-HTB1*) and its derivatives *BRE1 FLAG-htb1K123R*,  $\Delta$ *bre1 FLAG-HTB1* and  $\Delta$ *bre1 FLAG-htb1K123R* during chronological ageing. Data represent means  $\pm$  s.d. ( $n=6$ ). (B) The same strains were treated with 0.6 mM  $H_2O_2$  and survival was determined by clonogenicity. Data represent means  $\pm$  s.d. ( $n=6$ ; \*\*\* $P<0.001$ , \*\* $P<0.01$ ).

diminishes apoptotic death by mediating histone H2B ubiquitylation.

#### Death of $\Delta$ *bre1* cells depends on the yeast metacaspase Yca1p

Yeast cells lacking *BRE1* were sensitive to apoptotic stimuli and displayed morphological markers of apoptosis upon  $H_2O_2$  treatment and during chronological ageing (Figs 1 and 2). Apoptosis in yeast can occur in a caspase-dependent or caspase-independent manner (Madeo et al., 2009). To investigate whether the yeast metacaspase Yca1p is involved in the death of  $\Delta$ *bre1* cells, we generated a  $\Delta$ *yca1 $\Delta$ *bre1* double-mutant strain. The death rate during chronological ageing was decreased in  $\Delta$ *bre1 $\Delta$ *yca1* cells compared with  $\Delta$ *bre1* cells (47 $\pm$ 3% cell viability versus 21 $\pm$ 2% cell viability after 2 days in culture; Fig. 5A), indicating that Yca1p is required for cell death of  $\Delta$ *bre1* cells. However, the death of  $\Delta$ *bre1* cells is not exclusively Yca1p-dependent during chronological ageing as the double mutant  $\Delta$ *bre1 $\Delta$ *yca1* exhibited a higher death rate than wild-type cells (47 $\pm$ 3% cell viability versus 75 $\pm$ 5% cell viability after 2 days in culture; Fig. 5A). In addition,  $\Delta$ *bre1 $\Delta$ *yca1* cells displayed less ROS accumulation during chronological ageing when compared with  $\Delta$ *bre1* cells (41 $\pm$ 4% versus 77 $\pm$ 3% cells were DHE positive after 2 days in culture; Fig. 5B), but more than wild-type cells (41 $\pm$ 4% versus 15 $\pm$ 3% cells were DHE positive after 2 days in culture; Fig. 5B). Therefore, the death of  $\Delta$ *bre1* cells during chronological ageing is partially Yca1p dependent. Notably, unlike  $\Delta$ *bre1* cells,  $\Delta$ *bre1 $\Delta$ *yca1* cells did not display apoptotic DNA fragmentation during chronological ageing as detected by TUNEL labelling (supplementary material Fig. S1), suggesting that  $\Delta$ *bre1 $\Delta$ *yca1* cells die in a necrotic rather than apoptotic******




**Fig. 5. Death of  $\Delta$ *bre1* cells depends on the yeast caspase-like protein Yca1p.**

(A) Survival determined by clonogenicity during chronological ageing of wild-type (wt),  $\Delta$ *yca1*,  $\Delta$ *bre1* and  $\Delta$ *bre1 $\Delta$ *yca1* cells. Data represent means  $\pm$  s.d. ( $n=6$ ). (B) DHE-positive wild type (wt),  $\Delta$ *yca1*,  $\Delta$ *bre1* and  $\Delta$ *bre1 $\Delta$ *yca1* cells were quantified after 2 days in culture using flow cytometry. In each experiment, 10,000 cells were evaluated. Data represent means  $\pm$  s.d. ( $n=3$ ). (C) Wild-type (wt),  $\Delta$ *yca1*,  $\Delta$ *bre1* and  $\Delta$ *bre1 $\Delta$ *yca1* cells were exposed to 0.6 mM  $H_2O_2$  for 8 hours and survival was determined by clonogenicity. Data represent means  $\pm$  s.d. ( $n=6$ ). (D) DHE-positive cells were quantified after  $H_2O_2$  treatment by manually counting at least 500 cells. Data represent means  $\pm$  s.d. ( $n=3$ ).***

fashion. Annexin-V and propidium iodide (PI) costaining was used to further analyse apoptotic externalisation of phosphatidylserine and necrotic membrane permeabilisation.  $\Delta$ *bre1* cells, unlike  $\Delta$ *bre1 $\Delta$ *yca1* cells, displayed externalisation of phosphatidylserine as detected by annexin-V staining (supplementary material Fig. S1), further supporting the notion that  $\Delta$ *bre1* cells die in an apoptotic manner, whereas the death of  $\Delta$ *bre1 $\Delta$ *yca1* cells is of a necrotic nature.**

Next, we tested the response of  $\Delta$ *bre1 $\Delta$ *yca1* cells to  $H_2O_2$  exposure. As shown in Fig. 5C, the death rate after  $H_2O_2$  treatment was decreased in  $\Delta$ *bre1 $\Delta$ *yca1* cells when compared with  $\Delta$ *bre1* cells (57 $\pm$ 5% cell viability versus 32 $\pm$ 5% cell viability; Fig. 5C). Unlike during chronological ageing, the death of  $\Delta$ *bre1* cells seems to be exclusively Yca1p dependent after  $H_2O_2$  treatment, because the double-mutant  $\Delta$ *bre1 $\Delta$ *yca1* cells show survival rates similar to wild-type cells (57 $\pm$ 5% cell viability versus 50 $\pm$ 6% cell viability; Fig. 5C). In addition, the population of  $\Delta$ *bre1 $\Delta$ *yca1* cells with ROS accumulation after  $H_2O_2$  treatment was smaller when compared with  $\Delta$ *bre1* cells but similar to wild-type cells (23 $\pm$ 1% DHE-positive  $\Delta$ *bre1 $\Delta$ *yca1* cells compared with 53 $\pm$ 7% DHE-positive  $\Delta$ *bre1* and 23 $\pm$ 3% DHE-positive wild type cells; Fig. 5D). Taken together, our data indicate that Yca1p is required to activate apoptosis in  $\Delta$ *bre1* cells in response to  $H_2O_2$  treatment.*****

Apoptosis in yeast can also be Yca1p independent. The mitochondria-localised apoptosis-inducing factor Aif1p (Wissing



**Fig. 6. Cells lacking *BRE1* show increased caspase activity.** (A) Chronological aged wild-type and  $\Delta bre1$  cells and cells treated with 0.6 mM H<sub>2</sub>O<sub>2</sub> were labelled with FITC-VAD-fmk and propidium iodide (PI) and analysed by flow cytometry. One representative experiment from three independent experiments with similar results is shown. (B) Chronological aged wild-type and  $\Delta bre1$  cells and cells treated with 0.6 mM H<sub>2</sub>O<sub>2</sub> were incubated with the caspase substrate (L-Asp)<sub>2</sub> rhodamine 110 (D<sub>2</sub>R) and analysed by flow cytometry. One representative experiment from three independent experiments with similar results is shown.

et al., 2004), the endonuclease Nuc1p/EndoG (Buttner et al., 2007) and possibly the nuclear serine protease Nma111p (Fahrenkrog et al., 2004) can execute caspase-independent apoptosis. To test whether these pro-apoptotic factors are also involved in the Bre1p pathway, we generated distinct double-mutant strains,  $\Delta bre1\Delta nuc1$ ,  $\Delta bre1\Delta ai1$  and  $\Delta bre1\Delta nma111$ . As shown in supplementary material Fig. S2, none of these strains displayed better survival during chronological ageing or upon H<sub>2</sub>O<sub>2</sub> treatment when compared with  $\Delta bre1$  cells, indicating that these other pro-apoptotic factors do not contribute to the Bre1p pathway.

#### Cells lacking *BRE1* show increased caspase activity

Since disruption of *YCA1* desensitises  $\Delta bre1$  cells towards apoptotic stimuli, we asked whether or not the caspase activity of Yca1p is involved in cell death execution of  $\Delta bre1$  cells. To address this question, we tested whether  $\Delta bre1$  cells exhibited higher caspase activity after induction of apoptosis compared with wild-type cells. To monitor potential caspase activation, yeast cells were incubated with FITC-labelled VAD-fmk (FITC-VAD-fmk). FITC-VAD-fmk binds specifically to the active centre of metazoan caspases, which enables a flow-cytometric determination of cells with active caspases (Madedo et al., 2002). As FITC-VAD-fmk might have the limitation that it stains dead cells nonspecifically (Wysocki and Kron, 2004), we additionally used propidium iodide (PI) to distinguish between apoptotic (PI negative) and necrotic cells (PI positive). Wild-type and *BRE1*-disrupted cells were compared after stimulation with 0.6 mM H<sub>2</sub>O<sub>2</sub> or after 1 day of chronological ageing. As shown in Fig. 6A, after treatment with 0.6 mM H<sub>2</sub>O<sub>2</sub> 34% of  $\Delta bre1$  cells showed caspase activity (FITC positive, PI negative) (corresponding FACS profiles are found in supplementary material Fig. S3), whereas only 23% of wild-type cells exhibited caspase activity. Consistently, we monitored caspase activity in

about 21% of aged  $\Delta bre1$  cells, but only in 13% of wild-type cells (Fig. 6A). To confirm these findings, we used the caspase substrate (L-Asp)<sub>2</sub> rhodamine 110 (D<sub>2</sub>R), which is designated for the detection of caspase activity in mammalian cells. D<sub>2</sub>R is non-fluorescent, however, upon cleavage by a caspase, the released rhodamine 110 gives rise to a fluorescence signal, which enables a flow cytometric determination of caspase activity in cells. As shown in Fig. 6B, after 1 day of chronological ageing and after treatment with 0.6 mM H<sub>2</sub>O<sub>2</sub>,  $\Delta bre1$  cells exhibited higher caspase activity than wild-type cells. Taken together, our data show that  $\Delta bre1$  cells have higher caspase activity than wild-type cells, further supporting the notion that Bre1p acts in a Yca1p-dependent manner.

#### Discussion

*BRE1* encodes an evolutionarily conserved E3 ubiquitin ligase that, in yeast, catalyses monoubiquitylation of histone H2B at lysine 123 (K123). Histone H2B K123 ubiquitylation is involved in a variety of cellular processes, such as gene activation, gene silencing and checkpoint activation after DNA damage (Briggs et al., 2001; Dover et al., 2002; Game et al., 2006; Giannattasio et al., 2005; Henry et al., 2003; Kao et al., 2004; Mutiu et al., 2007; Sun and Allis, 2002; Wyce et al., 2007; Xiao et al., 2005). Here, we uncover a new role for Bre1p and show that *S. cerevisiae* Bre1p protects yeast cells from H<sub>2</sub>O<sub>2</sub>-induced cell death, whereas deletion of *BRE1* enhances cell death and leads to decreased lifespan during chronological ageing. Also, we show that Bre1p activity in yeast apoptosis requires its E3 ubiquitin ligase activity, thereby linking yeast apoptosis to histone H2B monoubiquitylation. Furthermore, we show that Bre1p protects yeast cells from death in a Yca1p-dependent manner.

#### A role for Bre1p in programmed cell death

Enhanced levels of Bre1p protect yeast from H<sub>2</sub>O<sub>2</sub>-induced cell death and diminish the development of apoptotic hallmarks, such as ROS accumulation and DNA single-strand breaks (Fig. 1). By contrast, cells lacking Bre1p are more sensitive to H<sub>2</sub>O<sub>2</sub> treatment (Fig. 1) and show a decreased lifespan during chronological ageing (Fig. 2A), which coincided with the appearance of apoptotic markers (Fig. 2B-D). A role for Bre1p in yeast cell death has not been assumed, but is consistent with the recent identification of its *C. elegans* homologue as a regulator of germ-cell apoptosis in worms (Lettre et al., 2004). Therefore, the anti-apoptotic function of Bre1p is probably evolutionary conserved and it will be interesting to see whether, for example, the human homologues of Bre1p, RNF20 and RNF40, are also implicated in apoptosis regulation.

The ability of Bre1p to reduce cell death is conferred by its E3 ubiquitin ligase activity and histone H2B ubiquitylation (Figs 3 and 4). Bre1p harbours a C-terminal zinc-binding motif known as the RING-finger domain (Hwang et al., 2003) that is frequently found in E3 ubiquitin ligases and is required for catalysing the transfer of ubiquitin from the E2 to the substrate (Deshaies and Joazeiro, 2009). RING domains appear to be crucial for apoptosis regulation, because members of the inhibitor-of-apoptosis protein (IAP) family comprise RING domains, which enable IAPs to mark other proteins such as caspases for proteasomal degradation (Wilson et al., 2002). Our data indicate that, in addition to polyubiquitylation, monoubiquitylation might have a role in apoptosis. We show that  $\Delta bre1$  cells complemented with the RING-finger mutants ProtA-Bre1p(C648G, C651G) and ProtA-

Bre1p(L650E) lack monoubiquitylated histone H2B (Fig. 3A) and exhibit increased apoptosis sensitivity similarly to *Δbre1* cells, whereas *Δbre1* cells complemented with a functional ProtA-Bre1p behave in a similar manner to wild-type cells (Fig. 3). These findings suggest that Bre1p requires its E3 ligase activity to confer H2B monoubiquitylation and apoptosis resistance and support the notion that RING domains have a major role in apoptosis regulation.

The importance of H2B monoubiquitylation for the anti-apoptotic activity of Bre1p is further supported by our observation that *htb1-K123R* mutant cells, which fail to ubiquitylate histone H2B, also exhibit increased apoptosis sensitivity, similarly to *Δbre1* cells (Fig. 4A,B). Interestingly, yeast cells that exhibit enhanced levels of ubiquitylated histone H2B owing to the lack of the ubiquitin-specific protease Ubp10p, which cleaves the ubiquitin moiety from histone H2B, are also prone to apoptosis (Bettiga et al., 2004; Emre et al., 2005; Gardner et al., 2005). These data indicate that high levels of histone H2B ubiquitylation can also predispose yeast to apoptotic stimuli in a similar manner to the lack of histone H2B ubiquitylation. However, this study did not address whether or not *Δubp10* cells show increased caspase activity due to high ubiquitylation levels of histone H2B. It is therefore possible that Ubp10p has other targets than ubiquitylated H2B and failure in deubiquitylation of these targets might cause sensitivity to apoptosis in these cells. To rule out this possibility, we analysed the apoptosis sensitivity of *Δbre1Δubp10*. We expected that this double mutant would exhibit increased apoptosis sensitivity when compared with *Δbre1* cells, if Ubp10p acted in a histone-H2B-independent manner. However, *Δbre1* and *Δbre1Δubp10* cells showed similar apoptosis sensitivity during chronological ageing (supplementary material Fig. S4A). After 2 days in culture, *BRE1*-null cells showed survival rates of 23±4% compared with 24±2% in *Δbre1Δubp10* cells (supplementary material Fig. S4A). In addition, *Δbre1Δubp10* and *Δbre1* cells showed similar survival rates after H<sub>2</sub>O<sub>2</sub> treatment [31±5% cell viability versus 28±3% cell viability (supplementary material Fig. S4B)]. These data suggest that *UBP10* disruption causes apoptosis sensitivity because of failures in histone H2B deubiquitylation. Therefore, the lack of histone H2B ubiquitylation, as well as high levels of histone H2B ubiquitylation, appear to predispose yeast to apoptotic stimuli, indicating that H2B monoubiquitylation needs to be tightly regulated to ensure cell survival.

The yeast metacaspase Yca1p appears to be essential for approximately 40% of the investigated cell-death scenarios in yeast (Madoe et al., 2009) and we show here that apoptosis in *Δbre1* cells in fact depends on Yca1p, because *YCA1* disruption leads to reduced apoptosis sensitivity in cells lacking Bre1p (Fig. 5). Furthermore, the death of *Δbre1* cells depends neither on other pro-apoptotic factors, such as EndoG, Nma111p nor on Aif1p (supplementary material Fig. S2), indicating that Bre1p anti-apoptotic activity is, at least in part, caspase dependent. Consistently, we show that *Δbre1* cells exhibit increased caspase activity compared with wild-type cells (Fig. 6), suggesting that the caspase-like activity of Yca1p is implicated in the death of *Δbre1* cells.

### Possible mechanisms

Our data suggest that lack of histone H2B monoubiquitylation leads to Yca1p-dependent apoptosis during chronological ageing and after H<sub>2</sub>O<sub>2</sub> treatment. Histone H2B ubiquitylation has a role in the DNA-damage response, as well as in transcriptional control (Briggs et al., 2001; Dover et al., 2002; Game et al., 2006;

Giannattasio et al., 2005; Henry et al., 2003; Kao et al., 2004; Mutiu et al., 2007; Sun and Allis, 2002; Wyce et al., 2007; Xiao et al., 2005). Therefore, cells lacking histone H2B ubiquitylation might undergo Yca1p-dependent apoptosis, either because of accumulated DNA damage, or because of alterations in transcription. Although H2B ubiquitylation has been implicated in transcriptional silencing (Briggs et al., 2001; Mutiu et al., 2007; Sun and Allis, 2002), other studies suggested a positive role for this modification in transcriptional initiation and elongation (Henry et al., 2003; Kao et al., 2004; Wyce et al., 2007; Xiao et al., 2005). Notably, the core apoptotic machinery, including caspases and other regulators of apoptosis, are regulated at the transcriptional level in higher eukaryotes (Zuckerman et al., 2009). Therefore, it is possible that this balance is disturbed in cells lacking H2B ubiquitylation and up- or down-regulated transcription of apoptotic regulators causes apoptosis of these cells. It would be interesting to see whether *YCA1* or other pro-apoptotic proteins are transcriptionally deregulated in *Δbre1* cells. Furthermore, comparison of global mRNA transcripts between wild-type and *Δbre1* cells could help to identify novel regulators of apoptosis.

Histone H2B ubiquitylation is required for Rad9p-mediated checkpoint activation after DNA damage and Rad51p-dependent DNA repair (Game et al., 2006; Giannattasio et al., 2005). Therefore, cells lacking histone H2B ubiquitylation might undergo Yca1p-dependent apoptosis because of accumulated DNA damage. However, neither *Δrad9* nor *Δrad51* cells exhibit apoptosis sensitivity similar to that observed in *Δbre1* cells (our unpublished results), indicating that apoptosis in *Δbre1* is not caused by defects in Rad9p or Rad51p pathways. Moreover, although *Δbre1* cells exhibit sensitivity towards DNA damage induced by methyl methanesulfonate, hydroxyurea and UV radiation, respectively, disruption of Yca1p does not lead to a rescue in survival of *Δbre1* cells under these conditions (data not shown). Therefore, we do not consider it likely that DNA damage causes Yca1p-dependent apoptosis in *Δbre1* cells.

In conclusion, we show that Bre1p confers resistance to apoptosis and Yca1p is required in the apoptosis pathway triggered by *BRE1* disruption. Bre1p is required for histone H2B ubiquitylation and its deletion, which influences transcriptional regulation and DNA repair, activates apoptosis. However, it remains to be seen whether transcription defects, failures in DNA repair, or both processes, activate the apoptotic program in cells lacking H2B ubiquitylation. Future studies in yeast will provide more details on the connection between transcription, DNA repair and apoptosis.

### Materials and Methods

#### Plasmids, yeast strains and culture conditions

To construct the plasmid pBF326, which encodes ProtA-Bre1p, the coding region of *BRE1* was amplified from genomic DNA isolated from BY4742 cells, using the following primers: 5'-CAT GCC ATG GCA ATG ACG GCC GAG CCT GCT A-3' and 5'-CGC GGA TCC TTA CAA GTG CAC TGT CAA TAA ATC-3'. The PCR product was digested with *NcoI* and *BamHI* and cloned into pNOPATA1L (Hellmuth et al., 1998). Plasmids pBF346 and pBF499 coding for ProtA-Bre1p(C648G,C651G) and ProtA-Bre1p(L650E), respectively, were constructed by site-directed polymerase chain reaction (PCR) mutagenesis using pBF326 as template according to the manufacturer's instructions (Stratagene, QuikChange Site-Directed Mutagenesis Kit).

Yeast strains are listed in supplementary material Table S1. BY4742 (MAT $\alpha$ ; his3 $\Delta$ 1; leu2 $\Delta$ 0; lys2 $\Delta$ 0; ura3 $\Delta$ 0) and derivative strain *Δbre1* were obtained from Euroscarf. *Δubp10* strain was derived from BY4742 and constructed according to Longtine et al. (Longtine et al., 1998). Construction of the *Δbre1Δyca1*, *Δbre1ΔendoG*, *Δbre1Δnma111*, *Δbre1Δaif1* and *Δbre1Δubp10* double-mutant strains was performed according to published methods (Gueldener et al., 2002). Yeast strains YZS276 (*FLAG-HTB1*) and YZS277 (*FLAG-hb1K123R*) were a gift from C. David Allis (Rockefeller University, New York, NY) (Sun and Allis, 2002a).

*Δbre1 FLAG-HTB1* and *Δbre1 FLAG-htb1K123R* were derived from YZS276 and YZS277, respectively and constructed as published (Longtine et al., 1998). *HTB1-GFP Δbre1* was derived from *Δbre1* strain and constructed as described (Longtine et al., 1998). Survival plating was conducted on YPAD (1% yeast extract, 2% peptone, and 2% glucose, 40 mg/ml Adenine) medium supplemented with 2% agar. For experiments testing oxygen stress and chronological lifespan, strains were grown in synthetic complete medium (SC) with 2% glucose (Fink, 1991). Transformation of yeast cells was performed by the lithium acetate procedure, as described (Gietz et al., 1992).

#### Survival plating and test for apoptotic markers

For experiments testing oxygen stress, cultures were inoculated at low cell density ( $2 \times 10^5$  cells/ml) in SCglu, grown to late log phase ( $OD_{600} \sim 2$ ) and exposed to 0.6 mM hydrogen peroxide ( $H_2O_2$ ) for 8 hours. For survival plating, yeast cultures were diluted in water, the cell concentration was determined using a Neubauer counting chamber and aliquots containing 500 cells were plated on YPAD plates. The number of colonies was determined after incubation for 2 days at 30°C. Percentage of cell survival was calculated for each strain by counting the number of colonies formed following  $H_2O_2$  treatment relative to untreated cells. Apoptotic tests using DHE staining, annexin-V and PI staining and TUNEL staining were performed as described previously (Belanger et al., 2009; Buttner et al., 2007). In each sample, 10,000 cells were evaluated using flow cytometry (FACS-Aria, BD) and processed using BD FACSDiva software. Alternatively, around 500 DHE- and TUNEL-stained cells, respectively, were counted manually. For chronological ageing experiments, cultures were inoculated from fresh overnight cultures at low cell density ( $1 \times 10^6$  cells/ml) and aliquots were taken to perform survival plating and tests for apoptotic markers as described above.

#### In vivo staining of caspase activity by flow cytometry

$5 \times 10^6$  cells were harvested, washed once in 1 ml PBS and incubated in PBS containing 10 μM FITC-VAD-fmk (CaspACE, Promega, Dübendorf) for 20 minutes at 30°C in the dark. Next, the cells were washed with PBS and resuspended in PBS containing 1 μg/ml propidium iodide (PI) and analysed by flow cytometry (FACS-Aria, BD). Cleavage of the caspase substrate (asparyl)-Rhodamine 110 ( $D_2R$ ) [CaspSCREEN (BioVision)] was measured by flow cytometry (FACS-Aria, BD) according to the manufacturer's instructions.

#### Immunoblotting

Protein lysates were prepared using a lysis buffer containing 100 mM NaCl, 50 mM Tris-HCl pH 7.5, 50 mM NaF, 5 mM EDTA and 0.1% IGEPAL. Lysate proteins were resolved on 12% acrylamide-Tris-HCl gels and transferred to an Immobilon PVDF membrane (Millipore, Billerica, MA). The filters were hybridised with anti-GFP antibody (Dianova, clone MA1-26343; Hamburg, Germany) and anti-protein-A antibody (Sigma), respectively and the peroxidase-conjugated secondary anti-mouse antibody (Sigma).

We thank C. D. Allis for providing plasmids YZS276 and YZS277 and Yvonne Lussi for critically reading the manuscript. This work was supported by grants from the Swiss National Science Foundation (to B.F.) as well as by the Kanton Basel Stadt and the M. E. Müller Foundation.

Supplementary material available online at

<http://jcs.biologists.org/cgi/content/full/123/11/1931/DC1>

#### References

- Ahn, S. H., Cheung, W. L., Hsu, J. Y., Diaz, R. L., Smith, M. M. and Allis, C. D. (2005). Sterile 20 kinase phosphorylates histone H2B at serine 10 during hydrogen peroxide-induced apoptosis in *S. cerevisiae*. *Cell* **120**, 25-36.
- Ahn, S. H., Diaz, R. L., Grunstein, M. and Allis, C. D. (2006). Histone H2B deacetylation at lysine 11 is required for yeast apoptosis induced by phosphorylation of H2B at serine 10. *Mol. Cell* **24**, 211-220.
- Belanger, K. D., Walter, D., Henderson, T. A., Yelton, A. L., O'Brien, T. G., Belanger, K. G., Geier, S. J. and Fahrenkrog, B. (2009). Nuclear localisation is crucial for the proapoptotic activity of the HtrA-like serine protease Nma111p. *J. Cell Sci.* **122**, 3931-3941.
- Bettiga, M., Calzari, L., Orlandi, I., Alberghina, L. and Vai, M. (2004). Involvement of the yeast metacaspase Yca1 in *ubpΔ*-programmed cell death. *FEMS Yeast Res.* **5**, 141-147.
- Briggs, S. D., Bryk, M., Strahl, B. D., Cheung, W. L., Davie, J. K., Dent, S. Y. R., Winston, F. and Allis, C. D. (2001). Histone H3 lysine 4 methylation is mediated by Set1 and required for cell growth and rDNA silencing in *Saccharomyces cerevisiae*. *Genes Dev.* **15**, 3286-3295.
- Burhans, W. C., Weinberger, M., Marchetti, M. A., Ramachandran, L., D'Urso, G. and Huberman, J. A. (2003). Apoptosis-like yeast cell death in response to DNA damage and replication defects. *Mutat. Res.* **532**, 227-243.
- Buttner, S., Eisenberg, T., Carmona-Gutierrez, D., Ruli, D., Knauer, H., Ruckentuhl, C., Sigrist, C., Wissing, S., Kollroser, M., Frohlich, K. U. et al. (2007). Endonuclease G regulates budding yeast life and death. *Mol. Cell* **25**, 233-246.
- Cheung, W. L., Ajiro, K., Samejima, K., Kloc, M., Cheung, P., Mizzen, C. A., Beeser, A., Etkin, L. D., Chernoff, J., Earnshaw, W. C. et al. (2003). Apoptotic phosphorylation of histone H2B is mediated by mammalian sterile twenty kinase. *Cell* **113**, 507-517.
- Deshaies, R. J. and Joazeiro, C. A. (2009). RING domain E3 ubiquitin ligases. *Annu. Rev. Biochem.* **78**, 399-434.
- Dover, J., Schneider, J., Tawiah-Boateng, M. A., Wood, A., Dean, K., Johnston, M. and Shilatifard, A. (2002). Methylation of histone H3 by COMPASS requires ubiquitination of histone H2B by Rad6. *J. Biol. Chem.* **277**, 28368-28371.
- Emre, N. C., Ingvarsdottir, K., Wyce, A., Wood, A., Krogan, N. J., Henry, K. W., Li, K., Marmorstein, R., Greenblatt, J. F., Shilatifard, A. et al. (2005). Maintenance of low histone ubiquitylation by Ubp10 correlates with telomere-proximal Sir2 association and gene silencing. *Mol. Cell* **17**, 585-594.
- Fadeel, B. and Orrenius, S. (2005). Apoptosis: a basic biological phenomenon with wide-ranging implications in human disease. *J. Intern. Med.* **258**, 479-517.
- Fahrenkrog, B., Sauder, U. and Aebi, U. (2004). The *S. cerevisiae* HtrA-like protein Nma111p is a nuclear serine protease that mediates yeast apoptosis. *J. Cell Sci.* **117**, 115-126.
- Fannjiang, Y., Cheng, W.-C., Lee, S. J., Qi, B., Pevsner, J., McCaffery, J. M., Hill, R. B., Basanez, G. and Hardwick, J. M. (2004). Mitochondrial fission proteins regulate programmed cell death in yeast. *Genes Dev.* **18**, 2785-2797.
- Fink, G. A. (1991). Guide to yeast genetics and molecular biology. *Methods Enzymol.* **194**, 1-863.
- Game, J. C., Williamson, M. S., Spicakova, T. and Brown, M. (2006). The RAD6/BRE1 histone modification pathway in *Saccharomyces* confers radiation resistance through a RAD51-dependent process that is independent of RAD18. *Genetics* **173**, 1951-1968.
- Gardner, R. G., Nelson, Z. W. and Gottschling, D. E. (2005). Ubp10/Dot4p regulates the persistence of ubiquitinated histone H2B: distinct roles in telomeric silencing and general chromatin. *Mol. Cell Biol.* **25**, 6123-6139.
- Giannattasio, M., Lazzaro, F., Plevani, P. and Muzi-Falconi, M. (2005). The DNA damage checkpoint response requires histone H2B ubiquitination by Rad6-Bre1 and H3 methylation by Dot1. *J. Biol. Chem.* **280**, 9879-9886.
- Gietz, D., St Jean, A., Woods, R. A. and Schiestl, R. H. (1992). Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**, 1425.
- Gourlay, C. W., Carpp, L. N., Timpson, P., Winder, S. J. and Ayscough, K. R. (2004). A role for the actin cytoskeleton in cell death and aging in yeast. *J. Cell Biol.* **164**, 803-809.
- Gueldener, U., Heinisch, J., Koehler, G. J., Voss, D. and Hegemann, J. H. (2002). A second set of loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. *Nucleic Acids Res.* **30**, e23.
- Hellmuth, K., Lau, D. M., Bischoff, F. R., Kunzler, M., Hurt, E. and Simos, G. (1998). Yeast Los1p has properties of an exportin-like nucleocytoplasmic transport factor for tRNA. *Mol. Cell Biol.* **18**, 6374-6386.
- Henry, K. W., Wyce, A., Lo, W. S., Duggan, L. J., Emre, N. C. T., Kao, C. F., Pillus, L., Shilatifard, A., Osley, M. A. and Berger, S. L. (2003). Transcriptional activation via sequential histone H2B ubiquitylation and deubiquitylation, mediated by SAGA-associated Ubp8. *Genes Dev.* **17**, 2648-2663.
- Herker, E., Jungwirth, H., Lehmann, K. A., Maldener, C., Frohlich, K. U., Wissing, S., Buttner, S., Fehr, M., Sigrist, S. and Madeo, F. (2004). Chronological aging leads to apoptosis in yeast. *J. Cell Biol.* **164**, 501-507.
- Huang, H.-K., Joazeiro, C. A. P., Bonfoco, E., Kamada, S., Levenson, J. D. and Hunter, T. (2000). The Inhibitor of Apoptosis, cIAP2, Functions as a Ubiquitin-Protein Ligase and Promotes in Vitro Monoubiquitination of Caspases 3 and 7. *J. Biol. Chem.* **275**, 26661-26664.
- Hwang, W. W., Venkatasubrahmanyam, S., Ianculescu, A. G., Tong, A., Boone, C. and Madhani, H. D. (2003). A conserved RING finger protein required for histone H2B monoubiquitination and cell size control. *Mol. Cell* **11**, 261-266.
- Kao, C. F., Hillyer, C., Tsukuda, T., Henry, K., Berger, S. and Osley, M. A. (2004). Rad6 plays a role in transcriptional activation through ubiquitylation of histone H2B. *Genes Dev.* **18**, 184-195.
- Laun, P., Pichova, A., Madeo, F., Fuchs, J., Ellinger, A., Kohlwein, S., Dawes, I., Frohlich, K. U. and Breitenbach, M. (2001). Aged mother cells of *Saccharomyces cerevisiae* show markers of oxidative stress and apoptosis. *Mol. Microbiol.* **39**, 1166-1173.
- Lee, J. C. and Peter, M. E. (2003). Regulation of apoptosis by ubiquitination. *Immunol. Rev.* **193**, 39-47.
- Lee, J. C., Schickling, O., Stegh, A. H., Oshima, R. G., Dinsdale, D., Cohen, G. M. and Peter, M. E. (2002). DEDD regulates degradation of intermediate filaments during apoptosis. *J. Cell Biol.* **158**, 1051-1066.
- Lettre, G., Kritikou, E. A., Jaeggi, M., Calixto, A., Fraser, A. G., Kamath, R. S., Ahringer, J. and Hengartner, M. O. (2004). Genome-wide RNAi identifies p53-dependent and -independent regulators of germ cell apoptosis in *C. elegans*. *Cell Death Differ.* **11**, 1198-1203.
- Longtine, M. S., McKenzie, A., 3rd, Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., Philippsen, P. and Pringle, J. R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* **14**, 953-961.
- Ludovico, P., Sousa, M. J., Silva, M. T., Leao, C. and Corte-Real, M. (2001). *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. *Microbiology* **147**, 2409-2415.
- Ludovico, P., Rodrigues, F., Almeida, A., Silva, M. T., Barrientos, A. and Corte-Real, M. (2002). Cytochrome c release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**, 2598-2606.
- Madeo, F., Frohlich, E. and Frohlich, K.-U. (1997). A yeast mutant showing diagnostic markers of early and late apoptosis. *J. Cell Biol.* **139**, 729-734.

- Madeo, F., Fröhlich, E., Ligr, M., Grey, M., Sigrist, S. J., Wolf, D. H. and Fröhlich, K. U. (1999). Oxygen stress: a regulator of apoptosis in yeast. *J. Cell Biol.* **145**, 757-767.
- Madeo, F., Herker, E., Maldener, C., Wissing, S., Lachelt, S., Herlan, M., Fehr, M., Lauber, K., Sigrist, S. J., Wesselborg, S. et al. (2002). A caspase-related protease regulates apoptosis in yeast. *Mol. Cell* **9**, 911-917.
- Madeo, F., Carmona-Gutierrez, D., Ring, J., Buttner, S., Eisenberg, T. and Kroemer, G. (2009). Caspase-dependent and caspase-independent cell death pathways in yeast. *Biochem. Biophys. Res. Commun.* **382**, 227-231.
- Mimnaugh, E. G., Kayastha, G., McGovern, N. B., Hwang, S. G., Marcu, M. G., Trepel, J., Cai, S. Y., Marchesi, V. T. and Neckers, L. (2001). Caspase-dependent deubiquitination of monoubiquitinated nucleosomal histone H2A induced by diverse apoptogenic stimuli. *Cell Death Differ.* **8**, 1182-1196.
- Mutiu, A. I., Hoke, S. M., Genereaux, J., Liang, G. and Brandl, C. J. (2007). The role of histone ubiquitylation and deubiquitylation in gene expression as determined by the analysis of an HTB1(K123R) *Saccharomyces cerevisiae* strain. *Mol. Genet. Genomics* **277**, 491-506.
- Robzyk, K., Recht, J. and Osley, M. A. (2000). Rad6-dependent ubiquitination of histone H2B in yeast. *Science* **287**, 501-504.
- Severin, F. F. and Hyman, A. A. (2002). Pheromone induces programmed cell death in *S. cerevisiae*. *Curr. Biol.* **12**, R233-R235.
- Sun, Z. W. and Allis, C. D. (2002). Ubiquitination of histone H2B regulates H3 methylation and gene silencing in yeast. *Nature* **418**, 104-108.
- Walter, D., Wissing, S., Madeo, F. and Fahrenkrog, B. (2006). The inhibitor-of-apoptosis protein Bir1p protects against apoptosis in *S. cerevisiae* and is a substrate for the yeast homologue of Omi/HtrA2. *J. Cell Sci.* **119**, 1843-1851.
- Weinberger, M., Ramachandran, L., Feng, L., Sharma, K., Sun, X., Marchetti, M., Huberman, J. A. and Burhans, W. C. (2005). Apoptosis in budding yeast caused by defects in initiation of DNA replication. *J. Cell Sci.* **118**, 3543-3553.
- Wilson, R., Goyal, L., Ditzel, M., Zachariou, A., Baker, D. A., Agapite, J., Steller, H. and Meier, P. (2002). The DIAP1 RING finger mediates ubiquitination of Dronc and is indispensable for regulating apoptosis. *Nat. Cell Biol.* **4**, 445-450.
- Wissing, S., Ludovico, P., Herker, E., Buttner, S., Engelhardt, S. M., Decker, T., Link, A., Proksch, A., Rodrigues, F., Corte-Real, M. et al. (2004). An AIF orthologue regulates apoptosis in yeast. *J. Cell Biol.* **166**, 969-974.
- Wood, A., Krogan, N. J., Dover, J., Schneider, J., Heidt, J., Boateng, M. A., Dean, K., Golshani, A., Zhang, Y., Greenblatt, J. F. et al. (2003a). Bre1, an E3 ubiquitin ligase required for recruitment and substrate selection of Rad6 at a promoter. *Mol. Cell* **11**, 267-274.
- Wood, A., Schneider, J., Dover, J., Johnston, M. and Shilatifard, A. (2003b). The Paf1 complex is essential for histone monoubiquitination by the Rad6-Bre1 complex, which signals for histone methylation by COMPASS and Dot1p. *J. Biol. Chem.* **278**, 34739-34742.
- Wyce, A., Xiao, T., Whelan, K. A., Kosman, C., Walter, W., Eick, D., Hughes, T. R., Krogan, N. J., Strahl, B. D. and Berger, S. L. (2007). H2B ubiquitylation acts as a barrier to Ctk1 nucleosomal recruitment prior to removal by Ubp8 within a SAGA-related complex. *Mol. Cell* **27**, 275-288.
- Wysocki, R. and Kron, S. J. (2004). Yeast cell death during DNA damage arrest is independent of caspase or reactive oxygen species. *J. Cell Biol.* **166**, 311-316.
- Xiao, T., Kao, C. F., Krogan, N. J., Sun, Z. W., Greenblatt, J. F., Osley, M. A. and Strahl, B. D. (2005). Histone H2B ubiquitylation is associated with elongating RNA polymerase II. *Mol. Cell Biol.* **25**, 637-651.
- Yamaki, M., Umehara, T., Chimura, T. and Horikoshi, M. (2001). Cell death with predominant apoptotic features in *Saccharomyces cerevisiae* mediated by deletion of the histone chaperone ASF1/CIA1. *Genes Cells* **6**, 1043-1054.
- Zuckerman, V., Wolyniec, K., Sionov, R. V., Haupt, S. and Haupt, Y. (2009). Tumour suppression by p53: the importance of apoptosis and cellular senescence. *J. Pathol.* **219**, 3-15.

**Table S1. Yeast strains used in this study**

<b>Yeast strain</b>	<b>Genotype</b>	<b>Source</b>
BY4742	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i>	<i>Euroscarf</i>
BFY177	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i>	<i>Euroscarf</i>
BFY324	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::LEU</i> , <i>YOR197W::kanMX4</i>	This study
BFY402	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YJL208C::kanMX4</i> , <i>YDL074C::LEU</i>	This study
BFY323	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YNL123W::kanMX4</i> , <i>YDL074C::LEU</i>	This study
BFY403	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YNR074C::kanMX4</i> , <i>YDL074C::LEU</i>	This study
YZS276	<i>MATa</i> , <i>ura3 1</i> , <i>trp1 1</i> , <i>leu2 3</i> , <i>his3 11</i> , <i>ade2 1</i> , <i>can1 100</i> , <i>hta1-</i> <i>htb1 ::LEU2 hta2-htb2 &lt;pZS145</i> <i>HTA1-Flag-HTB1,CEN, HIS3&gt;</i>	( <i>Sun and Allis,</i> 2002a)
YZS277	<i>MATa</i> , <i>ura3 1</i> , <i>trp1 1</i> , <i>leu2 3</i> , <i>his3 11</i> , <i>ade2 1</i> , <i>can1 100</i> , <i>hta1-</i> <i>htb1 ::LEU2 hta2-htb2 &lt;pZS146</i> <i>HTA1-Flag-htb1-K123R, CEN, HIS3&gt;</i>	( <i>Sun and Allis,</i> 2002a)
BFY508	<i>MATa</i> , <i>ura3 1</i> , <i>trp1 1</i> , <i>leu2 3</i> , <i>his3 11</i> , <i>ade2 1</i> , <i>can1 100</i> , <i>hta1-</i> <i>htb1 ::LEU2 hta2-htb2</i> , <i>YDL074c::kanMX4 &lt;pZS146 HTA1-</i> <i>Flag-htb1-K123R, CEN, HIS3&gt;</i>	This study
BFY507	<i>MATa</i> , <i>ura3 1</i> , <i>trp1 1</i> , <i>leu2 3</i> , <i>his3 11</i> , <i>ade2 1</i> , <i>can1 100</i> , <i>hta1-</i> <i>htb1 ::LEU2 hta2-htb2</i> , <i>YDL074c::kanMX4 &lt;pZS146 HTA1-</i> <i>Flag-htb1-K123R, CEN, HIS3&gt;</i>	This study
BFY191	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> ; <i>YDL074c::kanMX4</i> , <i>YDR224C:: YDR224C-GFP-HIS3-MX6</i> <i>&lt;PBF316 pNOPPATA1L-BRE1&gt;</i>	This study
BFY190	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4 &lt;PBF316</i>	This study

---

	<i>pNOPPATAIL-BRE1</i> >	
BFY192	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> < <i>PBF346</i> <i>pNOPPATAIL-bre1C648G,C651G</i> >	<i>This study</i>
BFY512	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> < <i>pBF499</i> <i>pNOPPATAIL-bre1L650E</i> >	<i>This study</i>
BFY193	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> , <i>YDR224C::</i> <i>YDR224C-GFP-HIS3-MX6</i> < <i>PBF346</i> <i>pNOPPATAIL-bre1C648G,C651G</i> >	<i>This study</i>
BFY510	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> , <i>YDR224C::</i> <i>YDR224C-GFP-HIS3-MX6</i> < <i>pBF499</i> <i>pNOPPATAIL-bre1L650E</i> >	<i>This study</i>
BFY199	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> < <i>PBF316 pNOPPATAIL-BRE1</i> >	<i>This study</i>
BFY200	<i>MAT<math>\alpha</math></i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> < <i>PBF005</i> <i>pNOPPATAIL</i> >	<i>This study</i>
BFY201	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> < <i>PBF005 pNOPPATAIL</i> >	<i>This study</i>

---

BFY514	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YNL186W::LEU</i>	<i>This study</i>
BFY515	<i>MAT_</i> , <i>his3 1</i> , <i>leu2 0</i> , <i>lys2 0</i> , <i>ura3 0</i> , <i>YDL074c::kanMX4</i> , <i>YNL186W::LEU</i>	<i>This study</i>

---