



Cover: NuSAP is essential for mitotic progression in early embryos, but not for spindle checkpoint activity. *Nusap1*-null mouse blastocysts display increased staining of the mitotic marker H3-P (pink) after treatment with the spindle poison nocodazole, indicative of a functional checkpoint. DNA (purple), F-actin (green). See article by A. Vanden Bosch et al. (pp. 3244–3255).

- Cell Science at a Glance**
- 3209 **Regulation of the Apaf-1–caspase-9 apoptosome.** Bratton, S. B. and Salvesen, G. S.
- Research Articles**
- 3215 **Golgi-associated GSK3 β regulates the sorting process of post-Golgi membrane trafficking.** Adachi, A., Kano, F., Tsuboi, T., Fujita, M., Maeda, Y. and Murata, M.
- 3226 **The exocytic gene *secA* is required for *Dictyostelium* cell motility and osmoregulation.** Zanchi, R., Howard, G., Bretscher, M. S. and Kay, R. R.
- 3235 **The recruitment of acetylated and unacetylated tropomyosin to distinct actin polymers permits the discrete regulation of specific myosins in fission yeast.** Coulton, A. T., East, D. A., Galinska-Rakoczy, A., Lehman, W. and Mulvihill, D. P.
- 3244 **NuSAP is essential for chromatin-induced spindle formation during early embryogenesis.** Vanden Bosch, A., Raemaekers, T., Denayer, S., Torrekens, S., Smets, N., Moermans, K., Dewerchin, M., Carmeliet, P. and Carmeliet, G.
- 3256 **The overexpression of the putative gut stem cell marker *Musashi-1* induces tumorigenesis through Wnt and Notch activation.** Rezza, A., Skah, S., Roche, C., Nadjar, J., Samarut, J. and Plateroti, M.
- 3266 **The Na⁺/H⁺ exchanger *Nhx1p* regulates the initiation of *Saccharomyces cerevisiae* vacuole fusion.** Qiu, Q.-S. and Fratti, R. A.
- 3276 **Arg206 of SNAP-25 is essential for neuroexocytosis at the *Drosophila melanogaster* neuromuscular junction.** Megighian, A., Scorzeto, M., Zanini, D., Pantano, S., Rigoni, M., Benna, C., Rossetto, O., Montecucco, C. and Zordan, M.
- 3284 **The role of *Schizosaccharomyces pombe dma1* in spore formation during meiosis.** Krapp, A., Cano del Rosario, E. and Simanis, V.
- 3294 **Regulation of bile canalicular network formation and maintenance by AMP-activated protein kinase and LKB1.** Fu, D., Wakabayashi, Y., Ido, Y., Lippincott-Schwartz, J. and Arias, I. M.
- 3303 **The nonsense-mediated decay pathway maintains synapse architecture and synaptic vesicle cycle efficacy.** Long, A. A., Mahapatra, C. T., Woodruff, E. A., III, Rohrbough, J., Leung, H.-T., Shino, S., An, L., Doerge, R. W., Metzstein, M. M., Pak, W. L. and Broadie, K.
- 3316 **PKC ζ regulates cell polarisation and proliferation restriction during mammary acinus formation.** Whyte, J., Thornton, L., McNally, S., McCarthy, S., Lanigan, F., Gallagher, W. M., Stein, T. and Martin, F.
- 3329 **Transmembrane domains control exclusion of membrane proteins from clathrin-coated pits.** Mercanti, V., Marchetti, A., Lelong, E., Perez, F., Orci, L. and Cosson, P.
- 3336 **Pro-cathepsin D interacts with the extracellular domain of the β chain of LRP1 and promotes LRP1-dependent fibroblast outgrowth.** Beaujoui, M., Prébois, C., Derocq, D., Laurent-Matha, V., Masson, O., Pattingre, S., Coopman, P., Bettache, N., Grossfield, J., Hollingsworth, R. E., Zhang, H., Yao, Z., Hyman, B. T., van der Geer, P., Smith, G. K. and Liaudet-Coopman, E.
- 3347 **Decrease of sialic acid residues as an *eat-me* signal on the surface of apoptotic lymphocytes.** Meesmann, H. M., Fehr, E.-M., Kierschke, S., Herrmann, M., Bilyy, R., Heyder, P., Blank, N., Krienke, S., Lorenz, H.-M. and Schiller, M.
- 3357 **WLS-dependent secretion of WNT3A requires Ser209 acylation and vacuolar acidification.** Coombs, G. S., Yu, J., Canning, C. A., Veltri, C. A., Covey, T. M., Cheong, J. K., Utomo, V., Banerjee, N., Zhang, Z. H., Juldco, R. C., Concepcion, G. P., Bugni, T. S., Harper, M. K., Mihalek, I., Jones, C. M., Ireland, C. M. and Virshup, D. M.
- 3368 **Src-dependent phosphorylation of ROCK participates in regulation of focal adhesion dynamics.** Lee, H.-H., Tien, S.-C., Jou, T.-S., Chang, Y.-C., Jhong, J.-G. and Chang, Z.-F.
- 3378 **AT-1 is the ER membrane acetyl-CoA transporter and is essential for cell viability.** Jonas, M. C., Pehar, M. and Puglielli, L.
- 3389 **PEX11 family members are membrane elongation factors that coordinate peroxisome proliferation and maintenance.** Koch, J., Pranjic, K., Huber, A., Ellinger, A., Hartig, A., Kragler, F. and Brocard, C.
- 3401 **Developmental regulation and extracellular release of a *VSG* expression-site-associated gene product from *Trypanosoma brucei* bloodstream forms.** Barnwell, E. M., van Deursen, F. J., Jeacock, L., Smith, K. A., Maizels, R. M., Acosta-Serrano, A. and Matthews, K.
- Erratum**
- 3412 **Protein mobilities and P-selectin storage in Weibel–Palade bodies.** Kiskin, N. I., Hellen, N., Babich, V., Hewlett, L., Knipe, L., Hannah, M. J. and Carter, T.
- Author Correction**
- 3413 **ERK5 is required for VEGF-mediated survival and tubular morphogenesis of primary human microvascular endothelial cells.** Roberts, O. Ll., Holmes, K., Müller, J., Cross, D. A. E. and Cross, M. J.