


Cover: The image shows co-expression of the neuronal SNARE protein syntaxin-1-GFP (green) and the adaptor protein Munc18-1 (red) in PC12 cells deficient in endogenous Munc18. A loop in Munc18-1 is crucial for neurosecretion but not syntaxin-1 trafficking. See article by S. Martin et al. (pp. 2353–2360).

Volume 126 (11) June 1, 2013

Sticky Wicket

2315 **The end of science II – facing forward.** Mole

Commentaries

2319  **Microtubule-based transport – basic mechanisms, traffic rules and role in neurological pathogenesis.** Franker, M. A. M. and Hoogenraad, C. C.

2331 **Using fly genetics to dissect the cytoskeletal machinery of neurons during axonal growth and maintenance.** Prokop, A., Beaven, R., Qu, Y. and Sánchez-Soriano, N.

2343 **An immunologist's guide to CD31 function in T-cells.** Marelli-Berg, F. M., Clement, M., Mauro, C. and Caligiuri, G.

Research Articles

2353 **The Munc18-1 domain 3a loop is essential for neuroexocytosis but not for syntaxin-1A transport to the plasma membrane.** Martin, S., Tomatis, V. M., Papadopoulos, A., Christie, M. P., Malintan, N. T., Gormal, R. S., Sugita, S., Martin, J. L., Collins, B. M. and Meunier, F. A.

2361 **Domain 3a of Munc18-1 plays a crucial role at the priming stage of exocytosis.** Han, G. A., Bin, N.-R., Kang, S.-Y. A., Han, L. and Sugita, S.

2372 **BBS7 is required for BBSome formation and its absence in mice results in Bardet-Biedl syndrome phenotypes and selective abnormalities in membrane protein trafficking.** Zhang, Q., Nishimura, D., Vogel, T., Shao, J., Swiderski, R., Yin, T., Searby, C., Carter, C. S., Kim, G., Bugge, K., Stone, E. M. and Sheffield, V. C.

2381 **ERK2 but not ERK1 mediates HGF-induced motility in non-small cell lung carcinoma cell lines.** Radtke, S., Milanovic, M., Rossé, C., De Rycker, M., Lachmann, S., Hibbert, A., Kermorgant, S. and Parker, P. J.

2392 **The kinesin-like protein TOP promotes Aurora localisation and induces mitochondrial, chloroplast and nuclear division.** Yoshida, Y., Fujiwara, T., Imoto, Y., Yoshida, M., Ohnuma, M., Hirooka, S., Misumi, O., Kuroiwa, H., Kato, S., Matsunaga, S. and Kuroiwa, T.

2401 **Intramolecular shielding maintains the ER Ca²⁺ sensor STIM1 in an inactive conformation.** Yu, F., Sun, L., Hubrack, S., Selvaraj, S. and Machaca, K.

2411 **CIP4 coordinates with phospholipids and actin-associated proteins to localize to the protruding edge and produce actin ribs and veils.** Saengsawang, W., Taylor, K. L., Lumbard, D. C., Mitok, K., Price, A., Pietila, L., Gomez, T. M. and Dent, E. W.

2424 **Rab35 establishes the EHD1-association site by coordinating two distinct effectors during neurite outgrowth.** Kobayashi, H. and Fukuda, M.

2436 **Negative regulation of p53 by Ras superfamily protein RBEL1A.** Lui, K., An, J., Montalbano, J., Shi, J., Corcoran, C., He, Q., Sun, H., Sheikh, M. S. and Huang, Y.

2446 **A novel role of microRNA146b in promoting mammary alveolar progenitor cell maintenance.** Elsarraj, H. S., Hong, Y., Valdez, K., Carletti, M., Salah, S. M., Raimo, M., Taverna, D., Prochasson, P., Bharadwaj, U., Twardy, D. J., Christenson, L. K. and Behbod, F.

2459 **Netrin-1-induced activation of Notch signaling mediates glioblastoma cell invasion.** Ylivinkka, I., Hu, Y., Chen, P., Rantanen, V., Hautaniemi, S., Nyman, T. A., Keski-Oja, J. and Hyytiäinen, M.

2470 **The *Drosophila* Arf GEF Steppke controls MAPK activation in EGFR signaling.** Hahn, I., Fuss, B., Peters, A., Werner, T., Sieberg, A., Gosejacob, D. and Hoch, M.

2480 **FE65 regulates and interacts with the Bloom syndrome protein in dynamic nuclear spheres – potential relevance to Alzheimer's disease.** Schrötter, A., Mastalski, T., Nensa, F. M., Neumann, M., Loosse, C., Pfeiffer, K., Magraoui, F. E., Platta, H. W., Erdmann, R., Theiss, C., Uszkoreit, J., Eisenacher, M., Meyer, H. E., Marcus, K. and Müller, T.

2493 **Microtubule motors mediate endosomal sorting by maintaining functional domain organization.** Hunt, S. D., Townley, A. K., Danson, C. M., Cullen, P. J. and Stephens, D. J.

2502 **The genetic depletion or the triptolide inhibition of TFIIF in p53-deficient cells induces a JNK-dependent cell death in *Drosophila*.** Villicaña, C., Cruz, G. and Zurita, M.

2516 **Conversion of genomic imprinting by reprogramming and redifferentiation.** Kim, M. J., Choi, H. W., Jang, H. J., Chung, H. M., Arauzo-Bravo, M. J., Schöler, H. R. and Do, J. T.

2525 **During post-natal human myogenesis, normal myotube size requires TRPC1- and TRPC4-mediated Ca²⁺ entry.** Antigny, F., Koenig, S., Bernheim, L. and Frieden, M.

2534 **Fine mapping of autophagy-related proteins during autophagosome formation in *Saccharomyces cerevisiae*.** Suzuki, K., Akioka, M., Kondo-Kakuta, C., Yamamoto, H. and Ohsumi, Y.