

Contents

VOLUME 92 PART 4 APRIL 1989

| | | |
|---|-----|-----|
| Book Reviews | 529 | |
| Verheijen, R., Kuijpers, H. J. H., van Driel, R., Beck, J. L. M., van Dierendonck, J. H., Brakenhoff, G. J. and Ramaekers, F. C. S. | | |
| Ki-67 detects a nuclear matrix-associated proliferation-related antigen. II. Localization in mitotic cells and association with chromosomes | 531 | |
| Traas, J. P., Burgain, S. and Dumas de Vaulx, R. | | |
| The organization of the cytoskeleton during meiosis in eggplant (<i>Solanum melongena</i> (L.): microtubules and F-actin are both necessary for coordinated meiotic division | 541 | |
| Slominski, A., Moellmann, G. and Kuklinska, E. | | |
| MSH inhibits growth in a line of amelanotic hamster melanoma cells and induces increases in cyclic AMP levels and tyrosinase activity without inducing melanogenesis | 551 | |
| Tienari, J., Virtanen, I. and Lehtonen, E. | | |
| Lectin binding of F9 embryonal carcinoma cells: evidence for population heterogeneity and developmentally regulated high- M_r cell surface proteins | 561 | |
| Tang, X., Hepler, P. K. and Scordilis, S. P. | | |
| Immunochemical and immunocytochemical identification of a myosin heavy chain polypeptide in <i>Nicotiana</i> pollen tubes | 569 | |
| Wang, H., Cutler, A. J. and Fowke, L. C. | | |
| High frequencies of preprophase bands in soybean protoplast cultures | 575 | |
| Cho, S.-O. and Wick, S. M. | | |
| Microtubule orientation during stomatal differentiation in grasses | 581 | |
| Prescott, A. R., Foster, K. E., Warn, R. M. and Gull, K. | | |
| Incorporation of tubulin from an evolutionarily diverse source, <i>Physarum polycephalum</i> , into the microtubules of a mammalian cell | 595 | |
| Díaz-Nido, J. and Avila, J. | | |
| Characterization of proteins immunologically related to brain microtubule-associated protein MAP-1B in non-neural cells | 607 | |
| Hollenbeck, P. J., Bershadsky, A. D., Pletjushkina, O. Y., Tint, I. S. and Vasiliev, J. M. | | |
| Intermediate filament collapse is an ATP-dependent and actin-dependent process | 621 | |
| Burkhardt, J. K. and Argon, Y. | | |
| Intracellular transport of the glycoprotein of VSV is inhibited by CCCP at a late stage of post-translational processing | | 633 |
| Burkhardt, J. K., Hester, S. and Argon, Y. | | |
| The glycoprotein of VSV accumulates in a distal Golgi compartment in the presence of CCCP | | 643 |
| Hunt, R. C., Dewey, A. and Davis, A. A. | | |
| Transferrin receptors on the surfaces of retinal pigment epithelial cells are associated with the cytoskeleton | | 655 |
| Skerrow, C. J., Clelland, D. G. and Skerrow, D. | | |
| Changes to desmosomal antigens and lectin-binding sites during differentiation in normal human epidermis: a quantitative ultrastructural study | | 667 |
| Bouziges, F., Simon-Assmann, P., Leberquier, C., Haffen, K. and Kedinger, M. | | |
| Glycosaminoglycan expression in intestinal epithelial skin-fibroblastic cell cocultures. Fibroblastic cell-mediated effects of glucocorticoids | | 679 |
| Harris, A. and Coleman, L. | | |
| Ductal epithelial cells cultured from human foetal epididymis and vas deferens: relevance to sterility in cystic fibrosis | | 687 |
| Dluzewski, A. R., Fryer, P. R., Griffiths, S., Wilson, R. J. M. and Gratzer, W. B. | | |
| Red cell membrane protein distribution during malarial invasion | | 691 |
| Bonifacino, J. S., Yuan, L. and Sandoval, I. V. | | |
| Internalization and recycling to serotonin-containing granules of the 80K integral membrane protein exposed on the surface of secreting rat basophilic leukaemia cells | | 701 |
| Vazquez, M. H., Phillips, D. M. and Wassarman, P. M. | | |
| Interaction of mouse sperm with purified sperm receptors covalently linked to silica beads | | 713 |
| Grant, S. G. and Worton, R. G. | | |
| Activation of the <i>hprt</i> gene on the inactive X chromosome in transformed diploid female Chinese hamster cells | | 723 |
| Index of Authors | | 733 |
| Index of Subjects | | 737 |
| List of Supplements | | |